



United States Department of Agriculture

---

# **Alcohol Consumption and All-Cause Mortality: A Systematic Review**

---

2020 Dietary Guidelines Advisory Committee,  
Beverages and Added Sugars Subcommittee

Published date: July 15, 2020

Nutrition Evidence Systematic Review  
Center for Nutrition Policy and Promotion  
Food and Nutrition Service  
U.S. Department of Agriculture  
Braddock Metro Center II  
1320 Braddock Place  
Alexandria, Virginia 22314

This systematic review was conducted by the 2020 Dietary Guidelines Advisory Committee in collaboration with the Nutrition Evidence Systematic Review (NESR) team at the Center for Nutrition Policy and Promotion, Food and Nutrition Service, U.S. Department of Agriculture (USDA). All systematic reviews from the 2020 Advisory Committee Project are available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews>.

Conclusion statements drawn as part of this systematic review describe the state of science related to the specific question examined. Conclusion statements do not draw implications, and should not be interpreted as dietary guidance. This portfolio provides the complete documentation for this systematic review. A summary of this review is included in the 2020 Advisory Committee's Scientific Report available at [www.DietaryGuidelines.gov](http://www.DietaryGuidelines.gov).

The contents of this document may be used and reprinted without permission. Endorsements by NESR, the Center for Nutrition Policy and Promotion, the Food and Nutrition Service, or the USDA of derivative products developed from this work may not be stated or implied.

**Suggested citation for this systematic review:** 2020 Dietary Guidelines Advisory Committee and Nutrition Evidence Systematic Review Team. Alcohol Consumption and All-Cause Mortality: A Systematic Review. 2020 Dietary Guidelines Advisory Committee Project. Alexandria, VA: U.S. Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy and Promotion, July 2020. Available at: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews>.

**Related citation:** Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.

In accordance with Federal civil rights law and USDA civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer, and lender.

## ACKNOWLEDGEMENTS

---

### **Beverages and Added Sugars Subcommittee:**

- Elizabeth Mayer-Davis, PhD, RD, University of North Carolina at Chapel Hill, Subcommittee Chair
- Heather Leidy, PhD, University of Texas at Austin
- Richard Mattes, PhD, MPH, RD, Purdue University
- Timothy Naimi, MD, MPH, Boston University
- Rachel Novotny, PhD, RDN, LD, University of Hawaii
- Barbara Schneeman, PhD, University of California, Davis, Chair of the 2020 Dietary Guidelines Advisory Committee

### **Nutrition Evidence Systematic Review (NESR) Team:**

- Brittany James Kingshipp, PhD, Analyst, Panum Group<sup>i</sup>
- Maureen Spill, PhD, Analyst, Panum Group<sup>i</sup>
- Natasha Chong Cole, PhD, MPH, RD, Analyst, Panum Group<sup>i</sup>
- Gisela Butera, MLIS, MEd, Systematic Review Librarian, Panum Group<sup>i</sup>
- Nancy Terry, MS, MLS, Biomedical Librarian, National Institutes of Health Library, U.S. Department of Health and Human Services (HHS)
- Julie Obbagy, PhD, RD, Project Lead, Office of Nutrition Guidance and Analysis (ONGA), Center for Nutrition Policy and Promotion (CNPP), Food and Nutrition Service (FNS), U.S. Department of Agriculture (USDA)

### **Federal Liaisons:**

- Jennifer Seymour, PhD, Division of Nutrition, Physical Activity, and Obesity, Centers for Disease Control and Prevention, HHS
- Julia Quam, MSPH, RDN, Division of Prevention Science (DPS), Office of Disease Prevention and Health Promotion (ODPHP), HHS
- Clarissa (Claire) Brown, MS, MPH, RD, ONGA, CNPP, FNS, USDA
- Meghan Adler, MS, RD, FADA, ONGA, CNPP, FNS, USDA

### **Project Leadership:**

- Eve Essery Stooddy, PhD, Designated Federal Officer and Director, ONGA, CNPP, FNS, USDA
- Janet de Jesus, MS, RD, Nutrition Advisor, DPS, ODPHP, Office of the Assistant Secretary for Health (OASH), HHS
- Richard (Rick) Olson, MD, MPH, Director, DPS, ODPHP, OASH, HHS

USDA and HHS implemented a process to identify topics and scientific questions to be examined by the 2020 Dietary Guidelines Advisory Committee. The Committee conducted its review of evidence in subcommittees for discussion by the full Committee during its public meetings. The role of the Committee members involved establishing all aspects of the protocol, which presented the plan for how they would examine the scientific evidence, including the inclusion and exclusion criteria;

---

<sup>i</sup> Under contract with the Food and Nutrition Service, United States Department of Agriculture.

reviewing all studies that met the criteria they set; deliberating on the body of evidence for each question; and writing and grading the conclusion statements to be included in the scientific report the 2020 Committee submitted to USDA and HHS. The NESR team with assistance from Federal Liaisons and Project Leadership, supported the Committee by facilitating, executing, and documenting the work necessary to ensure the reviews were completed in accordance with NESR methodology. More information about the 2020 Dietary Guidelines Advisory Committee, including the process used to identify topics and questions, can be found at [www.DietaryGuidelines.gov](http://www.DietaryGuidelines.gov). More information about NESR can be found at [NESR.usda.gov](http://NESR.usda.gov).

The Committee and NESR staff thank USDA's Agricultural Research Service for coordinating the peer review of this systematic review, and the Federal scientist peer reviewers for their time and expertise.

**FUNDING SOURCE:** United States Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy and Promotion, Alexandria, VA

# TABLE OF CONTENTS

Acknowledgements.....	3
Table of contents .....	5
Introduction .....	6
What is the relationship between alcohol consumption and all-cause mortality? .....	8
Plain language summary.....	8
Technical abstract .....	10
Full review .....	12
Systematic review question .....	12
Conclusion statements and grades .....	12
Summary of the evidence .....	12
Description of the evidence .....	13
Evidence synthesis .....	16
Research recommendations.....	21
Included articles.....	22
Methodology .....	85
Analytic framework.....	86
Literature search and screening plan .....	87
Inclusion and exclusion criteria .....	87
Electronic databases and search terms.....	89
Literature search and screening results .....	91
Excluded articles .....	92
Table 1: Summary of articles that address the relationship between alcohol consumption and all-cause mortality.....	27
Table 2: Risk of bias for observational studies examining alcohol consumption and all-cause mortality.....	83
Table 3: Inclusion and exclusion criteria .....	87
Table 4: Articles excluded after full text screening with rationale for exclusion.....	92
Figure 1: Analytic framework .....	86
Figure 2: Flow chart of literature search and screening results.....	91

## INTRODUCTION

---

This document describes a systematic review conducted to answer the following question: What is the relationship between alcohol consumption and all-cause mortality? This systematic review was conducted by the 2020 Dietary Guidelines Advisory Committee, supported by USDA's Nutrition Evidence Systematic Review (NESR).

More information about the 2020 Dietary Guidelines Advisory Committee is available at the following website: [www.DietaryGuidelines.gov](http://www.DietaryGuidelines.gov).

NESR specializes in conducting food- and nutrition-related systematic reviews using a rigorous, protocol-driven methodology. More information about NESR is available at the following website: [NESR.usda.gov](http://NESR.usda.gov).

NESR's systematic review methodology involves developing a protocol, searching for and selecting studies, extracting data from and assessing the risk of bias of each included study, synthesizing the evidence, developing conclusion statements, grading the evidence underlying the conclusion statements, and recommending future research. A detailed description of the systematic reviews conducted for the 2020 Dietary Guidelines Advisory Committee, including information about methodology, is available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews>. In addition, starting on page 85, this document describes the final protocol as it was applied in the systematic review. A description of and rationale for modifications made to the protocol are described in the 2020 Dietary Guidelines Advisory Committee Report, Part D: Chapter 11. Alcoholic Beverages.

## List of abbreviations

Abbreviation	Full name
ACM	All-cause mortality
CNPP	Center for Nutrition Policy and Promotion
CVD	Cardiovascular disease
DPS	Division of Prevention Science
FNS	Food and Nutrition Service
HDI	Human development index
HHS	United States Department of Health and Human Services
NESR	Nutrition Evidence Systematic Review
OASH	Office of the Assistant Secretary for Health
ODPHP	Office of Disease Prevention and Health Promotion
ONGA	Office of Nutrition Guidance and Analysis
PCS	Prospective cohort study
RCT	Randomized controlled trial
UK	United Kingdom
USDA	United States Department of Agriculture

# WHAT IS THE RELATIONSHIP BETWEEN ALCOHOL CONSUMPTION AND ALL-CAUSE MORTALITY?

---

## PLAIN LANGUAGE SUMMARY

### What is the question?

- The question is: What is the relationship between alcohol consumption and all-cause mortality?

### What is the answer to the question?

- **Primary comparisons (among those who currently drink alcohol)**
  - Moderate evidence indicates that higher average alcohol consumption is associated with an increased risk of all-cause mortality compared with lower average alcohol consumption among those who drink.
  - Moderate evidence indicates that binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) is associated with increased risk of all-cause mortality, and that more frequent binge drinking is associated with increased risk of all-cause mortality compared with less frequent or no binge drinking among those who drink.
- **Secondary comparison (between those who currently drink alcohol and those who have never consumed alcohol)**
  - Limited evidence suggests that low average alcohol consumption, particularly without binge drinking, is associated with a lower risk of all-cause mortality compared with never drinking alcohol. However, in light of the many scientific and public health issues associated with alcoholic beverages, any conclusions about low average consumption compared to never drinking alcohol require careful consideration.

### Why was this question asked?

- This important public health question was identified by the U.S. Departments of Agriculture (USDA) and Health and Human Services (HHS) to be examined by the 2020 Dietary Guidelines Advisory Committee.

### How was this question answered?

- The 2020 Dietary Guidelines Advisory Committee, Beverages and Added Sugars Subcommittee conducted a systematic review to answer this question with support from the Nutrition Evidence Systematic Review (NESR) team.

### What is the population of interest?

- The evidence includes alcohol consumption and all-cause mortality in primarily adults ages 21 years and older.

### What evidence was found?

- This review includes 60 articles.
- The review is divided into 2 parts: evidence that compares people who drink



different amounts of alcohol, and evidence that compares people who drink alcohol with people who never drank alcohol.

- All-cause mortality is the number of deaths from all causes (rather than from specific causes like cardiovascular disease or cancer) during a specific time period.
- For the primary comparisons among those who currently drink alcohol:
  - Higher compared to lower average volume of alcohol consumption was related to higher all-cause mortality.
  - More frequent compared with less frequent or no binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) was related to higher all-cause mortality.
- For the secondary comparison between those who currently drink alcohol and those who have never consumed alcohol:
  - About half of the studies reported significant findings that low average alcohol consumption (particularly without binge drinking) compared with never drinking alcohol was associated with lower risk of mortality.
  - About half of the studies indicated no significant relationship.
  - Studies had several limitations. Factors other than drinking alcohol may impact all-cause mortality, and these factors were not always carefully addressed by the studies. The studies tended to examine middle- and older-aged adults, only. Some studies only measured alcohol consumption once, and that one measurement may not reflect participants' usual consumption. Studies varied in how they defined a "low-average" volume of alcohol.

#### **How up-to-date is this systematic review?**

- This review searched for studies from January 2010 to March 2020.

# TECHNICAL ABSTRACT

## Background

- This important public health question was identified by the U.S. Departments of Agriculture (USDA) and Health and Human Services (HHS) to be examined by the 2020 Dietary Guidelines Advisory Committee.
- The 2020 Dietary Guidelines Advisory Committee, Beverages and Added Sugars Subcommittee conducted a systematic review to answer this question with support from the Nutrition Evidence Systematic Review (NESR) team.
- The goal of this systematic review was to examine the following question: What is the relationship between alcohol consumption and all-cause mortality?

## Conclusion statements and grades

- **Primary comparisons (among those who currently drink alcohol)**
  - Moderate evidence indicates that higher average alcohol consumption is associated with an increased risk of all-cause mortality compared with lower average alcohol consumption among those who drink. (Grade: Moderate)
  - Moderate evidence indicates that binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) is associated with increased risk of all-cause mortality, and that more frequent binge drinking is associated with increased risk of all-cause mortality compared with less frequent or no binge drinking among those who drink. (Grade: Moderate)
- **Secondary comparison (between those who currently drink alcohol and those who have never consumed alcohol)**
  - Limited evidence suggests that low average alcohol consumption, particularly without binge drinking, is associated with a lower risk of all-cause mortality compared with never drinking alcohol. However, in light of the many scientific and public health issues associated with alcoholic beverages, any conclusions about low average consumption compared to never drinking alcohol require careful consideration. (Grade: Limited)

## Methods

- A literature search was conducted using 3 databases (PubMed, Cochrane, and Embase) to identify articles that evaluated the intervention or exposure of alcohol consumption and the outcome of all-cause mortality. A manual search was conducted to identify articles that may not have been included in the electronic databases searched. Articles were screened by two NESR analysts independently for inclusion based on pre-determined criteria.
- Data extraction and risk of bias assessment were conducted for each included study, and both were checked for accuracy. The Committee qualitatively synthesized the body of evidence to inform development of a conclusion statement(s), and graded the strength of evidence using pre-established criteria for risk of bias, consistency, directness, precision, and generalizability.

## Summary of the evidence

- Sixty studies that met the inclusion criteria for this systematic review addressing alcohol consumption and all-cause mortality were identified through the literature search from January 2010 to March 2020.
  - The body of evidence included one Mendelian randomization study, one retrospective cohort study, and 58 prospective cohort studies. The evidence included no RCTs.
- Consistent evidence reported increased all-cause mortality among those with higher average volume of alcohol consumption compared to lower average alcohol consumption. Although consumption categories varied, among those who drank alcohol most studies found lower risk among men consuming within ranges up to 2 drinks per day and women consuming within ranges up to 1 drink per day compared to those consuming higher average amounts. Among studies assessing continuous distributions or based on dose-response relationships among narrower consumption ranges among men who drink, the lowest levels of risk were generally up to 1 or 1.5 drinks on average (depending on how consumption was categorized). Relatively few studies among women examined risk based on categories within the range of up to 1 drink per day on average.
- Consistent evidence among those who drink alcohol reported higher all-cause mortality with more frequent binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) compared with less frequent or no binge drinking.
- For the secondary comparison between current drinkers and never drinkers, the limited available evidence suggested that low average consumption was associated with lower risk of mortality compared with never drinking status. Included studies were a subset of the 60 studies above that were used to assess the primary comparisons of interest.
- Twenty-five studies compared those who consumed alcohol with never drinkers. Approximately half of the studies reported significant findings that low average alcohol consumption (particularly without binge drinking) was associated with reduced risk of all-cause mortality compared with never drinking alcohol, approximately half of the studies indicated no significant relationship, and two studies reported that low alcohol consumption was significantly associated with greater all-cause mortality compared to never drinking alcohol.
- Generally, the evidence was limited by inadequate adjustment for confounders, selection bias and limited generalizability (studies often included only middle- and older-aged adults), and potential misclassification or bias from an exposure assessment based on single-time measurements of alcohol consumption. Low average volume was classified variably.
- Because the studies provided no consistent definition or categorization of higher average or lower average consumption, these terms are used in a descriptive sense in the conclusion statement. However, across most studies definitions of binge drinking or levels that corresponded to binge drinking were generally consistent; thus binge drinking is defined based on a set number of drinks in the conclusion statement.

## **FULL REVIEW**

### **Systematic review question**

What is the relationship between alcohol consumption and all-cause mortality?

### **Conclusion statements and grades**

#### **Primary comparisons (among those who currently drink alcohol)**

Moderate evidence indicates that higher average alcohol consumption is associated with an increased risk of all-cause mortality compared with lower average alcohol consumption among those who drink. (Grade: Moderate)

Moderate evidence indicates that binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) is associated with increased risk of all-cause mortality, and that more frequent binge drinking is associated with increased risk of all-cause mortality compared with less frequent or no binge drinking among those who drink. (Grade: Moderate)

#### **Secondary comparison (between those who currently drink alcohol and those who have never consumed alcohol)**

Limited evidence suggests that low average alcohol consumption, particularly without binge drinking, is associated with a lower risk of all-cause mortality compared with never drinking alcohol. However, in light of the many scientific and public health issues associated with alcoholic beverages, any conclusions about low average consumption compared to never drinking alcohol require careful consideration. (Grade: Limited)

### **Summary of the evidence**

- Sixty studies that met the inclusion criteria for this systematic review addressing alcohol consumption and all-cause mortality were identified through the literature search from January 2010 to March 2020.<sup>1-60</sup>
  - The body of evidence included one Mendelian randomization study, one retrospective cohort study, and 58 prospective cohort studies. The evidence included no RCTs.
- Consistent evidence reported increased all-cause mortality among those with higher average volume of alcohol consumption compared to lower average alcohol consumption. Although consumption categories varied, among those who drank alcohol most studies found lower risk among men consuming within ranges up to 2 drinks per day and women consuming within ranges up to 1 drink per day compared to those consuming higher average amounts. Among studies assessing continuous distributions or based on dose-response relationships among narrower consumption ranges among men who drink, the lowest levels of risk were generally up to 1 or 1.5 drinks on average (depending on how consumption was categorized). Relatively few studies among women examined risk based on categories within the range of up to 1 drink per day on average.

- Consistent evidence among those who drink alcohol reported higher all-cause mortality with more frequent binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) compared with less frequent or no binge drinking.
- For the secondary comparison between current drinkers and never drinkers, the limited available evidence suggested that low average consumption was associated with lower risk of mortality compared with never drinking status. Included studies were a subset of the 60 studies above that were used to assess the primary comparisons of interest.
- Twenty-five studies compared those who consumed alcohol with never drinkers. Approximately half of the studies reported significant findings that low average alcohol consumption (particularly without binge drinking) was associated with reduced risk of all-cause mortality compared with never drinking alcohol, approximately half of the studies indicated no significant relationship, and two studies reported that low alcohol consumption was significantly associated with greater all-cause mortality compared to never drinking alcohol.
- Generally, the evidence was limited by inadequate adjustment for confounders, selection bias and limited generalizability (studies often included only middle- and older-aged adults), and potential misclassification or bias from an exposure assessment based on single-time measurements of alcohol consumption. Low average volume was classified variably.
- Because the studies provided no consistent definition or categorization of higher average or lower average consumption, these terms are used in a descriptive sense in the conclusion statement. However, across most studies definitions of binge drinking or levels that corresponded to binge drinking were generally consistent; thus binge drinking is defined based on a set number of drinks in the conclusion statement.

## **Description of the evidence**

### **Primary and secondary comparisons**

This systematic review examined available evidence on the relationship between alcohol consumption and all-cause mortality in adults ages 21 years and older. Studies that exclusively enrolled participants younger than age 21 years were excluded to focus on the primary population of interest.

The search range included peer-reviewed articles published from January 2000 to March 2020. However, due to time constraints, the Committee revised their protocol to focus the review on studies published from January 2010 to March 2020. Studies were included if they were conducted in countries categorized as high or very high on the Human Development Index<sup>ii</sup>. Study designs that were included were: RCTs, non-

---

<sup>ii</sup> The Human Development classification was based on the Human Development Index (HDI) ranking (1) from the year the study intervention occurred or data was collected. If the study did not report the year in which the

randomized controlled trials, prospective and retrospective cohort studies, nested case-control studies, and Mendelian randomization studies. Studies were included if the study participants were generally healthy or at risk for chronic disease and for cohort studies that had sample sizes greater than 1,000 enrolled participants. Relationships between differing average alcohol consumption or patterns among those who drink was examined as a primary comparison. Although the primary comparison was different levels of alcohol consumption among those who currently drink, relationships between those who currently drink and those who have never consumed alcohol (i.e., lifetime abstainers) were also examined as a secondary comparison. Of note, articles were excluded if the reference group were “non-drinkers” who comprised those who used to drink but no longer drink (“former drinkers”) or a combination of lifetime abstainers and former drinkers. In papers that met all inclusion criteria, if the analysis included a non-reference group that combined never drinkers and former drinkers, these data were not extracted and therefore not used during synthesis.

The body of evidence included 60 articles (see **Table 1**): one Mendelian randomization study,<sup>1</sup> one retrospective cohort study,<sup>16</sup> and 58 prospective cohort studies.<sup>2-15,17-60</sup> The evidence included no RCTs.

### **Population**

The Mendelian randomization study enrolled male participants of Caucasian origin who were 65 to 84 years old.<sup>1</sup> The study was conducted in Australia and the analytic sample size was 3496.

Among the cohort studies, many studies examined adults of middle age and older (i.e., ages 45 and older); however, some population-based studies of adults (e.g., ages 18 years and older) or more representative age groups (e.g., ages 35 to 75 years) were included for both average consumption and patterns of consumption. Analytic sample sizes ranged from 1,121 to 599,912 participants; the majority (76%) of studies had analytic sample sizes greater than 10,000 participants. The majority (53%) of the prospective studies were from Europe (Belarus, Denmark, France, Germany, Greece, Hungary, Italy, Lithuania, The Netherlands, Norway, Poland, Russia, Spain, Sweden, and the United Kingdom). Several studies (36%) were from the United States. Several other high or very high HDI countries were represented, including: Argentina, Australia, Brazil, Canada, Chile, China, Colombia, Korea, Singapore, and Turkey. Four studies were conducted in all-male samples,<sup>5,30,40,47</sup> and four studies were in all-female samples.<sup>27,33,44,48</sup>

---

intervention occurred or data was collected, the HDI classification for the year of publication was applied. HDI values are available from 1980, and then from 1990 to present. If a study was conducted in 2018 or 2019, the most current HDI classification was applied. If a study was conducted prior to 1990, the HDI classification from 1990 was applied. When a country was not included in the HDI ranking, the current country classification from the World Bank (2) is used instead; 1. UN Development Program. HDI 1990-2017 HDRO calculations based on data from UNDESA (2017a), UNESCO Institute for Statistics (2018), United Nations Statistics Division (2018b), World Bank (2018b), Barro and Lee (2016) and IMF (2018). Available from: <http://hdr.undp.org/en/data>; 2. The World Bank. World Bank country and lending groups. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-country-and-lending-groups>

### ***Intervention/exposure/comparator***

The interventions or exposures of interest were average consumption of alcoholic beverages and the pattern of consumption of alcoholic beverages (i.e., the number of drinks per drinking day or drinks per drinking occasion, such as binge drinking). Information on the type of beverage (e.g., beer, wine, spirits) was collected if available. The primary comparator of interest was differing average alcohol consumption or patterns among those who currently drink. The secondary comparison was between those who currently drink and those who have never consumed alcohol (i.e., lifetime abstainers). Studies were excluded if the reference group were “non-drinkers” that comprised of former drinkers or a combination of never and former drinkers.

The Mendelian randomization study examined average alcohol consumption as drinks per day on an average week and compared never drinkers, past drinkers, and different levels of daily consumption among those who currently drink.<sup>1</sup> Alcohol consumption was measured once at baseline, and the study did not control for pattern of alcohol consumption.

Among the 59 cohort studies, 40 studies examined average alcohol consumption only, two studies measured alcohol consumption patterns only, and 16 studies measured both but often in quantity-frequency combinations that made it difficult to disaggregate effects of quantity consumed per drinking day from effects of frequency. There were 25 studies that addressed the relationship between those who currently drink and never drinkers; one study did not provide data on the average amount consumed or drinking pattern among those who currently drink.<sup>40</sup> The majority of studies were based on categorical alcohol consumption (e.g., lifetime abstainer, light, moderate, and heavy drinkers); seven studies reported continuous consumption measures among those who currently drink.<sup>2,7,13,35,38,46,56</sup> Most studies (63%) did not control for average alcohol consumption or pattern of consumption in the analysis for pattern of consumption and average consumption, respectively. Furthermore, most studies (86%) assessed alcohol consumption only at one point in time (baseline). Three studies reported results on the type of alcoholic beverage (such as beer, cider, champagne, wine, liquor and spirits) in addition to overall alcohol consumption,<sup>10,35,56</sup> and one study reported results only by type of alcoholic beverage.<sup>46</sup>

### ***Outcome***

The endpoint outcome for this systematic review was all-cause mortality (i.e., total mortality), which was defined as the total number of deaths from all causes during a specific time period. Although some studies disaggregated causes of death, the outcome for this review did not include cause-specific mortality (i.e., total number of deaths from a specific disease, such as CVD or cancer).

There were 35 studies that reported all-cause mortality among men and women combined<sup>6-16,21,25,26,28,31,34-39,45,46,50-60</sup>; 26 studies provided results stratified by sex,<sup>2-4,9,11-13,15-18,20,22-24,28,29,32,34,41-43,49,56,58,59</sup> four studies were in all-male samples,<sup>5,30,40,47</sup> and four studies were in all-female samples.<sup>27,33,44,48</sup> Most studies directly measured all-cause or total mortality as the outcome; however, two studies measured all-cause mortality indirectly. Bellavia et al<sup>2</sup> measured differences in 10th survival percentile (point in time by which the first 10% of the cohort has died), and van den Brandt et al<sup>56</sup> measured the risk of reaching longevity until age 90.

## Evidence synthesis

### Primary comparisons (among those who currently drink alcohol)

The majority of studies examining the relationship between alcohol consumption and all-cause mortality found a statistically significant relationship (**Table 1**).

The Mendelian randomization study reported that among older men (70 to 89 years), having a genetic variant associated with reduced alcohol consumption was associated with lower total mortality.<sup>1</sup> However, the study did not distinguish between various levels of consumption by genetic variant.

Among the 59 cohort studies, over half reported a significant association between higher average volume of consumption and higher all-cause mortality compared to lower average volume consumption, with generally consistent dose-response relationships, at least with respect to point estimates. Among those who drink, most studies found lower risk among men consuming between 0 and 2 drinks per day, and women consuming between 0 and 1 drink per day. Among studies assessing continuous distributions or based on dose-response analyses among narrower average consumption ranges, approximately half of the studies found lower risks for men consuming at or below approximately 1 to 1.5 drinks per day compared to those consuming higher average amounts. Few studies among women examined categories of risk within the 0 to 1 range of average drinks.

In the body of evidence, 18 observational studies examined drinking patterns. Three studies assessed the relationship between the usual number of drinks typically consumed per drinking day or occasion and all-cause mortality. Among men, 3 studies found that consuming more than 2 drinks per drinking day was associated with higher mortality risk compared with consuming less; only 1 study examined differences among men comparing 1 vs 2 drinks. For women, both studies found increased risk for all-cause mortality with consumption greater than 1 drink. The majority of studies examining drinking patterns assessed relationships based on binge drinking (typically defined as consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion). Although not all studies defined binge drinking in the same manner, studies consistently found that among those who drink, binge drinking was associated with increased mortality risk compared to not binge drinking, and that more frequent binge drinking was associated with increased risk compared with less frequent binge drinking.

Funding sources were documented during data extraction for consideration when reviewing this evidence. Further, publication bias is always a consideration in systematic reviews.



## Assessment of the evidence<sup>iii</sup>

### 1. *Average consumption among those who currently drink alcohol:*

The following conclusion statement was assigned a grade of moderate: “Moderate evidence indicates that higher average alcohol consumption is associated with an increased risk of all-cause mortality compared with lower average alcohol consumption among those who drink.”

As outlined and described below, the body of evidence examining average consumption among those who currently drink alcohol and all-cause mortality was assessed for the following elements when grading the strength of evidence.

**Consistency:** A relatively large number of studies informed the primary comparisons, with generally consistent findings for both the United States and other high-income nations on the association between average alcohol consumption and all-cause mortality. However, studies used different levels of comparison (and terminology) to define lower versus higher average consumption. Further, for lower average consumption, some studies assessed a subgroup of those who consumed very small amounts of alcohol and/or had infrequent consumption. Although their risk was generally less than for higher average consumption, findings were inconsistent about whether they had higher risk than lower average volume drinkers. Not all studies reported consumption in grams of ethanol, but rather in drinks per day or per week.

**Directness:** The studies were designed to directly examine the relationships among alcohol consumption, the comparator, and all-cause mortality in the systematic review question. All studies assessed alcohol consumption, and most examined average consumption only with a few that accounted for volume and pattern concurrently.

**Precision:** The Mendelian randomization study had a relatively small size (<4000 participants), particularly for this type of study. However, there were numerous observational studies with relatively large sample sizes. Exposures were often categorical, and rarely continuous. Studies using “occasional or light” drinkers as the reference group had less precision to the extent that occasional drinkers are not a large proportion of most study populations; this makes it difficult to interpret overlapping confidence intervals between “moderate” and heavier drinkers since direct comparisons between those groups should yield narrower confidence intervals.

**Generalizability:** Since most studies were in older age cohorts, selection bias based on the age of the study cohorts is a possibility. A large percentage of alcohol-attributable deaths occur before age 50 years, and those who have been established moderate drinkers for longer periods or until later in life may be advantaged, socially, or in terms of health. Studies of older cohorts could lead to underestimation of alcohol-mortality associations compared with studies that are population-based.

---

<sup>iii</sup> A detailed description of the methodology used for grading the strength of the evidence is available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews> and in Part C of the following reference: Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.

**Risk of bias:** There were a number of potential risks of bias, or limitations across the body of evidence (see summary of risk of bias in **Table 2**). The evidence included no RCTs. Assessed studies did not control for multiple key confounders (e.g., socioeconomic factors, diet quality), and did not typically account for patterns of alcohol consumption. Failing to account for patterning as a confounder may affect the generalizability of studies because people participating in surveys or cohorts may have different patterns of consumption (i.e., less binge drinking) or alcohol-related problems compared to people in the general population, even among those with similar consumption levels. Selection bias is also a concern, mostly due to the age of participants as noted above, and measurement of exposure because it is almost always self-report. Furthermore, the majority of studies measured alcohol at one point in time contributing to risk of bias from deviations from the assessed exposure resulting in possible misclassification of typical average consumption. Measuring consumption at one point in time also precludes assessments related to drinking trajectories over time, or the effects of cumulative alcohol consumption over the life course. Alcohol consumption typically changes throughout the life course.

## ***2. Binge drinking among those who drink:***

The following conclusion statement was assigned a grade of moderate: “Moderate evidence indicates that binge drinking (consuming 5 or more drinks for men or 4 or more drinks for women during a drinking occasion) is associated with increased risk of all-cause mortality, and that more frequent binge drinking is associated with increased risk of all-cause mortality compared with less frequent or no binge drinking among those who drink.”

As outlined and described below, the body of evidence examining binge drinking among those who drink alcohol and all-cause mortality was assessed for the following elements when grading the strength of evidence.

**Consistency:** Findings were generally consistent in terms of the direction of association, magnitude and significance even though not all studies defined binge drinking the same way.

**Directness:** Studies were designed to directly examine the relationships among the alcohol pattern, comparator, and all-cause mortality in the systematic review question. However, not all studies measured alcohol pattern in the same manner and few accounted for volume and pattern concurrently.

**Precision:** The observational studies had relatively large sample sizes. However, exposures were often categorical, and therefore were less precise.

**Generalizability:** Studies may have included participants with less binge drinking or other risky patterns of consumption compared to studies of the general population, even among those with the same average consumption levels. This could lead to underestimates of alcohol-related risk in cohort or survey participants compared to the general population, and adversely affects the generalizability of findings. The majority of studies were conducted in European countries, and findings may not be generalizable to the United States. Confounders related to binge drinking may vary by gender, and few studies reported binge drinking in men and women separately.

**Risk of bias:** There were several potential risks of bias, or limitations across the body

of evidence (see summary of risk of bias in **Table 2**). There evidence included no RCTs. The risk of confounding bias was high overall. Assessed studies did not control for multiple key confounders (e.g., socioeconomic factors, diet quality), and did not typically account for average consumption. Despite the consistency and magnitude of findings, binge drinkers (or those consuming high average amounts) may have negative associated confounders. Selection bias is also a concern due to the age of participants, and measurement of exposure because it is almost always self-report. Furthermore, most studies measured alcohol at one point in time contributing to risk of bias from deviations from the assessed exposure resulting in possible misclassification of patterns of consumption compared to typical consumption over the life course.

### **Secondary comparison (between those who currently drink alcohol and those who have never consumed alcohol)**

There were 25 studies that compared those who currently drink alcohol and those who have never consumed alcohol (**Table 1**). Approximately half of the studies reported significant findings that low average alcohol consumption (particularly without binge drinking) was associated with reduced risk of all-cause mortality compared with never drinkers. Approximately half of the data indicated no significant relationship between low average alcohol consumption and never consuming alcohol on all-cause mortality. Two studies reported that low alcohol consumption was significantly associated with greater all-cause mortality compared to never drinkers.<sup>16,37</sup> Horvat et al<sup>16</sup> found that participants from Russia, but not Belarus or Hungary, who drank up to once a month had significantly higher risk of all-cause mortality compared to never drinkers. A study from the United Kingdom reported that occasional drinkers (those who declared not having drunk in the past 7 days) had reduced risk of all-cause mortality compared to never drinkers, but those who consumed alcohol within the UK guidelines (<14 units per week for women; <21 units per week for men) or more had a greater risk of all-cause mortality compared to never drinkers.<sup>37</sup> Three studies reported that “high” average drinkers had greater risk of all-cause mortality compared to never drinkers.<sup>17,35,51</sup>

There are particularities worth noting within this body of evidence, mainly regarding the population of those who never consumed alcohol (i.e., never drinkers). Definitions of never drinkers varied among studies. Furthermore, never drinkers may have more adverse confounding factors compared with those who drink, and all studies lacked inclusion of key confounders. A sizable fraction of self-described never drinkers may be former drinkers and are therefore misclassified. In addition, there were fewer studies of never drinkers than those of within-drinker comparisons; some studies were restricted to those who drink, while others assessed non-drinkers as a group and could not or did not disaggregate former drinkers from never drinkers.

Funding sources were documented during data extraction for consideration when reviewing this evidence. Further, publication bias is always a consideration in systematic reviews.

## Assessment of the evidence<sup>iv</sup>

### 3. Low average consumption vs never drinking alcohol:

The following conclusion statement was assigned a grade of limited: “Limited evidence suggests that low average alcohol consumption, particularly without binge drinking, is associated with a lower risk of all-cause mortality compared with never drinking alcohol. However, in light of the many scientific and public health issues associated with alcoholic beverages, any conclusions about low average consumption compared to never drinking alcohol require careful consideration.”

As outlined and described below, the body of evidence examining low average alcohol consumption vs never drinking alcohol and all-cause mortality was assessed for the following elements when grading the strength of evidence.

**Consistency:** The data was largely either significant indicating lower all-cause mortality risk for low average alcohol consumption compared to never drinking alcohol, or non-significant. Few studies reported significant findings in the opposite direction.

**Directness:** While the review methods focused only on data from “never drinkers” rather than “non-drinkers” (which combines never and former drinkers), studies varied in their assignment and definition of never drinkers. Further, in some studies, due to self-report, participants may have incorrectly classified themselves as “never drinkers”.

**Precision:** Since “never drinkers” were a relatively small proportion of the study populations, confidence intervals tended to be wide, thus limiting precision.

**Generalizability:** This evidence cannot be generalized to inform never drinkers who are considering drinking alcohol; nor can it be generalized to inform those who currently drink to become “never drinkers”. Many study populations were not representative of the age distribution of all adults who typically begin to consume alcohol.

**Risk of bias:** There were a number of potential risks of bias, or limitations across the body of evidence (see summary of risk of bias in **Table 2**). There evidence included no RCTs. Risk of bias is a concern for this body of evidence, particularly related to additional confounding issues with the “never drinker” population. All studies lacked inclusion of key pre-specified confounders. There are also concerns around selection bias due to the age of participants at the time of data assessment and measurement of exposure because it is almost always self-report. Furthermore, the majority of studies measured alcohol at one point in time contributing to risk of bias from deviations from the assessed exposure resulting in possible misclassification of alcohol consumption.

---

<sup>iv</sup> A detailed description of the methodology used for grading the strength of the evidence is available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews> and in Part C of the following reference: Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.

## Research recommendations

To better assess the relationship between alcohol consumption and all-cause mortality, future research may:

- Measure alcoholic beverage intake and consumption patterns with greater accuracy and determine their effect on human health through all appropriate life stages using more consistent and improved definitions, controls, and research methods, such as:
  - Stronger research designs, including randomized clinical trials, Mendelian randomization studies, and non-randomized intervention studies
  - Multiple measures of intake over time rather than at one time point
  - Distinction between lifelong never drinkers and either very low-level consumers or former drinkers who refer to themselves as never drinkers
  - Long timeframes for studies and better control for key confounders, including: sex, age, race/ethnicity, socioeconomic status, eating pattern or diet quality, physical activity, and smoking
  - Enrollment of younger and lower exposure drinkers who may have more adverse drinking outcomes
  - More research on patterns of consumption, including health outcomes in relation to the frequency and usual amount of alcohol consumed during drinking days
- Study the effects of alcohol consumption on:
  - Dietary patterns and overall diet quality
  - Growth, size, body composition, and risk of overweight and obesity
  - Risk of intermediate cardiovascular outcomes
  - Risk of type 2 diabetes, including markers for prediabetes
  - Risk of cancer, including mouth, throat, larynx, esophagus, colon and rectum, liver, and breast in women
  - Neurocognitive health, including alcohol consumption during lactation and neurocognitive health of the infant

## Included articles

1. Almeida OP, McCaul K, Hankey GJ, Yeap BB, Golledge J, Flicker L. Excessive alcohol consumption increases mortality in later life: a genetic analysis of the health in men cohort study. *Addict Biol.* 2017;22(2):570-578. doi: 10.1111/adb.12340.
2. Bellavia A, Bottai M, Wolk A, Orsini N. Alcohol consumption and mortality: a dose-response analysis in terms of time. *Ann Epidemiol.* 2014;24(4):291-296. doi: 10.1016/j.annepidem.2013.12.012.
3. Bergmann MM, Rehm J, Klipstein-Grobusch K, et al. The association of pattern of lifetime alcohol use and cause of death in the European prospective investigation into cancer and nutrition (EPIC) study. *Int J Epidemiol.* 2013;42(6):1772-1790. doi: 10.1093/ije/dyt154.
4. Bobak M, Malyutina S, Horvat P, et al. Alcohol, drinking pattern and all-cause, cardiovascular and alcohol-related mortality in Eastern Europe. *Eur J Epidemiol.* 2016;31(1):21-30. doi: 10.1007/s10654-015-0092-8.
5. Britton A, Marmot MG, Shipley MJ. How does variability in alcohol consumption over time affect the relationship with mortality and coronary heart disease? *Addiction.* 2010;105(4):639-645. doi: 10.1111/j.1360-0443.2009.02832.x.
6. Britton A, O'Neill D, Bell S. Underestimating the alcohol content of a glass of wine: the implications for estimates of mortality risk. *Alcohol Alcohol.* 2016;51(5):609-614. doi: 10.1093/alcalc/agw027.
7. Degerud E, Hoiseth G, Morland J, et al. Association of coincident self-reported mental health problems and alcohol intake with all-cause and cardiovascular disease mortality: A Norwegian pooled population analysis. *PLoS Med.* 2020;17(2):e1003030. doi: 10.1371/journal.pmed.1003030.
8. Evans-Polce RJ, Staff J, Maggs JL. Alcohol abstinence in early adulthood and premature mortality: do early life factors, social support, and health explain this association? *Soc Sci Med.* 2016;163:71-79. doi: 10.1016/j.socscimed.2016.06.052.
9. Ferrari P, Licaj I, Muller DC, et al. Lifetime alcohol use and overall and cause-specific mortality in the European Prospective Investigation into Cancer and nutrition (EPIC) study. *BMJ Open.* 2014;4(7):e005245. doi: 10.1136/bmjopen-2014-005245.
10. Gea A, Bes-Rastrollo M, Toledo E, et al. Mediterranean alcohol-drinking pattern and mortality in the SUN (Seguimiento Universidad de Navarra) Project: a prospective cohort study. *Br J Nutr.* 2014;111(10):1871-1880. doi: 10.1017/s0007114513004376.
11. Goulden R. Moderate alcohol consumption is not associated with reduced all-cause mortality. *Am J Med.* 2016;129(2):180-186.e184. doi: 10.1016/j.amjmed.2015.10.013.
12. Graff-Iversen S, Jansen MD, Hoff DA, et al. Divergent associations of drinking frequency and binge consumption of alcohol with mortality within the same cohort. *J Epidemiol Community Health.* 2013;67(4):350-357. doi: 10.1136/jech-2012-201564.
13. Hartz SM, Oehlert M, Horton AC, et al. Daily drinking is associated with increased mortality. *Alcohol Clin Exp Res.* 2018;42(11):2246-2255. doi: 10.1111/acer.13886.

14. Holahan CJ, Schutte KK, Brennan PL, Holahan CK, Moos BS, Moos RH. Late-life alcohol consumption and 20-year mortality. *Alcohol Clin Exp Res*. 2010;34(11):1961-1971. doi: 10.1111/j.1530-0277.2010.01286.x.
15. Holahan CJ, Schutte KK, Brennan PL, Holahan CK, Moos RH. Drinking level, drinking pattern, and twenty-year total mortality among late-life drinkers. *J Stud Alcohol Drugs*. 2015;76(4):552-558. doi: 10.15288/jsad.2015.76.552.
16. Horvat P, Stefler D, Murphy M, King L, McKee M, Bobak M. Alcohol, pattern of drinking and all-cause mortality in Russia, Belarus and Hungary: a retrospective indirect cohort study based on mortality of relatives. *Addiction*. 2018;113(7):1252-1263. doi: 10.1111/add.14189.
17. Jackson CL, Hu FB, Kawachi I, Williams DR, Mukamal KJ, Rimm EB. Black-white differences in the relationship between alcohol drinking patterns and mortality among US men and women. *Am J Public Health*. 2015;105 Suppl 3:S534-543. doi: 10.2105/ajph.2015.302615.
18. Jayasekara H, MacInnis RJ, Hodge AM, et al. Alcohol consumption for different periods in life, intake pattern over time and all-cause mortality. *J Public Health*. 2015;37(4):625-633. doi: 10.1093/pubmed/fdu082.
19. Jung EJ, Shin A, Park SK, et al. Alcohol consumption and mortality in the Korean Multi-Center Cancer Cohort Study. *J Prev Med Public Health*. 2012;45(5):301-308. doi: 10.3961/jpmph.2012.45.5.301.
20. Kabat GC, Matthews CE, Kamensky V, Hollenbeck AR, Rohan TE. Adherence to cancer prevention guidelines and cancer incidence, cancer mortality, and total mortality: a prospective cohort study. *Am J Clin Nutr*. 2015;101(3):558-569. doi: 10.3945/ajcn.114.094854.
21. Kerr WC, Greenfield TK, Bond J, Ye Y, Rehm J. Racial and ethnic differences in all-cause mortality risk according to alcohol consumption patterns in the national alcohol surveys. *Am J Epidemiol*. 2011;174(7):769-778. doi: 10.1093/aje/kwr147.
22. Keyes KM, Calvo E, Ornstein KA, et al. Alcohol consumption in later life and mortality in the United States: results from 9 waves of the Health and Retirement Study. *Alcohol Clin Exp Res*. 2019;43(8):1734-1746. doi: 10.1111/acer.14125.
23. Knott CS, Coombs N, Stamatakis E, Biddulph JP. All cause mortality and the case for age specific alcohol consumption guidelines: pooled analyses of up to 10 population based cohorts. *BMJ*. 2015;350:h384. doi: 10.1136/bmj.h384.
24. Kunzmann AT, Coleman HG, Huang WY, Berndt SI. The association of lifetime alcohol use with mortality and cancer risk in older adults: a cohort study. *PLoS Med*. 2018;15(6):e1002585. doi: 10.1371/journal.pmed.1002585.
25. Lantz PM, Golberstein E, House JS, Morenoff J. Socioeconomic and behavioral risk factors for mortality in a national 19-year prospective study of U.S. adults. *Soc Sci Med*. 2010;70(10):1558-1566. doi: 10.1016/j.socscimed.2010.02.003.
26. Li Y, Pan A, Wang DD, et al. Impact of healthy lifestyle factors on life expectancies in the US population. *Circulation*. 2018;138(4):345-355. doi: 10.1161/circulationaha.117.032047.
27. Licaj I, Sandin S, Skeie G, Adami HO, Roswall N, Weiderpass E. Alcohol consumption over time and mortality in the Swedish Women's Lifestyle and Health cohort. *BMJ Open*. 2016;6(11):e012862. doi: 10.1136/bmjopen-2016-012862.

28. Lindschou Hansen J, Tolstrup JS, Jensen MK, et al. Alcohol intake and risk of acute coronary syndrome and mortality in men and women with and without hypertension. *Eur J Epidemiol*. 2011;26(6):439-447. doi: 10.1007/s10654-011-9564-7.
29. Luksiene D, Tamosiunas A, Virviciute D, Radisauskas R. The prognostic value of combined smoking and alcohol consumption habits for the estimation of cause-specific mortality in middle-age and elderly population: results from a long-term cohort study in Lithuania. *Biomed Res Int*. 2017;2017:9654314. doi: 10.1155/2017/9654314.
30. Lundin A, Mortensen LH. Mortality from alcohol consumption and alcohol use disorder: findings from the Vietnam Experience Study. *Drug Alcohol Depend*. 2015;151:135-143. doi: 10.1016/j.drugalcdep.2015.03.013.
31. Lundin A, Mortensen LH, Halldin J, Theobald H. The relationship of alcoholism and alcohol consumption to all-cause mortality in forty-one-year follow-up of the Swedish REBUS sample. *J Stud Alcohol Drugs*. 2015;76(4):544-551. doi: 10.15288/jsad.2015.76.544.
32. McCullough ML, Patel AV, Kushi LH, et al. Following cancer prevention guidelines reduces risk of cancer, cardiovascular disease, and all-cause mortality. *Cancer Epidemiol Biomarkers Prev*. 2011;20(6):1089-1097. doi: 10.1158/1055-9965.Epi-10-1173.
33. Midlov P, Calling S, Memon AA, Sundquist J, Sundquist K, Johansson SE. Women's health in the Lund area (WHILA)--Alcohol consumption and all-cause mortality among women--a 17 year follow-up study. *BMC Public Health*. 2016;16:22. doi: 10.1186/s12889-016-2700-2.
34. Muller DC, Murphy N, Johansson M, et al. Modifiable causes of premature death in middle-age in Western Europe: results from the EPIC cohort study. *BMC Med*. 2016;14:87. doi: 10.1186/s12916-016-0630-6.
35. Ortola R, Garcia-Esquinas E, Lopez-Garcia E, Leon-Munoz LM, Banegas JR, Rodriguez-Artalejo F. Alcohol consumption and all-cause mortality in older adults in Spain: an analysis accounting for the main methodological issues. *Addiction*. 2019;114(1):59-68. doi: 10.1111/add.14402.
36. Pan XF, Li Y, Franco OH, Yuan JM, Pan A, Koh WP. Impact of combined lifestyle factors on all-cause and cause-specific mortality and life expectancy in Chinese: the Singapore Chinese Health Study. *J Gerontol A Biol Sci Med Sci*. 2019. doi: 10.1093/gerona/glz271.
37. Perreault K, Bauman A, Johnson N, Britton A, Rangul V, Stamatakis E. Does physical activity moderate the association between alcohol drinking and all-cause, cancer and cardiovascular diseases mortality? A pooled analysis of eight British population cohorts. *Br J Sports Med*. 2017;51(8):651-657. doi: 10.1136/bjsports-2016-096194.
38. Plunk AD, Syed-Mohammed H, Cavazos-Rehg P, Bierut LJ, Grucza RA. Alcohol consumption, heavy drinking, and mortality: rethinking the j-shaped curve. *Alcohol Clin Exp Res*. 2014;38(2):471-478. doi: 10.1111/acer.12250.
39. Powell L, Feng Y, Duncan MJ, Hamer M, Stamatakis E. Does a physically active lifestyle attenuate the association between alcohol consumption and mortality risk? Findings from the UK biobank. *Prev Med*. 2020;130:105901. doi: 10.1016/j.ypmed.2019.105901.



40. Rhee CW, Kim JY, Park BJ, Li ZM, Ahn YO. Impact of individual and combined health behaviors on all causes of premature mortality among middle aged men in Korea: the Seoul Male Cohort Study. *J Prev Med Public Health*. 2012;45(1):14-20. doi: 10.3961/jpmph.2012.45.1.14.
41. Ricci C, Schutte AE, Schutte R, Smuts CM, Pieters M. Trends in alcohol consumption in relation to cause-specific and all-cause mortality in the United States: a report from the NHANES linked to the US mortality registry. *Am J Clin Nutr*. 2020;111(3):580-589. doi: 10.1093/ajcn/nqaa008.
42. Rosella LC, Kornas K, Huang A, Grant L, Bornbaum C, Henry D. Population risk and burden of health behavioral-related all-cause, premature, and amenable deaths in Ontario, Canada: Canadian Community Health Survey-linked mortality files. *Ann Epidemiol*. 2019;32:49-57.e43. doi: 10.1016/j.annepidem.2019.01.009.
43. Rostron B. Alcohol consumption and mortality risks in the USA. *Alcohol Alcohol*. 2012;47(3):334-339. doi: 10.1093/alcalc/agr171.
44. Rundberg J, Nilsson PM, Samsioe G, Ojehagen A. Alcohol use and early mortality in Swedish middle-aged women: nine-year follow-up of the Women's Health in Lund Area study. *Scand J Public Health*. 2014;42(4):344-348. doi: 10.1177/1403494814523343.
45. Schoenborn CA, Stommel M, Ward BW. Mortality risks associated with average drinking level and episodic heavy drinking. *Subst Use Misuse*. 2014;49(10):1250-1258. doi: 10.3109/10826084.2014.891620.
46. Schutte R, Papageorgiou M, Najlah M, et al. Drink types unmask the health risks associated with alcohol intake - prospective evidence from the general population. *Clin Nutr*. 2020. doi: 10.1016/j.clnu.2020.02.009.
47. Shuval K, Barlow CE, Chartier KG, Gabriel KP. Cardiorespiratory fitness, alcohol, and mortality in men: the Cooper Center longitudinal study. *Am J Prev Med*. 2012;42(5):460-467. doi: 10.1016/j.amepre.2012.01.012.
48. Simpson TL, Rillamas-Sun E, Lehavot K, et al. Alcohol consumption levels and all-cause mortality among women veterans and non-veterans enrolled in the Women's Health Initiative. *Gerontologist*. 2016;56 Suppl 1:S138-149. doi: 10.1093/geront/gnv667.
49. Skov-Ettrup LS, Eliassen M, Ekholm O, Gronbaek M, Tolstrup JS. Binge drinking, drinking frequency, and risk of ischaemic heart disease: a population-based cohort study. *Scand J Public Health*. 2011;39(8):880-887. doi: 10.1177/1403494811425605.
50. Sluik D, Boeing H, Li K, et al. Lifestyle factors and mortality risk in individuals with diabetes mellitus: are the associations different from those in individuals without diabetes? *Diabetologia*. 2014;57(1):63-72. doi: 10.1007/s00125-013-3074-y.
51. Smyth A, Teo KK, Rangarajan S, et al. Alcohol consumption and cardiovascular disease, cancer, injury, admission to hospital, and mortality: a prospective cohort study. *Lancet*. 2015;386(10007):1945-1954. doi: 10.1016/s0140-6736(15)00235-4.
52. Soedamah-Muthu SS, De Neve M, Shelton NJ, Tielemans SM, Stamatakis E. Joint associations of alcohol consumption and physical activity with all-cause and cardiovascular mortality. *Am J Cardiol*. 2013;112(3):380-386. doi: 10.1016/j.amjcard.2013.03.040.

53. Stringhini S, Dugravot A, Shipley M, et al. Health behaviours, socioeconomic status, and mortality: further analyses of the British Whitehall II and the French GAZEL prospective cohorts. *PLoS Med*. 2011;8(2):e1000419. doi: 10.1371/journal.pmed.1000419.
54. Syden L, Landberg J. The contribution of alcohol use and other lifestyle factors to socioeconomic differences in all-cause mortality in a Swedish cohort. *Drug Alcohol Rev*. 2017;36(5):691-700. doi: 10.1111/dar.12472.
55. Tevik K, Selbaek G, Engedal K, Seim A, Krokstad S, Helvik AS. Mortality in older adults with frequent alcohol consumption and use of drugs with addiction potential - The Nord Trondelag Health Study 2006-2008 (HUNT3), Norway, a population-based study. *PLoS One*. 2019;14(4):e0214813. doi: 10.1371/journal.pone.0214813.
56. van den Brandt PA, Brandts L. Alcohol consumption in later life and reaching longevity: the Netherlands Cohort Study. *Age Ageing*. 2020. doi: 10.1093/ageing/afaa003.
57. van Hedel K, van Lenthe FJ, Oude Groeniger J, Mackenbach JP. What's the difference? A gender perspective on understanding educational inequalities in all-cause and cause-specific mortality. *BMC Public Health*. 2018;18(1):1105. doi: 10.1186/s12889-018-5940-5.
58. Warren Andersen S, Zheng W, Sonderman J, et al. Combined impact of health behaviors on mortality in low-income Americans. *Am J Prev Med*. 2016;51(3):344-355. doi: 10.1016/j.amepre.2016.03.018.
59. Wood AM, Kaptoge S, Butterworth AS, et al. Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599 912 current drinkers in 83 prospective studies. *Lancet*. 2018;391(10129):1513-1523. doi: 10.1016/s0140-6736(18)30134-x.
60. Xi B, Veeranki SP, Zhao M, Ma C, Yan Y, Mi J. Relationship of alcohol consumption to all-cause, cardiovascular, and cancer-related mortality in U.S. adults. *J Am Coll Cardiol*. 2017;70(8):913-922. doi: 10.1016/j.jacc.2017.06.054.

**Table 1: Summary of articles that address the relationship between alcohol consumption and all-cause mortality<sup>v</sup>**

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Almeida, 2017<sup>1</sup></b> Mendelian Randomization Study (PCS), Health in Men Study (HIMS) Australia N=3,496  <b>Participants:</b> ~77y (65-84y), 100% male; excluded: those with missing data on alcohol consumption or <i>ADH1B</i> rs1229984 G>A genotype, men of possible non-Caucasian origin	<b>Baseline:</b> 1996-1998 (for alcohol intake); 2001-2004 for DNA <b>Follow-up:</b> 2012, mean: 8y  <b>Average intake: drinks/d on an average week</b> <ul style="list-style-type: none"> <li><b>Never drinker (REF)</b></li> <li>Past: none in preceding year</li> <li>≤2 drinks/d</li> <li>2-4 drinks/d</li> <li>4-6 drinks/d</li> <li>&gt;6 drinks/d</li> </ul> <i>ADH1B</i> rs1229984 G>A genotype assessed at first follow-up visit (2001-2004) (6.4% were carriers and consumed significantly less alcohol)	<ul style="list-style-type: none"> <li>Never drinker (REF)</li> <li>Past: NS</li> <li>≤2 drinks/d: NS</li> <li>2-4 drinks/d: NS</li> <li>4-6 drinks/d: NS</li> <li><b>&gt;6 drinks/d: 1.87 (1.11, 3.12)</b></li> </ul> <b>ACM HR by genotype</b> Carriers have lower mortality risk if alcohol is not consumed "We found evidence of a statistically significant interaction between alcohol use group and being a carrier."	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), physical activity, smoking  <b>Other factors considered:</b> anthropometry  <b>Other:</b> prevalent coronary heart disease and stroke	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern</li> <li>Older cohort: Potential survival bias</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>6-year time lag between alcohol assessment and DNA assessment</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

<sup>v</sup> Abbreviations/short-hand: ACM: all-cause mortality; ADG: aggregated diagnosis group; adj: adjusted; AHEI: alternate healthy eating index; Avg: average; B: beta; BMI: body mass index; CHD: coronary heart disease; CI: confidence interval; CVD: cardiovascular disease; d: day(s); DGA: Dietary Guidelines for Americans; DM: diabetes mellitus; ECG: electrocardiogram; EER: estimated energy requirement; ETOH: ethyl alcohol; FFQ: food frequency questionnaire; g: gram(s); HEI: healthy eating index; HR: hazard ratio; HRT: hormone replacement therapy; IADL: instrumental activities of daily living; kg/m<sup>2</sup>: kilograms per meters squared; ml: milliliters; mo: month(s); N/A: not applicable; NS: not significant; NR: not reported; occ: occasion; OR: odds ratio; PAR: population-attributable risks; PCS: prospective cohort study; PD: percentile difference; pts: points; RCT: randomized controlled trial; REF: reference group; RR: risk ratio; SD: standard deviation; SE: standard error; SES: socioeconomic status; UK: United Kingdom; unadj: unadjusted; wk: week(s); y: year(s)  
Red font indicates a statistically significant detrimental relationship, and green font indicates a statistically significant beneficial relationship.

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Bellavia, 2014<sup>2</sup></b> Prospective Cohort Study, Cohort of Swedish Men and Swedish Mammography Cohort Sweden N=67,706  <b>Participants:</b> 45-83y (~60y), 46% female; excluded participants who reported incorrect or missing personal numbers, those who died before start of follow-up, those who had diabetes or any history of CVD and cancer, and those with missing information on all alcohol-related items; former drinkers were also excluded from the analysis	<b>Baseline:</b> 1997-1998 <b>Follow-up:</b> 2012 (15y)  <b>Average intake: frequency (FFQ) + amount (per occasion)</b> <ul style="list-style-type: none"> <li><b>Lifetime abstainers (REF)</b></li> <li>&lt;1.5 g/d (women)</li> <li>1.5-5 g/d (women) or &lt;5 g/d (men)</li> <li>5-10 g/d</li> <li>10-15 g/d</li> <li>&gt;15 g/d (women) or 15-20 g/d (men)</li> <li>20-25 g/d (men)</li> <li>25-30 g/d (men)</li> <li>&gt;30 g/d (men)</li> </ul> 1 drink = 12g ETOH	<b>Differences in 10<sup>th</sup> survival percentile</b> , Laplace regression, 10 <sup>th</sup> PD, months (95% CI) <b>Males:</b> <ul style="list-style-type: none"> <li>Lifetime abstainers (REF)</li> <li>&lt;5: NS</li> <li>5-10: 9 (-1, 18)</li> <li>10-15: 16 (6, 25)</li> <li>15-20: 14 (3, 24)</li> <li>20-25: NS</li> <li>25-30: NS</li> <li>&gt;30: NS</li> </ul> <b>Females:</b> <ul style="list-style-type: none"> <li>Lifetime abstainers (REF)</li> <li>&lt;1.5: 15 (8, 23)</li> <li>1.5-5: 17 (9, 24)</li> <li>5-10: 22 (13, 30)</li> <li>10-15: 13 (3, 23)</li> <li>&gt;15: 16 (5, 26)</li> </ul> <b>HR (95% CI)</b> <b>Males:</b> <ul style="list-style-type: none"> <li>Lifetime abstainers (REF)</li> <li>&lt;5: NS</li> <li>5-10: 0.86 (0.76, 0.97)</li> <li>10-15: 0.77 (0.68, 0.88)</li> <li>15-20: 0.80 (0.70, 0.93)</li> <li>20-25: NS</li> <li>25-30: NS</li> <li>&gt;30: NS</li> </ul> <b>Females:</b> <ul style="list-style-type: none"> <li>Lifetime abstainers (REF)</li> <li>&lt;1.5: 0.81 (0.73, 0.89)</li> <li>1.5-5: 0.77 (0.70, 0.85)</li> <li>5-10: 0.72 (0.64, 0.81)</li> <li>10-15: 0.74 (0.64, 0.85)</li> <li>&gt;15: 0.78 (0.68, 0.91)</li> </ul>	<b>Key confounders:</b> Sex, age, SES, physical activity, smoking  <b>Other factors considered:</b> age distribution of the study sample, anthropometry, diabetes  <b>Other:</b> N/A	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol pattern</li> <li>Exposure data only measured once</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Bergmann, 2013<sup>3</sup></b> Prospective Cohort Study, European Prospective Investigation into Cancer and nutrition (EPIC) Denmark, France, Germany, Greece, Italy, Netherlands, Spain, and UK N=380,395</p> <p><b>Participants:</b> 25-70y (~52y), 71% female; excluded cohorts of France (female members of a health insurance for school employees), Utrecht (breast cancer screening attendees), Spain (mainly blood donors and their spouses), and Oxford (mainly vegetarian and health-conscious people), and study centers where information on alcohol use in the past was not collected; participants with no follow-up or information on vital status, those with missing data on dietary or lifestyle questionnaire or alcohol use in the past, and those whose ratio of energy intake to EER was in top or bottom 1%</p>	<p><b>Baseline:</b> 1992-2000 <b>Follow-up:</b> 12.6y</p> <p><b>Average intake: glasses/d by beverage type, converted to g/d—reflects past year avg and avg during each prior decade of life</b></p> <ul style="list-style-type: none"> <li>Never: no alcohol use at all points in time</li> <li>Former light to moderate: ≤60 g/d (men) or ≤30 g/d (women) in the past but not at time of enrollment</li> <li>Former heavy: &gt;60 g/d (men) or &gt;30 g/d (women) in the past but not at time of enrollment</li> <li><b>Light (REF):</b> ≤2 g/d (men) or ≤1 g/d (women)</li> <li>Below rec. limits: ≤24 g/d (men) or ≤12 g/d (women)</li> <li>Light to moderate: ≤60 g/d (men) or ≤30 g/d (women)</li> <li>Occasionally heavy: &gt;60 g/d (men) or &gt;30 g/d (women)</li> <li>Heavy: &gt;60 g/d (men) or &gt;30 g/d (women)</li> </ul>	<p><b>Men</b> Never: NS <b>Former light to mod: 1.32 (1.11, 1.57)</b> <b>Former heavy: 2.00 (1.61, 2.50)</b> Light (REF) <b>Below rec. limits: 0.86 (0.74, 0.99)</b> Light to mod: NS Occasionally heavy: NS <b>Heavy: 1.57 (1.26, 1.96)</b></p> <p><b>Women</b> <b>Never: 1.16 (1.04, 1.30)</b> <b>Former light to mod: 1.23 (1.09, 1.39)</b> <b>Former heavy: 1.74 (1.22, 2.48)</b> Light (REF) <b>Below rec. limits: 0.91 (0.84, 0.99)</b> Light to mod: NS Occasionally heavy: NS Heavy: NS</p>	<p><b>Key confounders:</b> Sex, age, SES (education), eating pattern or diet quality, physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry, hypertension, diabetes</p> <p><b>Other:</b> Center; self-reported cancer, myocardial infarction, or stroke; in women: menopausal status, menopausal hormone use, number of live births</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, alcohol intake</li> <li>Length of follow-up differs among participants</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Bobak, 2016<sup>4</sup></b> Prospective Cohort Study, Health, Alcohol and Psychosocial factors in Eastern Europe Russia, Lithuania, Poland N=34,304  <b>Participants:</b> 45-69y, 53% female, excluded those without linkage to death registry	<b>Baseline:</b> 2002-2005 <b>Follow-up:</b> 2010-2011  <b>Average daily alcohol intake for men/women:</b> <ul style="list-style-type: none"> <li>0 g/d (combines never + former)</li> <li><b>&lt;10/5 g/d, REF</b></li> <li>10-60/5-20 g/d</li> <li>&gt;60/20 g/d</li> </ul> <b>Drinking frequency:</b> <ul style="list-style-type: none"> <li>Never</li> <li><b>&lt;1/mo, REF</b></li> <li>1-3/mo</li> <li>1-4/wk</li> <li>5+/wk</li> </ul> <b>Drinking pattern for men/women:</b> <ul style="list-style-type: none"> <li><b>Light drinker (REF):</b> &lt;2/0.5 drinks per occasion</li> <li>Moderate drinker (&lt;4/2 drinks per occasion)</li> <li>Occasional heavy drinker (&gt;4 drinks per occasion &lt;1/wk)</li> <li>Regular heavy drinker (&gt;4 drinks per &gt;1/wk)</li> </ul> Binge drinking: >100/60 g of ethanol in one episode at least monthly ( <b>non-binge drinker, REF</b> )	Average daily alcohol intake for women (all NS in men): <ul style="list-style-type: none"> <li>&lt;5 g/d, REF</li> <li>5-20 g/d: NS</li> <li><b>&gt;20 g/d: 1.92 (1.25, 2.93)</b></li> </ul> Drinking frequency in men (similar in women): <ul style="list-style-type: none"> <li>&lt;1/mo, REF</li> <li>1-3/mo: NS</li> <li>1-4/wk: NS</li> <li>5+/wk: NS</li> </ul> Drinking pattern in men (similar in women): <ul style="list-style-type: none"> <li>Light drinker, REF</li> <li>Moderate drinker: NS</li> <li>Occasional heavy drinker: NS</li> <li>Regular heavy drinker: NS</li> </ul> Binge drinking in men (similar in women): <ul style="list-style-type: none"> <li>Non-binge drinker, REF</li> <li>Binge-drinker: NS</li> </ul>	<b>Key confounders:</b> Sex, age, SES (education, marital status, economic activity, asset score, subjective hardship score), eating pattern or diet quality, physical activity, smoking  <b>Other factors considered:</b> anthropometry  <b>Other:</b> Marital status, prevalent CVD or cancer, depressive symptoms, study center	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>Older cohort: potential survival bias</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>Lowest level alcohol consumers likely included never and former drinkers</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Britton, 2010<sup>5</sup></b> Prospective Cohort Study, Whitehall II cohort study of British civil servants United Kingdom N= 5,411</p> <p><b>Participants:</b> 35-55y (~43y), 100% male; analysis restricted to those who were free of CHD at baseline and those who attended ≥3 phases to look at variability in alcohol intake; women were excluded since there were too few CHD events; men who reported on ≥3 separate occasions that alcohol frequency in last year was “none” were deemed “never drinkers” and excluded from analysis</p>	<p><b>Baseline:</b> 1985-1988; Follow-up (exp data): Phase 2 (1989-90), Phase 3 (1991-94), Phase 5 (1997-99), and Phase 7 (2003-04); <b>Follow-up</b> (outcome): 2009</p> <p><b>Variability in average alcohol intake (SD of grouped alcohol levels)</b></p> <ul style="list-style-type: none"> <li>• <b>Low (REF):</b> alcohol variability (SD≤0.5)</li> <li>• Medium alcohol variability (0.5&lt;SD≤1.1)</li> <li>• High alcohol variability (SD&gt;1.1)</li> </ul> <p><b>Average alcohol consumption (baseline and average among drinkers)</b></p> <ul style="list-style-type: none"> <li>• None</li> <li>• 1-7 units/wk</li> <li>• <b>Moderate drinkers (REF):</b> 8-14 units/wk</li> <li>• 15-21 units/wk</li> <li>• 22-28 units/wk</li> <li>• 29-35 units/wk</li> <li>• ≥36 units/wk</li> </ul>	<p><u>Variability in intake</u> Low (REF) Medium: 1.00 (0.77, 1.29) <b>High: 1.52 (1.07, 2.17)</b></p> <p><u>Alcohol consumption</u> Figure only; Data NR</p>	<p><b>Key confounders:</b> Sex, age, SES (civil service employment grade of low/med/high), smoking, average alcohol intake during follow-up</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> CHD</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Britton, 2016<sup>6</sup></b> Prospective Cohort Study, Whitehall II cohort study of British civil servants United Kingdom N= 7,010</p> <p><b>Participants:</b> 47-67y, 29% female</p>	<p><b>Baseline:</b> 1997-1999 <b>Follow-up:</b> 2015 (16-18y)</p> <p><u><b>Average intake: reported for previous week by beverage type, converted to g/week</b></u></p> <ul style="list-style-type: none"> <li>• Non-drinker</li> <li>• Former drinker: current non-drinker but indicated alcohol consumption in the past</li> <li>• <b>Moderate (REF):</b> within guidelines of 1–168 g (1–21 units) of ethanol per week for men and 1–112 g (1–14 units) for women)</li> <li>• Heavy: 169–407 g (22–50 units) for men and 113–287 (15–35 units) for women)</li> <li>• Very heavy: 408+g (51+ units) for men and 288+g (36+ units) for women</li> </ul> <p><u>Pre- and post-change conversion:</u> before and after the legal introduction of standardized wine serving sizes in the UK and the shift from 125 to 175 ml glasses becoming the normative standard serving size; 1 glass of wine = 8g ETOH before 16g after</p>	<p><u>Pre-change conversion</u> 1 glass of wine = 8g ETOH:</p> <ul style="list-style-type: none"> <li>• Non-drinker: 1.17 (0.91, 1.51)</li> <li>• Former drinker: 1.22 (0.97, 1.54)</li> <li>• Moderate: REF</li> <li>• Heavy: 1.05 (0.87, 1.26)</li> <li>• <b>Very heavy: 2.26 (1.67, 3.06)</b></li> </ul> <p><u>Post-change conversion</u> 1 glass of wine = 16g ETOH:</p> <ul style="list-style-type: none"> <li>• Non-drinker: 1.14 (0.88, 1.48)</li> <li>• Former drinker: 1.19 (0.94, 1.51)</li> <li>• Moderate: REF</li> <li>• Heavy: 0.90 (0.75, 1.08)</li> <li>• <b>Very heavy: 1.55 (1.23, 1.94)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (civil service employment grade of low/med/high), eating pattern or diet quality, physical activity, smoking</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>



Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Degerud, 2020<sup>7</sup></b>  Prospective Cohort Study, The Age 40 Program; the Cohort of Norway (CONOR)  Norway  N= 243,372</p> <p><b>Participants:</b> ~44y, 52% female, mean BMI 25.7 kg/m<sup>2</sup>; excluded participants with missing or inconsistent survey data on alcohol, mental health, or covariates</p>	<p><b>Baseline:</b> 1994-2002  <b>Follow-up:</b> 2014 (~17y)</p> <p><b>Average intake:</b> combined frequency &amp; amount</p> <ul style="list-style-type: none"> <li>Current abstainers: includes never and former drinkers (data not extracted)</li> <li><b>Low (REF):</b> &lt;2 g/d</li> <li>Light: 2-11.99 g/d</li> <li>Moderate: 12-23.99 g/d</li> <li>High: ≥24 g/d</li> <li>Per 5 g/d increase among current drinkers</li> </ul>	<ul style="list-style-type: none"> <li>Low: REF</li> <li><b>Light: 0.93 (0.90, 0.96), p&lt;0.001</b></li> <li>Moderate: 1.03 (0.97, 1.09), P=0.386</li> <li><b>High: 1.33 (1.19, 1.48), P&lt;0.001</b></li> <li><b>Per 5 g/d increase among current drinkers: 1.03 (1.02, 1.04), P&lt;0.001</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, SES, physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry, diabetes, lipids (total cholesterol concentration, triglyceride concentration), family history of chronic disease (CHD, CVD)</p> <p><b>Other:</b> marital status, resting heart rate, mental health problems</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol pattern</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>Length of follow-up differs among participants</li> <li>Individuals excluded for missing values were on average older, more likely to be male, and more likely to have died from any cause and CVD</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Evans-Polce, 2016<sup>8</sup></b>  Prospective Cohort Study,  National Child Development  Study (NCDS)  United Kingdom  N= 11,469</p> <p><b>Participants:</b> All 33y at baseline;  51% female</p>	<p><b>Baseline:</b> 33y  <b>Follow-up:</b> 51y (18y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• Abstainers: 0 units/wk reported and reported frequency as “never” (combined never + former)</li> <li>• <b>Light drinkers (REF):</b> 1-14 units/week for females; 1-21 units/week for males</li> <li>• Moderate drinkers: 15-28 units/week for females; 22-35 units/week for males</li> <li>• Heavy drinkers: (&gt;28 units/week for females; &gt;35 units/week for males)</li> </ul> <p>Unit = approximately equal to half a pint of beer (284 ml), a small glass of wine (125 ml), or a standard pub measure of distilled spirits (25 ml)</p>	<ul style="list-style-type: none"> <li>• Light drinkers: REF</li> <li>• Moderate drinkers: 1.09 (0.71, 1.68)</li> <li>• <b>Heavy drinkers: 1.90 (1.25, 2.88), P&lt;0.05</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, SES (early life social class, parent education, &amp; home ownership; participant (adult) education &amp; employment status), smoking, alcohol intake (at 16y, 23y)</p> <p><b>Other factors considered:</b> diabetes</p> <p><b>Other:</b> social support, “physical health” (heart problems, epilepsy, asthma, congenital malformations, psychiatric disorders; at 16y, 23y, 33y)</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity, alcohol pattern</li> <li>• Exposure measurements from baseline and earlier ages included, but no additional measurements during follow up</li> <li>• Exposure data measurement tool not validated</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Ferrari, 2014<sup>9</sup></b> Prospective Cohort Study, European Prospective Investigation into Cancer and nutrition (EPIC) Denmark, France, Germany, Greece, Italy, Netherlands, Spain, and United Kingdom N=349,730</p> <p><b>Participants:</b> 25-70y (~52y), 71% female; excluded entire cohorts (Naples, Italy; Bilthoven, Netherlands, Sweden, and Norway) because no information was collected on past alcohol use, and participants with incomplete vital status information, those who had not filled out dietary or lifestyle questionnaires, or whose ratio of energy intake to EER was in top or bottom 1%, those that reported cancer, diabetes, myocardial infarction, heart disease or stroke.</p>	<p><b>Baseline:</b> 1992-2000 <b>Follow-up:</b> 12.6y</p> <p><b>Average intake:</b> Lifetime intake: Avg of weekly consumption at ages 20, 30, 40, &amp; 50y</p> <ul style="list-style-type: none"> <li>Never drinkers (potentially includes never + former)</li> <li><b>Moderate drinkers (REF):</b> 0.1-4.9 g/d</li> <li>5-14.9 g/d</li> <li>15-29.9 g/d</li> <li>Heavy drinkers: ≥30 g/d (women) or 30-59.9 g/d (men)</li> <li>Extreme drinkers: ≥60 g/d (men)</li> </ul>	<p><b>Women</b> 0.1-4.9 g/d (REF) 5-14.9 g/d: NS 15-29.9 g/d: NS <b>≥30 g/d: 1.27 (1.13, 1.43)</b></p> <p><b>Men</b> 0.1-4.9 g/d (REF) <b>5-14.9 g/d: 0.93 (0.87, 0.99)</b> 15-29.9 g/d: NS <b>30-59.9 g/d: 1.15 (1.06, 1.24)</b> <b>≥60 g/d: 1.53 (1.39, 1.68)</b></p>	<p><b>Key confounders:</b> Sex, age, SES, smoking, alcohol pattern</p> <p><b>Other factors considered:</b> total energy intake, anthropometry, diabetes</p> <p><b>Other:</b> Center; in women: menopausal status, ever use of replacement hormones, number of full-term pregnancies</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity</li> <li>Exposure data only measured once</li> <li>Length of follow-up differs among participants</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Gea, 2014<sup>10</sup></b> Prospective Cohort Study, Seguimiento Universidad de Navarra (SUN) Project Spain N=18,394</p> <p><b>Participants:</b> ~44y, 60% female; excluded participants with total daily energy intake out of percentiles 1 and 99</p>	<p><b>Baseline:</b> 2009 <b>Follow-up:</b> ~12y</p> <p><b>Average intake</b> Adherence to Mediterranean alcohol-drinking pattern (MADP):</p> <ul style="list-style-type: none"> <li>Abstainers (combines never + former)</li> <li>Low (0-2 pts)</li> <li>Moderate-low (3-4pts)</li> <li>Moderate-high (5-6pts)</li> <li><b>High (7-9pts, REF)</b></li> </ul> <p><b>Moderate alcohol intake:</b></p> <ul style="list-style-type: none"> <li>Low: 5 g/d women; &lt;10 g/d men</li> <li>Moderate: 5-25 g/d women, 10-50 g/d men</li> <li>High: &gt; 25 g/d women, &gt;50 g/d men</li> </ul>	<p><b>By 2pt increase in MADP score among drinkers: 0.75 (0.62, 0.89)</b></p> <p><b>Adherence to MADP:</b> <b>Low: 3.09 (1.74, 5.50)</b> Mod-low: NS Mod-high: NS High (REF):</p> <p><b>Alcohol intake:</b> Moderate (REF) Low: NS <b>High: 2.07 (1.20, 3.57)</b></p> <p>Proportion of alcohol consumed from spirits or wine was not significantly associated with mortality.</p>	<p><b>Key confounders:</b> Sex, age, eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> total energy intake, anthropometry, hypertension, diabetes, beverage type (e.g., beer, wine, spirits)</p> <p><b>Other:</b> Hypercholesterolemia, cancer, watching television</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, SES</li> <li>Exposure data only measured once</li> <li>Length of follow-up differs among participants</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Goulden, 2016<sup>11</sup></b>  Prospective Cohort Study, Health and Retirement Study (HRS)  United States  N= 24,029</p> <p><b>Participants:</b> ~66y (all &gt;50y), 58% female, 75% white, BMI ~27.4 kg/m<sup>2</sup></p>	<p><b>Baseline:</b> 1995-2004  <b>Follow-up:</b> 2012</p> <p><b>Average intake:</b> (avg from 3 interviews over 4y prior to baseline—each interview reflects avg intake from last 3 months)</p> <ul style="list-style-type: none"> <li>Non-drinker: abstinent over 4y; From data waves 4-7, separated into: <ul style="list-style-type: none"> <li>Lifetime non-drinker</li> <li>Former drinker now abstainer</li> </ul> </li> <li><b>Occasional (REF):</b> reported drinking on at least 1 occasion, but never more than &lt;1/wk</li> <li>&lt;7 drinks/wk</li> <li>7-&lt;14 drinks/wk</li> <li>14-&lt;21 drinks/wk</li> <li>≥21 drinks/wk</li> </ul>	<p><u>Full sample (N=24,029):</u></p> <ul style="list-style-type: none"> <li>Occasional: REF</li> <li>&lt;7 drinks/wk: 1.02 (0.94, 1.11)</li> <li><b>7-&lt;14 drinks/wk: 1.14 (1.02, 1.28)</b></li> <li>14-&lt;21 drinks/wk: 1.13 (0.93, 1.35)</li> <li><b>≥21 drinks/wk: 1.45 (1.16, 1.81)</b></li> </ul> <p><i>Data stratified by age &amp; sex available in paper</i></p> <p><u>Separation of non-drinker (N=8,452):</u></p> <ul style="list-style-type: none"> <li>Lifetime non-drinker: 1.16 (0.95, 1.43)</li> <li><b>Former drinker now abstainer: 1.26 (1.05, 1.53)</b></li> <li>Occasional: ref</li> <li>&lt;7 drinks/wk: 0.99 (0.82, 1.21)</li> <li>7-&lt;14 drinks/wk: 1.21 (0.91, 1.61)</li> <li>14-&lt;21 drinks/wk: 1.30 (0.88, 1.91)</li> <li>≥21 drinks/wk: 1.49 (0.97, 2.29)</li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (income quintile, wealth quintile), physical activity, smoking, alcohol pattern (binge drinking)</p> <p><b>Other factors considered:</b> anthropometry, hypertension, diabetes</p> <p><b>Other:</b> religiosity, self-rated health, frequency of inpatient and emergency department or clinic visits, symptoms (shortness of breath, fatigue, and pain), diagnoses (cancer, lung disease, psychiatric disease, stroke, heart disease, and other diseases), mobility, activities of daily living, instrumental activities of daily living, and cognitive level</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> eating pattern or diet quality</li> <li>Older cohort: Survival bias</li> <li>Exposure data measurement tool not validated</li> <li>Different length of follow up across participants (2-12y)</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Graff-Iversen, 2013<sup>12</sup></b> Prospective Cohort Study Norway N= 12,472</p> <p><b>Participants:</b> 20-62y, 49% female</p>	<p><b>Baseline:</b> 1987-1988 <b>Follow-up:</b> ~20.4y</p> <p><b>Frequency of alcohol use:</b> Abstainers (REF): did not differentiate never from former drinkers, therefore data not extracted</p> <p><b>Binge drinking:</b></p> <ul style="list-style-type: none"> <li>• <b>Not in the last year (REF)</b></li> <li>• A few times last year</li> <li>• 1-3 times/mo</li> <li>• 1-2 times/wk</li> <li>• ≥3 times/wk</li> </ul> <p>Frequency not reported</p>	<p><b>Binge drinking:</b></p> <p><b>MEN:</b></p> <ul style="list-style-type: none"> <li>• Not in the last year: REF</li> <li>• A few times last year: 1.14 (1.00, 1.31)</li> <li>• <b>1-3 times/mo: 1.29 (1.09, 1.52)</b></li> <li>• <b>1-2 times/wk: 1.43 (1.11, 1.84)</b></li> <li>• <b>≥3 times/wk: 2.68 (1.88, 3.82)</b></li> <li>• Frequency not reported: 1.23 (0.95, 1.60)</li> </ul> <p><b>WOMEN:</b></p> <ul style="list-style-type: none"> <li>• Not in the last year: REF</li> <li>• <b>A few times last year: 1.26 (1.06, 1.50)</b></li> <li>• <b>1-3 times/mo: 1.50 (1.06, 2.14)</b></li> <li>• <b>≥1 time/wk: 2.06 (1.17, 3.62)</b></li> <li>• Frequency not reported: 1.16 (0.92, 1.45)</li> </ul>	<p><b>Key confounders:</b> Sex, age, SES, smoking, alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> hypertension, blood pressure, diabetes, medications,</p> <p><b>Other:</b> myocardial infarction, angina, stroke, angina symptoms, systolic blood pressure</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity</li> <li>• Exposure data measurement tool not validated</li> <li>• No preregistered data analysis plan</li> </ul>

**Hartz, 2018<sup>13</sup>**

Prospective Cohort Study,  
National Health Interview Survey  
(NHIS) + Veterans Health  
Administration (VA) clinics

United States

N=340,668 (NHIS), 93,653 (VA)

Participants: NHIS: 47y, 18-85y;  
VA: 53y; 55% female (NHIS),  
11% female (VA)

Excluded: those with missing  
alcohol-related data (NHIS)

**Baseline:** 1997-2009 (NHIS),  
2008 (VA)

**Follow-up:** 2011 (NHIS), 2016  
(VA)

**Pattern of intake + frequency:**

- **Frequency of non-binge drinking associated with lowest risk of ACM: 1-2 drinks ~2-3 times/wk, REF**
  - 3.2 times/wk (NHIS)
  - 2-3 times/wk (VA)
- 1-2 drinks 5 times/wk
- 1-2 drinks 6 times/wk
- 1-2 drinks 7 times/wk
- Once weekly
- Monthly
- Never drinker (NHIS)
- Former drinker (NHIS)
- Binge drinking (≥5 drinks)
  - Continuous, days/mo (NHIS)
  - Categorical (VA)
- Non-drinker (VA): does not distinguish never/former

**NHIS:***Frequency of non-binge drinking*

- 1-2 drinks 3.2/wk, REF
- **Frequency of drinking (non-binge) linear term: 1.04 (1.01, 1.07)**
- **Frequency of drinking (non-binge) cubic term: 1.08 (1.081, 1.082)**
- **Frequency of binge drinking linear term: 1.10 (1.09, 1.10)**
- **1-2 drinks 5/wk: 1.05 (P=0.003)**
- **1-2 drinks 6/wk: 1.12 (P<0.0001)**
- **1-2 drinks 7/wk: 1.26 (P<0.0001)**
- Once/wk: 1.04 (P=0.05)
- **Once/mo: 1.06 (P<0.01)**
- **Never: 1.24 (1.18, 1.29)**
- **Former: 1.14 (1.07, 1.22)**

*Quantity of non-binge drinking*

- 1-2 drinks 3.2/wk, REF
- **3-4 drinks: 1.29 (1.22, 1.37)**

*Frequency of binge drinking*

- 1-2 drinks 3.2/wk, REF
- **1.10 (1.09, 1.10)**

**VA:***Frequency of non-binge drinking*

- 1-2 drinks 2-3 times/wk, REF
- **1-2 drinks ≥4/wk: 1.23 (1.04, 1.45)**
- 1-2 drinks 2-4/mo: NS
- ≤Monthly: NS

*Quantity of non-binge drinking*

- Typically 1-2 drinks/occ, REF
- **Typically 3-4 drinks/occ: 1.81 (1.28, 1.56)**

**Key confounders:** Sex, age, race/ethnicity, SES (NHIS: education, employment status, food stamp eligibility), physical activity (NHIS), smoking (NHIS), alcohol pattern (for alcohol intake), alcohol intake (for alcohol pattern)

**Other factors considered:** age distribution of the study sample, diabetes

**Other:** Region of country, survey administration year, perceived health status, medical comorbidities

- **Key confounders NOT accounted for:** SES (VA), eating pattern or diet quality, physical activity (VA), smoking (VA)
- Exposure data measurement tool not validated (NHIS)
- Exposure data only measured once
- No information on non-completers
- No preregistered data analysis plan

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
		<p>Frequency of binge drinking</p> <ul style="list-style-type: none"> <li>• Never (REF)</li> <li>• <b>Monthly: 1.41 (1.28, 1.56)</b></li> <li>• <b>Weekly: 1.79 (1.57, 2.04)</b></li> <li>• <b>Daily/Almost Daily: 2.63 (2.34, 2.97)</b></li> </ul> <p>Secondary analyses in paper stratified by gender</p>		
<p><b>Holahan, 2010<sup>14</sup></b> Prospective Cohort Study United States N=1,824</p> <p><b>Participants:</b> 61y (55-65y), 37% female, 91% Caucasian Excluded lifetime abstainers and those with missing data</p>	<p><b>Baseline:</b> NR <b>Follow-up:</b> 20y</p> <p><b>Average intake:</b> avg/wk by beverage type + frequency/wk, converted to g/d</p> <ul style="list-style-type: none"> <li>• Abstainers (0 g/d) (this group is 100% former drinkers, lifetime abstainers were excluded)</li> <li>• Light drinkers: consuming &lt;1 drink/d (&gt;0 to &lt;14 g/d)</li> <li>• <b>Moderate drinkers (REF):</b> 1 to &lt;3 drinks/d (14 to &lt;42 g/d)</li> <li>• Heavy drinkers: ≥3 drinks/d (≥42 g/d)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Light: 1.12 (0.94, 1.35), P&lt;0.05</b></li> <li>• Moderate (REF)</li> <li>• <b>Heavy: 1.42 (1.17, 1.73), P&lt;0.001</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, SES, physical activity, smoking, alcohol pattern</p> <p><b>Other factors considered:</b> anthropometry, hypertension, diabetes</p> <p><b>Other:</b> Marital status, Other health factors (cancer, heart problems, stroke, anemia, bronchitis, kidney problems, ulcers), physical ailments, depressive symptoms, avoidance coping, number of close friends, quality of friend support</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality</li> <li>• Older sample: Potential survival bias</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>



Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Holahan, 2015<sup>15</sup></b> Prospective Cohort Study United States N=1,121</p> <p><b>Participants:</b> 61y (55-65y), 36% female, excluded those who had not consumed alcohol in the last year</p>	<p><b>Baseline:</b> NR <b>Follow-up:</b> 20y</p> <p><b>Level and pattern:</b></p> <ul style="list-style-type: none"> <li>• Moderate/regular</li> <li>• Moderate/heavy episodic</li> <li>• High/regular</li> <li>• High/heavy episodic</li> </ul> <p><b>Average</b> level of drinking, men/women:</p> <ul style="list-style-type: none"> <li>• Moderate: 1-2/0.5-1 drink/d</li> <li>• High: &gt;2/&gt;1 drink/d</li> </ul> <p>Pattern of drinking, men/women:</p> <ul style="list-style-type: none"> <li>• Regular: &lt;5/&lt;4 drinks in an occasion</li> <li>• Heavy episodic: ≥5/≥4 drinks in an occasion</li> </ul>	<p><b>Level and pattern:</b></p> <ul style="list-style-type: none"> <li>• High/regular, REF</li> <li>• High/heavy episodic: NS</li> <li>• Moderate/regular, REF</li> <li>• <b>High/regular: 1.73 (1.20, 2.47)</b></li> <li>• Moderate/heavy episodic, REF</li> <li>• High/heavy episodic: NS</li> </ul> <p>Interactions with gender (NS) and baseline medical conditions were examined in the paper (drinking level was more strongly linked to mortality among regular drinkers with fewer medical conditions)</p>	<p><b>Key confounders:</b> Sex, age, SES (family income, education, marital status), physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> medical conditions (cancer, diabetes, heart problems, stroke, high blood pressure, anemia, bronchitis, kidney problems, and ulcers), anthropometry</p> <p><b>Other:</b> Number of close friends, quality of friend support, avoidance coping, depressive symptoms</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality</li> <li>• Older sample: Potential survival bias</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan, but clearly followed preplanned comparisons</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Horvat, 2018<sup>16</sup></b> Retrospective Cohort Study PrivMort Hungary, Russia, Belarus N=124,150</p> <p><b>Participants:</b> 35-69y, 41% female; <b>exclusions:</b> missing data on mortality, drinking frequency, or covariates</p>	<p><b>Baseline:</b> 1998 <b>Follow up:</b> 2013</p> <p><b>Frequency of intake:</b> Reported by female relative</p> <ul style="list-style-type: none"> <li>• Never drank</li> <li>• Former</li> <li>• <b>≤1x/month (REF)</b></li> <li>• 2-4x/month</li> <li>• Several times/wk</li> <li>• Almost every day (men)</li> </ul> <p><b>Pattern of intake:</b> Forms of risky drinking in paper</p> <p>Data on various forms of risky drinking are also available in paper</p>	<p><b>Men</b> ≤1x/month (REF) Never drank</p> <ul style="list-style-type: none"> <li>• <b>Russia: 0.85 (0.78, 0.93)</b></li> <li>• Belarus: 0.92 (0.83, 1.01)</li> <li>• Hungary: 0.93 (0.87, 1.00)</li> </ul> <p>Former</p> <ul style="list-style-type: none"> <li>• <b>Russia: 0.66 (0.61, 0.72)</b></li> <li>• <b>Belarus: 0.59 (0.52, 0.67)</b></li> <li>• <b>Hungary: 0.66 (0.59, 0.74)</b></li> </ul> <p>2-4x/month</p> <ul style="list-style-type: none"> <li>• <b>Russia: 1.38 (1.31, 1.46)</b></li> <li>• <b>Belarus: 1.20 (1.13, 1.27)</b></li> <li>• Hungary: 1.07 (0.99, 1.15)</li> </ul> <p>Several times/wk</p> <ul style="list-style-type: none"> <li>• <b>Russia: 1.85 (1.73, 1.97)</b></li> <li>• <b>Belarus: 1.65 (1.52, 1.80)</b></li> <li>• <b>Hungary: 1.37 (1.27, 1.48)</b></li> </ul> <p>Almost every day</p> <ul style="list-style-type: none"> <li>• <b>Russia: 1.98 (1.81, 2.17)</b></li> <li>• <b>Belarus: 1.85 (1.63, 2.10)</b></li> <li>• <b>Hungary: 1.65 (1.52, 1.79)</b></li> </ul> <p><b>Women</b> ≤1x/month (REF) Never drank</p> <ul style="list-style-type: none"> <li>• <b>Russia: 0.91 (0.85, 0.98)</b></li> <li>• Belarus: 1.03 (0.96, 1.12)</li> <li>• Hungary: 0.96 (0.86, 1.06)</li> </ul> <p>Former</p> <ul style="list-style-type: none"> <li>• <b>Russia: 0.72 (0.57, 0.90)</b></li> <li>• <b>Belarus: 0.63 (0.44, 0.91)</b></li> <li>• <b>Hungary: 0.66 (0.47, 0.92)</b></li> </ul> <p>2-4x/month</p> <ul style="list-style-type: none"> <li>• <b>Russia: 1.42 (1.19, 1.68)</b></li> <li>• Belarus: 1.28 (0.99, 1.66)</li> <li>• <b>Hungary: 1.45 (1.20, 1.75)</b></li> </ul> <p>Several times/wk</p> <ul style="list-style-type: none"> <li>• <b>Russia: 2.70 (2.06, 3.53)</b></li> <li>• <b>Belarus: 4.02 (2.77, 5.84)</b></li> <li>• <b>Hungary: 2.10 (1.71, 2.58)</b></li> </ul> <p>Almost every day: N/A</p>	<p><b>Key confounders:</b> Sex, age, SES (education, marital status), smoking</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> Country of residence, relation to informant, age of informant</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> Race/ethnicity, eating pattern or diet quality, physical activity, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Jackson, 2015<sup>17</sup></b> Prospective Cohort Study, National Health Interview Survey (NHIS) United States N=152,180  <b>Participants:</b> ~45y, 57% female, Adults aged ≥18 years; excluded participants with missing data on alcohol consumption or mortality	<b>Baseline:</b> 1997-2002 <b>Follow-up:</b> 2006 (mean 6.4y)  <b>Pattern of intake + frequency:</b> <ul style="list-style-type: none"> <li>• <b>Never drinkers (REF):</b> ≤12 drinks during lifetime</li> <li>• Former drinkers: ≥12 drinks in lifetime but none during previous year</li> <li>• Infrequent drinkers: ≥12 drinks in lifetime but never ≥12 in a single year</li> <li>• 1-2 drinks ≤2 days/wk (men) or 1 drink ≤2 days/wk (women)</li> <li>• 1-2 drinks 3-7 days/wk (men) or 1 drink 3-7 days/wk (women)</li> <li>• ≥3 drinks ≤2 days/wk (men) or ≥2 drinks ≤2 days/wk (women)</li> <li>• ≥3 drinks 3-7 days/wk (men) or ≥2 drinks 3-7 days/wk (women)</li> </ul>	<b>White Men</b> (n=55,700) Never (REF) <b>Former: 1.17 (1.02, 1.35)</b> Infrequent: NS 1-2 drinks ≤2 d/wk: NS 1-2 drinks 3-7 d/wk: NS <b>≥3 drinks ≤2 d/wk: 0.84 (0.72, 0.97)</b> ≥3 drinks 3-7 d/wk: NS  <b>Black Men</b> (n=9593) Never (REF) Former: NS Infrequent: NS 1-2 drinks ≤2 d/wk: NS 1-2 drinks 3-7 d/wk: NS ≥3 drinks ≤2 d/wk: NS <b>≥3 drinks 3-7 d/wk: 1.44 (1.02, 2.04)</b>  <b>White Women</b> (n=70,669) Never (REF) Former: NS Infrequent: NS <b>1-2 drinks ≤2 d/wk: 0.79 (0.70, 0.88)</b> <b>1-2 drinks 3-7 d/wk: 0.71 (0.61, 0.82)</b> <b>≥3 drinks ≤2 d/wk: 0.62 (0.51, 0.76)</b> ≥3 drinks 3-7 d/wk: NS  <b>Black Women</b> (n=16,218) Never (REF) Former: NS Infrequent: NS 1-2 drinks ≤2 d/wk: NS 1-2 drinks 3-7 d/wk: NS <b>≥3 drinks ≤2 d/wk: 0.53 (0.33, 0.85)</b> ≥3 drinks 3-7 d/wk: NS	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, household income, marital status, poverty status, employment status), physical activity, smoking  <b>Other factors considered:</b> distribution of the study sample, anthropometry  <b>Other:</b> Visited health care professional, self-reported health status	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>

<p><b>Jayasekara, 2015<sup>18</sup></b>  Prospective Cohort Study,  Melbourne Collaborative Cohort  Study (MCCS)  Australia  N=39,577</p> <p>Participants: 40-69y; 60% female  Excluded those with missing  alcohol intake or implausibly high  intake, extreme energy intake, or  missing data for any potential  confounding variable</p>	<p><b>Baseline:</b> 1990-1994  <b>Follow-up:</b> 2008</p> <p><b>Average intake (by beverage  type) + frequency, converted  to g/d</b></p> <ul style="list-style-type: none"> <li>• <b>Lifetime abstainer (REF):</b>  &lt;12 drinks in any past year</li> <li>• &gt;0-9 g/d</li> <li>• 10-19 g/d</li> <li>• 20-39 g/d</li> <li>• 40-59 g/d</li> <li>• 60-79 g/d</li> <li>• ≥80 g/d</li> </ul>	<p><b>Men</b>  <i>Average lifetime intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.83 (0.73, 0.94)  10-19 g/d: 0.87 (0.76, 0.99)  20-39 g/d: 0.81 (0.71, 0.92)  40-59 g/d: NS  60-79 g/d: NS  ≥80 g/d: 1.46 (1.13, 1.88)</p> <p><i>Usual current intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.87 (0.79, 0.97)  10-19 g/d: NS  20-39 g/d: 0.79 (0.70, 0.90)  40-59 g/d: NS  60-79 g/d: NS  ≥80 g/d: NS</p> <p><i>Maximum usual past intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.85 (0.74, 0.97)  10-19 g/d: 0.83 (0.73, 0.96)  20-39 g/d: 0.79 (0.69, 0.90)  40-59 g/d: NS  60-79 g/d: NS  ≥80 g/d: 1.26 (1.08, 1.49)</p> <p><b>Women</b>  <i>Average lifetime intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.85 (0.77, 0.94)  10-19 g/d: NS  20-39 g/d: NS  ≥40 g/d: NS</p> <p><i>Usual current intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.88 (0.79, 0.98)  10-19 g/d: 0.82 (0.70, 0.95)  20-39 g/d: NS  ≥40 g/d: NS</p> <p><i>Maximum usual past intake (g/d)</i>  Abstainer (REF)  &gt;0-9 g/d: 0.86 (0.77, 0.96)  10-19 g/d: 0.85 (0.74, 0.98)  20-39 g/d: NS  ≥40 g/d: NS</p>	<p><b>Key confounders:</b> Sex, age,  race/ethnicity, SES (education,  residential address-based  measure), physical activity,  smoking</p> <p><b>Other factors considered:</b> total  energy intake (from food),  anthropometry</p> <p><b>Other:</b> Household size, fruit and  vegetable consumption,  saturated fat</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT  accounted for:</b> eating pattern  or diet quality, alcohol pattern</li> <li>• Exposure data measurement  tool not validated</li> <li>• Exposure data only measured  once</li> <li>• No preregistered data analysis  plan</li> </ul>
---	--	--	--	---

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Jung, 2012<sup>19</sup></b> Prospective Cohort Study, Korean Multi-center Cancer Cohort (KMCC) Korea N= 16,320  <b>Participants:</b> ≥20y (41% ≥60y), 61% female; excluded those with missing or no information on alcohol consumption habits, and those <20y of age	<b>Baseline:</b> 1993-2004 <b>Follow-up:</b> median 9.3y  <b>Consumption status</b> <ul style="list-style-type: none"> <li>• <b>Never (REF)</b></li> <li>• Past</li> <li>• Current</li> </ul> <b>Average intake (by beverage type) + frequency, converted to g/wk</b> <ul style="list-style-type: none"> <li>• Never drinker</li> <li>• <b>0.01-90 g/wk (REF)</b></li> <li>• 90.01-252 g/wk</li> <li>• 252.0-504 g/wk</li> <li>• &gt;504.01 g/wk</li> </ul>	<b>Consumption status</b> Never (REF) <b>Past: 1.72 (1.38, 2.14)</b> <b>Current: 1.21 (1.06, 1.39)</b>  <b>Average intake (g/wk)</b> 0.01-90 (REF) Never drinker: NS 90.01-252 g/wk: NS 252.01-504 g/wk: NS <b>&gt;504.01 g/wk: 1.39 (1.05, 1.83)</b>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), smoking  <b>Other factors considered:</b> age distribution of the study sample, anthropometry  <b>Other:</b> geographic area	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, physical activity, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• Length of follow-up differs among participants</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>
<b>Kabat, 2015<sup>20</sup></b> Prospective Cohort Study, NIH-AARP Diet and Health Study United States N=476,396  Participants: 50-71y; 40% female; excluded those who had questionnaires completed by proxy respondents, had a history of cancer, died or moved out of study area before study entry, were identified as having cancer through death reports only (n = 2152), had zero follow-up time, had a log-transformed calorie intake of >3 SDs beyond mean, or were missing information needed to construct score (adherence to American Cancer Society guidelines)	<b>Baseline:</b> 1995-1996 <b>Follow-up:</b> 2009 (13.6y)  <b>Average intake:</b> <ul style="list-style-type: none"> <li>• <b>Heavier (REF):</b> ≥3 drinks/d (men); ≥2 drinks/d (women)</li> <li>• Never: nondrinkers (never+former)</li> <li>• Lighter: 1-2 drinks/d (men); 1 drink/d (women)</li> </ul>	<b>Men</b> Heavier (REF) <b>Lighter: 0.91 (0.89, 0.94)</b>  <b>Women</b> Heavier (REF) <b>Lighter: 0.92 (0.89, 0.96)</b>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES, eating pattern or diet quality, physical activity, smoking  <b>Other factors considered:</b> anthropometry  <b>Other:</b> marital status	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> alcohol pattern</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data only measured once</li> <li>• Nondrinker group likely combines never and former drinkers</li> <li>• No information on non-completers</li> <li>• Trial registry does not have data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Kerr, 2011<sup>21</sup></b> Prospective Cohort Study, National Alcohol Surveys (NAS) United States N=10,146  <b>Participants:</b> 42y, 57% female	<b>Baseline:</b> 1984, 1995 <b>Follow-up:</b> 2006 (22y)  <b>Average intake:</b> <ul style="list-style-type: none"> <li>• Ex-drinker</li> <li>• Lifetime abstainer</li> <li>• Occasional drinkers (0-&lt;2/mo)</li> <li>• <b>Moderate drinkers (REF):</b> 2-&lt;30/mo, women; 2-&lt;60/mo, men</li> <li>• Heavy drinkers (30-&lt;120/mo, women, 60-&gt;120/mo, men)</li> <li>• Heavy, ≥120 drinks/mo</li> </ul> Alcohol consumption, <b>pattern:</b> <ul style="list-style-type: none"> <li>• Former drinkers with no alcohol problems</li> <li>• Former drinkers with alcohol problems</li> <li>• Lifetime abstainers</li> <li>• Occasional drinkers, no problems</li> <li>• Occasional drinkers, with problems</li> <li>• <b>Moderate drinkers (REF):</b> no ≥5 drink days</li> <li>• Moderate drinkers, with ≥5 drink/days</li> <li>• Heavy drinkers, no ≥8 drink days</li> <li>• Heavy drinkers, with ≥8 drink days</li> </ul>	<b>Consumption Volume (unadj):</b> <ul style="list-style-type: none"> <li>• Ex-drinker: NS (<b>Black: 0.71 (0.56, 0.91), Hispanic: 1.44 (1.03, 2.01), White: 1.50 (1.20, 1.87)</b>)</li> <li>• Lifetime abstainer: NS</li> <li>• Occasional drinkers: NS</li> <li>• Moderate drinkers (REF)</li> <li>• Heavy drinkers: NS</li> <li>• Heavy drinkers, ≥120/mo: NS</li> </ul> <b>Consumption Pattern (adj):</b> <ul style="list-style-type: none"> <li>• Former drinkers with no alcohol problems: NS (<b>Black: 0.66 (0.53, 0.87); Hispanic: 1.56 (1.07, 2.27); White: 1.59 (1.25, 2.02)</b>)</li> <li>• <b>Former drinkers with alcohol problems: 1.58 (1.21, 2.07) (Black: 1.76 (1.10, 2.93), NS for Hispanic, White)</b></li> <li>• Lifetime abstainers: NS (<b>White: 1.33 (1.06, 1.68)</b>); NS for Hispanic, Black)</li> <li>• Occasional drinkers, no problems: NS (<b>White: 1.39 (1.09, 1.77)</b>); NS for Hispanic, Black)</li> <li>• Occasional drinkers, with problems: NS (<b>White: 4.28 (2.36, 7.79)</b>); NS for Hispanic, Black)</li> <li>• Moderate drinkers, no ≥5 drink days (REF)</li> <li>• <b>Moderate drinkers, with ≥5 drink/days: 1.86 (1.29, 2.69) (Black: 1.94 (1.16, 3.23), NS: for Hispanic, White)</b></li> <li>• Heavy drinkers, no ≥8 drink days: NS</li> <li>• Heavy drinkers, with ≥8 drink days: NS (<b>White: 1.98 (1.15, 3.42)</b>); NS for Hispanic, Black)</li> </ul>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, income), alcohol intake (for pattern exposure), smoking  <b>Other factors considered:</b> N/A  <b>Other:</b> Region of the United States, born in the United States, depression, impulsivity scale, use of illegal drugs or cannabis	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, physical activity</li> <li>• Exposure data only measured once</li> <li>• No information on criteria used to select participants into the analysis or on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Keyes, 2019<sup>22</sup></b> Prospective Cohort Study, Health and Retirement Study United States N=7,904</p> <p><b>Participants:</b> ~61y (all &gt;56y), 54% female, 73% white, 68% overweight/obese</p>	<p><b>Baseline:</b> 1998 <b>Follow-up:</b> 15y</p> <p><b>Average + Pattern</b></p> <ul style="list-style-type: none"> <li>Lifetime abstainers: &lt;12 drinks in lifetime</li> <li>Current abstainers: did not drink in the current wave, but drank in the past or we cannot rule out that they drank in the past</li> <li><b>Occasional drinkers (REF):</b> drinking &lt;1 d/wk (e.g., 1-2/mo), not bingeing &gt; 5 drinks in a single day for men or &gt;4 drinks in a single day for women, and drinking a max of 3 drinks/d for men and a max of 2 drinks/d for women</li> <li>Moderate drinkers: drinking 1 to 2 (women) or 1 to 3 (men) drinks ≥1 d/wk, and not bingeing &gt;5 drinks in a single day for men or &gt;4 drinks in a single day for women</li> <li>Heavy drinkers: men consuming &gt;3 drinks/d or bingeing &gt;5 drinks in 1 day and for women as consuming &gt;2 drinks/day or bingeing &gt;4 drinks in 1 day</li> </ul>	<p><b>Men</b></p> <ul style="list-style-type: none"> <li>Lifetime abstainers: 1.08 (0.87, 1.34)</li> <li><b>Current abstainers: 1.25 (1.02, 1.53)</b></li> <li>Occasional drinkers: REF</li> <li><b>Moderate drinkers: 0.74 (0.60, 0.91)</b></li> <li>Heavy drinkers: 0.97 (0.68, 1.40)</li> </ul> <p><b>Women</b></p> <ul style="list-style-type: none"> <li><b>Lifetime abstainers: 1.35 (1.08, 1.68)</b></li> <li><b>Current abstainers: 1.34 (1.08, 1.67)</b></li> <li>Occasional drinkers: REF</li> <li>Moderate drinkers: 0.82 (0.63, 1.07)</li> <li>Heavy drinkers: 1.07 (0.54, 2.11)</li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, wealth), physical activity, smoking, alcohol pattern (def of exposure included binge drinking)</p> <p><b>Other factors considered:</b> anthropometry, hypertension, blood pressure, diabetes</p> <p><b>Other:</b> depressive symptoms; difficulty in activities of daily living, since previous wave: arthritis, cancer, heart disease, stroke, lung disease, psychiatric problem</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> eating pattern or diet quality</li> <li>Older cohort: Potential survival bias</li> <li>Exposure data measurement tool not validated</li> <li>No preregistered data analysis plan</li> </ul>

**Knott, 2015<sup>23</sup>**

Prospective Cohort Study, Health Survey for England

England

N= 18,368 and 34,523

**Participants:** ≥50y (50-64y: ~53%; ≥65y: ~47%)

~55% female

**Baseline:** 1998-2008

**Follow-up:** 2011 (Median: 6-10y)

**Avg weekly consumption,** 1998-2002, N=18,368:

- **Never drinker (REF):** reported does not drink nowadays and has always been a non-drinker
- Former drinker: reported does not drink nowadays but has not always been a non-drinker
- Not in past 12 mo: reported does drink nowadays but not in past 12 mo
- <1 occasion/mo
- 1-2 occasions/mo
- 0.1–5 units/wk
- 5.1–10 units/wk
- 10.1–15 units/wk
- 15.1–20 units/wk
- >20 units/wk

**Heaviest day** (of the past week), 1999-2008, N=34,523:

- **Never drinker (REF)**
- Former drinker:
- Not in past 12 mo:
- <1 occasion/mo
- 1-2 occasions/mo
- >2 occasions/mo
- 0.1-1.5 units
- 1.6-3.0 units
- 3.1-4.5 units
- >4.5 units

**Men by age: (only significant comparisons reported)**

Men, 50-64y:

Weekly:

- Never drinker: REF
- **15.1–20 units/wk: 0.49 (0.26, 0.91), P=0.03**

Heaviest day:

- Never drinker: REF
- **0.1-1.5 units: 0.43 (0.21, 0.87), P=0.02**

Men, >65y:

Weekly:

- Never drinker: REF
- All NS

Heaviest day:

- Never drinker: REF
- All NS

**Women by age:**

Women, 50-64y:

Weekly:

- Never drinker: REF
- **Former drinker: 1.79 (1.02, 3.16), P=0.04**

Heaviest day:

- Never drinker: REF
- All NS

Women, >65y:

Weekly:

- Never drinker: REF
- **Former drinker: 1.23 (1.01, 1.51), P=0.04**
- **Not in past 12 mo: 1.72 (1.03, 2.88), P=0.04**
- **<1 occ/mo: 0.83 (0.70, 0.99), P=0.04**
- **1-2 occ/mo: 0.74 (0.60, 0.91), P=0.01**
- **0.1-5.0 units/wk: 0.77 (0.64, 0.92), P=0.01**
- **5.1–10 units/wk: 0.77 (0.63, 0.94), P=0.01**

**Key confounders:** Sex, age, race/ethnicity, SES (education, social class, employment status), smoking

**Other factors considered:** anthropometry

**Other:** government office region, marital status

- **Key confounders NOT accounted for:** eating pattern or diet quality, physical activity, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)
- Older cohort: Potential survival bias
- Exposure data measurement tool not validated
- Exposure data only measured once
- Length of follow-up differs among participants
- No information on non-completers
- No preregistered data analysis plan



Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
		<p>Heaviest day:</p> <ul style="list-style-type: none"> <li>• Never drinker: REF</li> <li>• &lt;1 occ/mo: 0.79 (0.66, 0.94), P=0.01</li> <li>• 1-2 occ/mo: 0.70 (0.56, 0.86), P&lt;0.01</li> <li>• 0.1-1.5 units: 0.76 (0.63, 0.91), P&lt;0.01</li> <li>• 1.6-3.0 units: 0.73 (0.60, 0.90), P&lt;0.01</li> <li>• 3.1-4.5 units: 0.58 (0.39, 0.87), P=0.01</li> <li>• &gt;4.5 units: 0.63 (0.42, 0.95), P=0.03</li> </ul> <p>Analyses including those &lt;50y included in supplement document</p>		

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Kunzmann, 2018<sup>24</sup></b> Prospective Cohort Study (secondary analysis of RCT) United States N=99,654  <b>Participants:</b> 55-74y (~65y); 69% female; exclusions: cancer diagnosis or death before exposure assessment, missing data on exposure and covariates	<b>Baseline:</b> Variable (began 1998) <b>Follow up:</b> ~13y / 2009  <b>Average lifetime intake:</b> average of retrospective reports of intake at ages 18-24, 25-39, 40-54, and ≥55y: <ul style="list-style-type: none"> <li>• Never: reported no intake on all assessments</li> <li>• Infrequent: &gt;0-&lt;1 drink/wk</li> <li>• <b>Light (REF):</b> 1-&lt;3 drink/wk</li> <li>• Somewhat light: 3-&lt;5 drink/wk</li> <li>• Light-moderate: 5-&lt;7 drink/d</li> <li>• Moderate: 1-&lt;2 drink/d</li> <li>• Heavy: 2-&lt;3 drink/d</li> <li>• Very heavy: 3+ drink/d</li> </ul> These frequencies take into account seasonal differences by assessing summer separately	<b>Men</b> <ul style="list-style-type: none"> <li>• <b>Never: 1.25 (1.11, 1.40)</b></li> <li>• <b>Infrequent: 1.14 (1.04, 1.24)</b></li> <li>• Light: (REF)</li> <li>• Somewhat light: 0.95 (0.87, 1.05)</li> <li>• Light-moderate: 1.05 (0.95, 1.16)</li> <li>• Moderate: 1.03 (0.95, 1.13)</li> <li>• <b>Heavy: 1.19 (1.07, 1.32)</b></li> <li>• <b>Very heavy: 1.36 (1.23, 1.50)</b></li> </ul> <b>Women</b> <ul style="list-style-type: none"> <li>• <b>Never: 1.29 (1.14, 1.46)</b></li> <li>• <b>Infrequent: 1.23 (1.12, 1.35)</b></li> <li>• Light: (REF)</li> <li>• Somewhat light: 1.01 (0.88, 1.15)</li> <li>• Light-moderate: 1.10 (0.93, 1.30)</li> <li>• Moderate: 1.11 (0.95, 1.30)</li> <li>• <b>Heavy: 1.38 (1.07, 1.78)</b></li> <li>• <b>Very heavy: 1.99 (1.51, 2.64)</b></li> </ul>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, marital status), smoking  <b>Other factors considered:</b> total energy intake, anthropometry  <b>Other:</b> Study center, randomization group, year of exposure assessment, HRT use (women), intake of red meat, processed meat, coffee, fruit, vegetable, fiber, calcium	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> Eating pattern or diet quality, physical activity, alcohol pattern</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data only measured once but does include retrospective data</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Lantz, 2010<sup>25</sup></b> Prospective Cohort Study Americans' Changing Lives (ACL) longitudinal study United States N=3,617</p> <p><b>Participants:</b> ≥25y, 53% female, 84% non-Hispanic White</p>	<p><b>Baseline:</b> 1986, 1989, 1994, 2001/2002 <b>Follow-up:</b> 2005 (~19y)</p> <p><b>Average intake</b> (Drinks past month)</p> <ul style="list-style-type: none"> <li>None/non-drinker (0 drinks in the past month) (cannot distinguish never from former drinkers)</li> <li><b>Moderate drinkers (REF):</b> 1-79 drinks the past month</li> <li>Heavy drinkers (80 or more drinks in the past month)</li> </ul>	<ul style="list-style-type: none"> <li>Moderate (REF)</li> <li>Heavy: 1.13 (0.68, 1.88)</li> </ul> <p>Stratified by age: &lt;55y, 55y+</p> <ul style="list-style-type: none"> <li>&lt;55y, Heavy: 1.16 (0.50, 2.69)</li> <li>+55y, Heavy: 1.07 (0.59, 1.96)</li> </ul>	<p><b>Key confounders:</b> sex, age, race/ethnicity, SES (education, income), physical activity, smoking</p> <p><b>Other factors considered:</b> age distribution of the study sample, anthropometry</p> <p><b>Other:</b> residence (rural, suburban, central city), physical impairment, self-rated health</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern</li> <li>Exposure data measurement tool not validated</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>
<p><b>Li, 2018<sup>26</sup></b> Prospective Cohort Study NHS and HPFS United States N=123,219</p> <p><b>Participants:</b> 47y (women, 30-55y) and 54y (men, 40-75y), 64% female; excluded participants who were underweight, missing data, or had implausible energy intake</p>	<p><b>Baseline:</b> 1980 (women), 1986 (men) <b>Follow-up:</b> 2014 (33.9y (women), 27.2y (men))</p> <p><b>Avg intake</b> across assessments occurring every 4y</p> <ul style="list-style-type: none"> <li>0 g/d alcohol (likely combined never + former, not extracted)</li> <li>1-4.9 g/d alcohol</li> <li><b>5-14.9 g/d alcohol, REF</b></li> <li>15-29.9 g/d alcohol</li> <li>≥30 g/d alcohol</li> </ul>	<ul style="list-style-type: none"> <li>1-4.9 g/d alcohol: 1.03 (1.00, 1.06)</li> <li>5-14.9 g/d alcohol (REF)</li> <li>15-29.9 g/d alcohol: 0.99 (0.96, 1.03)</li> <li><b>≥30 g/d alcohol: 1.25 (1.19, 1.30)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity</p> <p><b>Other factors considered:</b> medications (current multivitamin use; current aspirin use), family history of diabetes mellitus, myocardial infarction, or cancer</p> <p><b>Other:</b> menopausal status and hormone use (women only)</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> SES, eating pattern or diet quality, physical activity, smoking, alcohol pattern</li> <li>Not clear if the "0 g/d" group included both never and former drinkers</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Licaj, 2016<sup>27</sup></b>  Prospective Cohort Study,  Swedish Women's Lifestyle and  Heath cohort  Sweden  N= 33,404</p> <p><b>Participants:</b> ~41y (30-49y),  100% female</p>	<p><b>Baseline:</b> 1991-1992  <b>Follow-up:</b> 2012 (~20y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>0 g/d: abstainers (not extracted because combines never and former drinkers)</li> <li><b>"Light" (REF):</b> 0.1–1.49 g/day</li> <li>1.5–4.9, g/day</li> <li>5–9.9 g/day</li> <li>10–14.9 g/day</li> <li>15+ g/day</li> </ul> <p><b>Change from 1991-2004:</b></p> <ul style="list-style-type: none"> <li><b>Maintain low (REF):</b> &lt;5g/d</li> <li>Stop drinking</li> <li>Start drinking</li> <li>Never at both times</li> <li>Maintain high, ≥5g/d</li> <li>Increased from &lt;5 to ≥5g/d</li> <li>Decreased from ≥5 to &lt;5g/d</li> </ul>	<p><u>Average intake:</u> HR (95% CI)  NS</p> <p><u>Change from 1991-2004:</u></p> <ul style="list-style-type: none"> <li>Maintain low, &lt;5g/d: Ref</li> <li><b>Stop drinking: 1.49 (1.10, 2.02)</b></li> <li>Start drinking: 1.17 (0.92, 1.48)</li> <li><b>Never at both times: 1.53 (1.20, 1.96)</b></li> <li>Maintain high, ≥5g/d: 1.00 (0.84, 1.18)</li> <li>Increased from &lt;5 to ≥5g/d: 0.82 (0.68, 1.00)</li> <li>Decreased from ≥5 to &lt;5g/d: 1.17 (0.89, 1.54)</li> </ul>	<p><b>Key confounders:</b> Sex, age, SES (education), physical activity, smoking</p> <p><b>Other factors considered:</b>  energy intake not from alcohol, anthropometry, family history of chronic disease (breast cancer)</p> <p><b>Other:</b> number of children, age at first birth, oral contraceptive use and duration, menopausal status at baseline and age at menopause</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>Exposure data measurement tool not validated</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Lindshou Hansen, 2011<sup>28</sup></b> Prospective Cohort Study Diet, Cancer, and Health study Denmark N=55,462</p> <p><b>Participants:</b> 50-64y (~56y); 53% female; eligible participants were born in Denmark and had no previous cancers; excluded participants with missing information on relevant questionnaire data, acute coronary syndrome (ACS) diagnosis before baseline or were registered with ACS diagnosis during follow-up, whose medical record could not be retrieved and whose endpoint could not be validated</p>	<p><b>Baseline:</b> 1993-1997 <b>Follow-up:</b> 7.6y</p> <p><b>Average intake:</b> drinks / week</p> <ul style="list-style-type: none"> <li>• &lt;1 (never+former combined)</li> <li>• <b>1-6 (REF)</b></li> <li>• 7-13</li> <li>• 14-20</li> <li>• 21-27</li> <li>• 28-34 (men) or ≥28 (women)</li> <li>• ≥35</li> </ul>	<p>HR reported in figure only and stratified by those with and without hypertension (Figure 2).</p> <p>Text states: "All-cause mortality risk according to alcohol intake was J-shaped in men and women (data not shown). In analyses including hypertension, the association between alcohol and risk of mortality was also J-shaped and the risk among participants with hypertension was higher than for participants without hypertension at all levels of alcohol intake (Figure 2)."</p>	<p><b>Key confounders:</b> Sex, age, SES (education), eating pattern or diet quality, physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry, hypertension, diabetes</p> <p><b>Other:</b> hypercholesterolemia; for women: menopausal status and use of hormone replacement therapy</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, alcohol pattern</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data only measured once</li> <li>• Length of follow-up differs among participants</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>
<p><b>Luksiene, 2017<sup>29</sup></b> Prospective Cohort Study, Multinational Monitoring of Trends and Determinants in Cardiovascular Disease (MONICA) Lithuania N=6,729</p> <p><b>Participants:</b> 35-64y, 52% female; excluded those with incomplete information on analyzed covariates; women not included in multivariate analysis due to small numbers of events, and current smokers and drinkers</p>	<p><b>Baseline:</b> 1983-2002 <b>Follow-Up:</b> 21y</p> <p><b>Average intake</b></p> <ul style="list-style-type: none"> <li>• <b>None (REF):</b> likely combined never + former</li> <li>• <b>1-14 units/wk (REF)</b></li> <li>• &gt;14 units/wk</li> </ul>	<p><b>Men</b> 1-14 units/wk (REF) <b>&gt;14 units/wk: 1.34 (1.13, 1.59)</b></p> <p><b>Women</b> 1-14 units/wk (REF) &gt;14 units/wk: 0.64 (0.24, 1.70)</p> <p><b>Multivariate analysis in men by smoking status are in the paper.</b></p>	<p><b>Key confounders:</b> Sex</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> age, race/ethnicity, SES, eating pattern or diet quality, physical activity, smoking, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Lundin, 2015a</b><sup>30</sup> Prospective Cohort Study, Vietnam Experience Study (VES) United States N= 4,462</p> <p><b>Participants: Vietnam-era Army Vets:</b> mean 38.3y (31.1-49.0y), 0% female; Alcohol use disorder prevalence: 46.7% Inclusion criteria were: (1) entering military service between January 1, 1965 and December 31, 1971; (2) served only one term of enlistment; (3) served at least 16 weeks of active duty; 4) earned a military specialty other than “trainee” or “duty soldier” and at discharge from active duty had a military paygrade no higher than sergeant</p>	<p><b>Baseline:</b> 1985 <b>Follow-up:</b> 2000</p> <p><b>Average intake:</b> <b>1-7 drinks/wk (REF)</b> 8-10 drinks/wk 10-14 drinks/wk 15-28 drinks/wk ≥29 drinks/wk Non-drinkers (does not distinguish between never and former; data not extracted)</p> <p><b>Binge frequency:</b> <b>Drinker, non-binger (REF)</b> 1-3/mo 4-7/mo 8-15/mo ≥16/mo Non-drinkers (does not distinguish between never and former; data not extracted)</p>	<p><i>Average intake:</i> 1-7 drinks/wk: REF All NS</p> <p><i>Binge frequency:</i> Drinker, non-binger: REF All NS</p>	<p><b>Key confounders:</b> Sex, race/ethnicity, SES (education, marital status), smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> depressive disorder, antisocial personality disorder</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> age, eating pattern or diet quality, physical activity</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Lundin, 2015b</b><sup>31</sup> Prospective Cohort Study, REBUS Sweden N= 1,895</p> <p><b>Participants:</b> 18–65y, 18-25y: 14%; 26-35y: 20%; 36-45y: 16%; 46-55y: 22%; 56-65y: 29%; 51% female</p>	<p><b>Baseline:</b> 1969-1970 <b>Follow-up:</b> 2011 (41-42y)</p> <p><b>Average intake: Frequency + amount per occasion</b></p> <ul style="list-style-type: none"> <li>• Lifetime abstainer</li> <li>• Former drinker</li> <li>• <b>Low consumption (REF):</b> 1-49g/wk</li> <li>• Moderate consumption: 50-139 g/wk</li> <li>• High consumption: ≥139 g/wk</li> </ul>	<ul style="list-style-type: none"> <li>• Lifetime abstainer: 0.83 (0.59, 1.18)</li> <li>• <b>Former drinker: 1.86 (1.27, 2.73)</b></li> <li>• Low consumption: REF</li> <li>• Moderate consumption: 0.89 (0.72, 1.10)</li> <li>• High consumption: 1.01 (0.76, 1.35)</li> </ul>	<p><b>Key confounders:</b> Sex, age, SES (social class, school level, marital status); Note: Age controlled for in 5 groups: 18-25, 26-35, 36-45, 46-55, and 56-65 years</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> marital status, depressive neurosis, illicit substance use disorder, psychopathic personality disorder</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>
<p><b>McCullough, 2011</b><sup>32</sup> Prospective Cohort Study, Cancer Prevention Study II (CPS-II) Nutrition Cohort United States N=111,966</p> <p><b>Participants:</b> 50-74y (~63y); 55% female; 98% White; excluded participants from analysis if they reported a history of cancer (except nonmelanoma skin cancer), myocardial infarction, or stroke; an extreme value for height or weight, and below-normal weight (BMI &lt;18.5 kg/m<sup>2</sup>); current smokers and individuals with missing information on smoking, physical activity, and alcohol use or who poorly completed FFQ.</p>	<p><b>Baseline:</b> 1992-1993 <b>Follow-up:</b> 2006 (~14y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• Nondrinker (likely never+former combined)</li> <li>• ≤2 drinks/d (men); ≤1 drink/d (women)</li> <li>• <b>&gt;2 drinks/d (men); &gt;1 drink/d (women) (REF)</b></li> </ul>	<p><u><b>Men</b></u> &gt;2 drinks/d (REF) <b>≤2 drinks/d: 0.86 (0.81, 0.92)</b></p> <p><u><b>Women</b></u> &gt;1 drinks/d (REF) ≤1 drinks/d: 0.99 (0.11, 1.08)</p>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), eating pattern or diet quality, physical activity, smoking</p> <p><b>Other factors considered:</b> total energy intake, anthropometry, medications (nonsteroidal anti-inflammatory drug use)</p> <p><b>Other:</b> multivitamin use, menopausal hormone therapy</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> Alcohol pattern</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Midlov, 2016<sup>33</sup></b> Prospective Cohort Study, Women's Health in Lund Area (WHILA) Sweden N=6,353  <b>Participants:</b> 50-59y, 100% female; excluded those with missing data on explanatory variables	<b>Baseline:</b> 1995-2000 <b>Follow-up:</b> 2015 (~17y)  <b>Average intake:</b> <ul style="list-style-type: none"> <li>None (never+former)</li> <li><b>0.1-11.9 g/d (REF)</b></li> <li>≥12 g/d</li> </ul>	<ul style="list-style-type: none"> <li>0.1-11.9 g/d (REF)</li> <li><b>≥12 g/d: 1.49 (1.18, 1.89)</b></li> </ul>	<b>Key confounders:</b> Sex, age, SES (education, marital status), physical activity, smoking  <b>Other factors considered:</b> anthropometry, diabetes  <b>Other:</b> marital status, ischemic heart disease	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol pattern</li> <li>Older sample: potential survival bias</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>
<b>Muller, 2016<sup>34</sup></b> Prospective Cohort Study, European Prospective Investigation into Cancer and Nutrition (EPIC) Denmark, France, Germany, Greece, Italy, The Netherlands, Spain, Sweden, and the United Kingdom N=246,906  <b>Participants:</b> 50y (40-50y: 25%; 50-60y: 47%; 60-70y: 24%), 65% female, excluded participants who were ≥75 or <40 years old, and those with missing questionnaire information, blood pressure measurements, or follow-up information	<b>Baseline:</b> 1992-2000 <b>Follow-up:</b> 2008-2010 (~11.5y)  <b>Average intake:</b> <ul style="list-style-type: none"> <li>0 drinks/d (never+former)</li> <li>0-0.5 drinks/d</li> <li><b>0.5-1 drinks/d (REF)</b></li> <li>1-2 drinks/d</li> <li>2-6 drinks/d</li> <li>&gt;6 drinks/d (women), 6-10 drinks/d (men)</li> <li>&gt;10 drinks/d (men)</li> </ul>	<b>Total Sample</b> <ul style="list-style-type: none"> <li><b>0-0.5: 1.17 (1.10, 1.25)</b></li> <li>0.5-1 drinks/d (REF)</li> <li>1-2: 0.97 (0.91, 1.04)</li> <li>2-6: 1.07 (1.00, 1.14)</li> <li><b>&gt;6 (women), 6-10 (men): 1.51 (1.38, 1.66)</b></li> <li><b>&gt;10 (men): 2.47 (2.12, 2.88)</b></li> </ul> <b>Men</b> <ul style="list-style-type: none"> <li><b>0-0.5: 1.26 (1.14, 1.38)</b></li> <li>0.5-1 drinks/d (REF)</li> <li>1-2: 0.97 (0.89, 1.07)</li> <li>2-6: 1.04 (0.96, 1.14)</li> <li><b>6-10: 1.48 (1.32, 1.65)</b></li> <li><b>&gt;10: 2.43 (2.07, 2.85)</b></li> </ul> <b>Women</b> <ul style="list-style-type: none"> <li><b>0-0.5: 1.13 (1.04, 1.22)</b></li> <li>0.5-1 drinks/d (REF)</li> <li>1-2: 0.98 (0.90, 1.08)</li> <li><b>2-6: 1.16 (1.05, 1.27)</b></li> <li><b>&gt;6: 1.72 (1.38, 2.15)</b></li> </ul>	<b>Key confounders:</b> Sex, age, eating pattern or diet quality, physical activity, smoking  <b>Other factors considered:</b> anthropometry, blood pressure  <b>Other:</b> Country of recruitment	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, SES, alcohol pattern</li> <li>Exposure data only measured once</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>



Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Ortolá, 2019<sup>35</sup></b> Prospective Cohort Study ENRICA Spain N=3,045  <b>Participants:</b> ≥60y (21% ≥75y), 54% female; excluded participants without information on alcohol consumption, missing data on confounders, and those who died in first year of follow-up	<b>Baseline:</b> 2008-2010 <b>Follow-up:</b> median 7.8y (assessed in 2012, '15, and '17)  <b>Average intake and pattern:</b> <ul style="list-style-type: none"> <li>Continuous (standard units/d)</li> <li><b>Never drinker (REF):</b> 0 g/d</li> <li>Ex-drinker: 0 g/d and answered "I used to drink alcohol, but I quit"</li> <li>Occasional drinker: &gt;0-&lt;1.43 g/d (~1 drink/wk)</li> <li>Light drinker: men ≥1.43-&lt;20 g/d, women ≥1.43-&lt;10 g/d,</li> <li>Moderate drinker: men ≥20-&lt;40 g/d, women ≥10-&lt;24 g/d</li> <li>Heavy/binge drinker: men ≥40 g/d, women ≥24 g/d and/or ≥80g (men) or ≥60g (women) in 1 drinking session during the past 30 days</li> </ul> <b>Lifetime consumption (per decade) was also assessed and used to reclassify consumption category in separate analysis.</b>	<i>By baseline alcohol consumption</i> Never (REF), HR (95% CI) <b>Ex-drinker: 1.49 (1.03, 2.14), P&lt;0.05</b>  Occasional: NS Light: NS Moderate: NS Heavy/binge: NS P for trend (excluding ex-drinkers): NS  <i>By lifetime alcohol consumption</i> Never (REF) Occasional: NS Light: NS Moderate: NS <b>Heavy/binge: 1.85 (1.07, 3.23)</b> P for trend: NS  <i>By lifetime alcohol consumption, with occasional drinkers as reference:</i> NS for all  <i>Change per alcohol standard unit/d:</i> Baseline intake: NS Cumulative intake: NS  <i>By wine consumption among drinkers:</i> NS	<b>Key confounders:</b> Sex, age, SES (education), diet quality (Mediterranean Diet Adherence Screener excluding wine component), physical activity, smoking, alcohol pattern  <b>Other factors considered:</b> anthropometry, hypertension, blood pressure, medications  <b>Other:</b> Time watching TV and other sedentary behaviors, morbidity (diagnosed previously with CVD, cancer, chronic respiratory disease, osteomuscular disease, or depression requiring treatment), self-rated health, functional limitations in IADL	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity</li> <li>Exposure assessed at baseline only but did assess lifetime consumption retrospectively at that point</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Pan, 2019<sup>36</sup></b>  Prospective Cohort Study  Singapore Chinese Health Study (SCHS)  Singapore  N=44,052</p> <p><b>Participants:</b> ~55y (45-74y), 55% female; excluded: participants with a history of invasive cancer (except nonmelanoma skin cancer), coronary heart disease, stroke, diabetes at baseline, whose weight or height data were missing, with extreme energy intakes (&lt;600 or &gt;3,000 kcal for women; &lt;700 or &gt;3,700 kcal for men), and those without information of menopausal status at recruitment</p>	<p><b>Baseline:</b> 1993-1998  <b>Follow up:</b> 12/2016</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• <b>Healthy (REF):</b> light to moderate drinking <ul style="list-style-type: none"> <li>○ Men: &gt;0 to ≤14 drinks/wk</li> <li>○ Women: &gt;0 to ≤7 drinks/wk</li> </ul> </li> <li>• Heavy <ul style="list-style-type: none"> <li>○ Men: &gt;14 drinks/wk</li> <li>○ Women: &gt;7 drinks/wk</li> </ul> </li> <li>• Light to moderate vs. Heavy</li> </ul>	<ul style="list-style-type: none"> <li>• Light to moderate (REF)</li> <li>• <b>Heavy: 1.38 (1.25, 1.53)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, marital status), eating pattern or diet quality (AHEI score), physical activity, smoking</p> <p><b>Other factors considered:</b> total energy intake, age distribution of the study sample, diabetes</p> <p><b>Other:</b> Year of interview, marital status, sleep duration, menopausal status (women)</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> pattern of alcohol intake</li> <li>• Older cohort: Possible survival bias</li> <li>• Exposure data collection tool not validated</li> <li>• Exposure not well-defined: frequency not amount</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Perreault, 2017<sup>37</sup></b> Prospective Cohort Study Health Survey for England (HSE) and Scottish Health Survey (SHS) United Kingdom N=36,370  <b>Participants:</b> ~56y (>40y), NR% female, excluded those with missing data	<b>Baseline:</b> 1994-8-9, 2003-4-6 (HSE), 1998, 2003 (SHS) <b>Follow-up:</b> 2009 (HSE), 2011 (SHS) (353,049 person-years follow-up)  <b>Average intake:</b> <ul style="list-style-type: none"> <li>• <b>Never drunk alcohol, REF</b></li> <li>• Ex-drinkers</li> <li>• Occasional drinkers</li> <li>• Within guidelines (&lt;14 units/wk (women); &lt;21 (men))</li> <li>• Hazardous (14–35 units/wk (women); 21–49 (men))</li> <li>• Harmful (&gt;35 (women); &gt;49 (men))</li> </ul>	<ul style="list-style-type: none"> <li>• Never drunk alcohol, REF: 1.00</li> <li>• <b>Ex-drinkers: 1.37 (1.21, 1.55)</b></li> <li>• <b>Occasional drinkers: 0.83 (0.74, 0.94)</b></li> <li>• <b>Within: 1.13 (1.02, 1.24)</b></li> <li>• <b>Hazardous: 1.19 (1.06, 1.35)</b></li> <li>• <b>Harmful: 1.64 (1.38, 1.95)</b></li> <li>• <b>P&lt;0.001</b></li> </ul> <p>Results were similar, but attenuated among those who meet physical activity requirements</p>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES, physical activity, smoking  <b>Other factors considered:</b> BMI  <b>Other:</b> Psychological distress/depression score, CVD, cancer	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, alcohol intake (for pattern exposure)</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>
<b>Plunk, 2014<sup>38</sup></b> Prospective Cohort Study National Health Interview Survey (NHIS) United States N=128,203  <b>Participants:</b> 18-64y (24% >50y), 48% female	<b>Baseline:</b> 1997-2001 <b>Follow-up:</b> 2006  <b>Pattern of intake:</b> <ul style="list-style-type: none"> <li>• <b>Lifetime abstainers (REF):</b> &lt;12 drinks in lifetime</li> <li>• Past-year abstainer status (12+ drinks in lifetime, but none in past year)</li> <li>• Heavy drinking days (# of days w/ 5+ drinks were consumed)</li> <li>• Non-heavy drinking days (total # of drinking days minus heavy drinking days)</li> </ul>	<p>Lifetime abstainers = REF</p> <p><b>Non-heavy drinking frequency:</b>  <b>B= -0.278, SE= 0.063, P&lt;0.001</b></p> <p><b>Heavy drinking frequency:</b>  <b>B=0.084, SE=0.015, P&lt;0.001</b></p>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES, smoking, alcohol intake (for pattern exposure)  <b>Other factors considered:</b> anthropometry  <b>Other:</b> Survey wave, census region, poor health status	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, physical activity</li> <li>• No information provided regarding criteria used to select participants into the analysis</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Powell, 2020<sup>39</sup></b> Prospective Cohort Study UK Biobank United Kingdom N= 297,988</p> <p><b>Participants:</b> ~56y (40-69y), ~57% female; excluding those with major cardiovascular event or diagnosed with cancer prior to baseline; participants with missing data in covariates included in the analyses or implausible values of sedentary behavior</p>	<p><b>Baseline:</b> 2006-2010 <b>Follow-up:</b> 2016 (6-10y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• <b>Never drinker (REF)</b></li> <li>• Previous-drinkers</li> <li>• Within guidelines: &lt;14 UK units alcohol/wk</li> <li>• Above guidelines: ≥14 and &lt;28 UK units alcohol/wk</li> <li>• Double the guidelines or more: ≥28 UK units alcohol/wk</li> </ul> <p>1 unit = 8g ETOH</p>	<ul style="list-style-type: none"> <li>• Never drinker: REF</li> <li>• <b>Previous drinkers: 1.41 (1.21, 1.63)</b></li> <li>• <b>Within guidelines: 0.83 (0.73, 0.93)</b></li> <li>• Above guidelines: 0.91 (0.80, 1.03)</li> <li>• <b>Double guidelines: 1.22 (1.06, 1.40)</b></li> <li>• <b>P&lt;0.001</b></li> </ul> <p><b>Interactions between alcohol intake and physical activity level are presented in the paper.</b></p>	<p><b>Key confounders:</b> Sex, age, SES (Townsend deprivation index), physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry</p> <p><b>Other:</b> Daily fruit and vegetable consumption, sedentary behavior, baseline hypertension and baseline diabetes</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, smoking, alcohol pattern (for intake exposure)</li> <li>• Exposure data collection methods NR</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>
<p><b>Rhee, 2012<sup>40</sup></b> Prospective Cohort Study Seoul Male Cohort Study Korea N=14,205</p> <p><b>Participants:</b> ~50y (40-59y), 100% male; Criteria for eligible population were males aged 40 to 59 years, residing in or near Seoul, who were beneficiaries of the Korea Medical Insurance Corporation, and who were confirmed to be disease-free at a biennial health checkup conducted in 1990 (excluded those with cancer, CVD, or myocardial infarction)</p>	<p><b>Baseline:</b> 1992 <b>Follow-up:</b> 1993-2008</p> <p><b>Neither average nor pattern</b></p> <ul style="list-style-type: none"> <li>• <b>Never (REF)</b></li> <li>• Former</li> <li>• Current</li> </ul>	<ul style="list-style-type: none"> <li>• Never (REF)</li> <li>• Former: NS</li> <li>• <b>Current: 1.40 (1.07, 1.83)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), physical activity, smoking</p> <p><b>Other factors considered:</b> age distribution of the study sample, anthropometry, hypertension, diabetes</p> <p><b>Other:</b> sleep hours</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• Length of follow-up differs among participants</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Ricci, 2020<sup>41</sup></b> Prospective Cohort Study NHANES United States N=34,672  <b>Participants:</b> ~47y (≥18y), 52% female; excluded participants with missing information on mortality status, alcohol use, and covariates	<b>Baseline:</b> 1999-2014 <b>Follow-up:</b> 2015 (~7.8y)  <b>Average intake, g/d:</b> <ul style="list-style-type: none"> <li>• &lt;0.1</li> <li>• <b>0.1 to &lt;14, REF</b></li> <li>• 14 to &lt;28</li> <li>• ≥28</li> </ul> <b>Pattern of intake, drinks/d:</b> <ul style="list-style-type: none"> <li>• <b>1, REF</b></li> <li>• 2</li> <li>• 3</li> <li>• ≥4</li> </ul> Consuming alcohol in excess of the 2015-2020 DGAs: <u>Men:</u> <ul style="list-style-type: none"> <li>• Mean alcohol intake &gt;28 g/d (2 standard drinks)</li> <li>• &gt;2 drinks during a drinking day</li> <li>• &gt;1 drink during a drinking day</li> </ul> <u>Women:</u> <ul style="list-style-type: none"> <li>• Mean alcohol intake &gt;14 g/d (1 standard drink)</li> <li>• &gt;1 drink during a drinking day</li> </ul>	<b>MEN:</b> Average intake, g/d: <ul style="list-style-type: none"> <li>• <b>&lt;0.1: 1.40 (1.20, 1.63)</b></li> <li>• 0.1 to &lt;14: REF</li> <li>• 14 to &lt;28: 1.17 (0.94, 1.46)</li> <li>• <b>≥28: 1.55 (1.23, 1.96)</b></li> </ul> Pattern of intake, drinks/d: <ul style="list-style-type: none"> <li>• 1: REF</li> <li>• 2: 0.83 (0.69, 1.00)</li> <li>• 3: 0.96 (0.76, 1.21)</li> <li>• <b>≥4: 1.34 (1.18, 1.53)</b></li> </ul> Alcohol intake >28 g/d: <b>HR: 1.87 (1.52, 2.30); PAR: 8.8 (5.5, 12.6)</b>  >2 drinks during a drinking day: <b>HR: 1.39 (1.24, 1.55), PAR: 19.2 (12.8, 25.1)</b>  >1 drink during a drinking day: <b>HR: 1.17 (1.04, 1.33); PAR: 13 (3.4, 22.3)</b>  <b>WOMEN:</b> Average intake, g/d: <ul style="list-style-type: none"> <li>• <b>&lt;0.1: 1.21 (1.01, 1.44)</b></li> <li>• 0.1 to &lt;14: REF</li> <li>• <b>14 to &lt;28: 1.34 (0.95, 1.87)</b></li> <li>• <b>≥28: 1.38 (1.05, 1.83)</b></li> </ul> Pattern of intake, drinks/d: <ul style="list-style-type: none"> <li>• 1: REF</li> <li>• 2: 1.16 (0.93, 1.44)</li> <li>• 3: 1.38 (0.96, 1.98)</li> <li>• <b>≥4: 1.59 (1.38, 1.83)</b></li> </ul> Alcohol intake >14 g/d: <b>HR: 1.47 (1.16, 1.85); PAR: 5.5 (1.9, 9.5)</b>  >1 drink during a drinking day: <b>1.49 (1.30, 1.71); PAR: 25.8 (17.6, 33.5)</b>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)  <b>Other factors considered:</b> Energy intake, BMI, systolic and diastolic blood pressure, CVD (CAD, CHF, MI, stroke), cancer  <b>Other:</b> Dietary fiber, survey	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, physical activity</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Rosella, 2019<sup>42</sup></b> Prospective Cohort Study Canadian Community Health Survey (CCHS) Canada N=149,262</p> <p><b>Participants:</b> 45y (all ≥18y), 54% female; excluded participants &lt;18y at baseline, pregnant women, death before baseline date</p>	<p><b>Baseline:</b> 2000 <b>Follow-up:</b> 2015 (mean ~5.7y)</p> <p><b>Average intake (except 'heavy' category, which combines avg + pattern)</b></p> <ul style="list-style-type: none"> <li>• <b>Light (REF):</b> <ul style="list-style-type: none"> <li>○ Men: 1-3 drinks/wk</li> <li>○ Women: 1-2 drinks/wk</li> </ul> </li> <li>• Moderate: <ul style="list-style-type: none"> <li>○ Men: 4-21 drinks/wk</li> <li>○ Women: 3-14 drinks/wk</li> </ul> </li> <li>• Heavy: <ul style="list-style-type: none"> <li>○ Men: ≥21 drinks/wk or bingeing behavior on weekly basis (≥5 drinks/occasion)</li> <li>○ Women: ≥14 drinks/wk or bingeing behavior on weekly basis (≥5 drinks/occasion)</li> </ul> </li> <li>• Nondrinker: no alcohol in last 12mo or drinks &lt;weekly (combines former &amp; never, so data were not extracted)</li> </ul>	<p><b>All-Cause Mortality</b></p> <p><b>Male:</b> Light (16% of male sample), REF <b>Moderate (25%): 0.85 (0.78, 0.93)</b> Heavy (12%): 1.08 (0.95, 1.23)</p> <p><b>Female:</b> Light (11% of female sample), REF Moderate (17%): 0.94 (0.83, 1.06) <b>Heavy (4%): 1.56 (1.29, 1.90)</b></p> <p><b>Premature Mortality (&lt;75y)</b></p> <p><b>Male:</b> Light = REF Moderate: 0.99 (0.81, 1.22) Heavy: 0.91 (0.73, 1.13)</p> <p><b>Female:</b> Light = REF Moderate: 0.93 (0.71, 1.21) Heavy: 1.05 (0.65, 1.69)</p> <p><b>Data are also presented on amenable vs. nonamenable deaths in the paper.</b></p>	<p><b>Key confounders:</b> Sex, age (ACM analysis only), SES (household income), physical activity, smoking, alcohol pattern</p> <p><b>Other factors considered:</b> ADG score (likely included hypertension, blood pressure, diabetes, glucose, lipids)</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> Race/ethnicity, eating pattern or diet quality</li> <li>• Validity of exposure assessment tool unclear</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Rostron, 2012<sup>43</sup></b> Prospective Cohort Study National Health Interview Survey (NHIS) United States N=237,859  <b>Participants:</b> >18y, NR% female Excluded those with missing data	<b>Baseline:</b> 1997-2004 <b>Follow-up:</b> 2006  <b>Average intake:</b> <ul style="list-style-type: none"> <li>• Never drinker (&lt;12 drinks in life)</li> <li>• Former drinker (0 drinks during last year)</li> <li>• <b>Infrequent drinker (REF):</b> &lt;12 drinks in any year</li> <li>• Light drinker (1 drink per drinking day), occasional or regular</li> <li>• Moderate drinker (2 drinks/d), occasional or regular</li> <li>• Heavy drinker (3+ drinks/d), occasional or regular</li> </ul>	<b>Males</b> <ul style="list-style-type: none"> <li>• Never: NS</li> <li>• <b>Former: 1.24 (1.08, 1.43)</b></li> <li>• Infrequent (REF)</li> <li>• Light: NS</li> <li>• Occasional light: NS</li> <li>• Regular light: NS</li> <li>• Moderate: NS</li> <li>• Occasional moderate: NS</li> <li>• Regular moderate: NS</li> <li>• Heavy: NS</li> <li>• Occasional heavy: NS</li> <li>• <b>Regular heavy: 1.31 (1.12, 1.54)</b></li> </ul> <b>Females</b> <ul style="list-style-type: none"> <li>• <b>Never: 1.34 (1.19, 1.50)</b></li> <li>• <b>Former: 1.40 (1.25, 1.57)</b></li> <li>• Infrequent (REF)</li> <li>• Light: NS</li> <li>• Occasional light: NS</li> <li>• Regular light: NS</li> <li>• Moderate: NS</li> <li>• <b>Occasional moderate: 0.86 (0.74, 0.99)</b></li> <li>• <b>Regular moderate: 1.35 (1.09, 1.67)</b></li> <li>• <b>Heavy: 1.19 (1.02, 1.40)</b></li> <li>• Occasional heavy: NS</li> <li>• <b>Regular heavy: 1.61 (1.29, 2.02)</b></li> </ul>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES, smoking  <b>Other factors considered:</b> health status, anthropometry  <b>Other:</b> N/A	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, physical activity, alcohol intake</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Rundberg, 2014<sup>44</sup></b> Prospective Cohort Study, Women's Health in Lund Area (WHILA) Sweden N=6917  <b>Participants:</b> 50-59y, 100% female	<b>Baseline:</b> 1995-2000 <b>Follow-up:</b> 2009  <b>Average intake:</b> <ul style="list-style-type: none"> <li>Non-drinkers (0g, 0 drinks per wk), combines never + former</li> <li><b>Moderate Drinkers (REF):</b> 1–108 g, &lt;9 drinks/wk</li> <li>Risk drinkers (&gt;109 g, &gt;9 drinks/wk)</li> </ul>	<ul style="list-style-type: none"> <li>Moderate drinker (REF)</li> <li>Risk drinker: NS</li> </ul>	<b>Key confounders:</b> Sex, age, SES, smoking, alcohol intake  <b>Other factors considered:</b> hypertension, diabetes, medications  <b>Other:</b> Social network, household composition, regular medical control for physical or mental disease	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity</li> <li>No information provided regarding criteria used to select participants into the analysis</li> <li>Exposure data measurement tool not validated</li> <li>Not clear if the “non-drinkers” group included both never and former drinkers</li> <li>Exposure data only measured once</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>
<b>Schoenborn, 2014<sup>45</sup></b> Prospective Cohort Study, National Health Interview Survey (NHIS) United States N= 140,741  <b>Participants:</b> adults, current drinkers (participant characteristics NR)	<b>Baseline:</b> 1997-2004 <b>Follow-up:</b> 2006  <b>Current average drinking:</b> <ul style="list-style-type: none"> <li><b>Infrequent/light (REF):</b> ≤3 drinks per week</li> <li>Moderate: 3-7 drinks per week, women; 3-14 drinks per week, men</li> <li>Heavier: &gt;7 drinks per week, women; &gt;14 drinks per week, men</li> </ul> <b>Episodic heavy drinking:</b> <ul style="list-style-type: none"> <li>&lt;3d of 5+ drinks in the past year: REF</li> <li>≥3d of 5+ drinks in past year</li> </ul>	<u>Current average drinking</u> <ul style="list-style-type: none"> <li>Infrequent/light: REF</li> <li>Moderate: 0.94 (0.87, 1.03)</li> <li><b>Heavier: 1.34 (1.17, 1.53), P&lt;0.001</b></li> </ul> <u>Episodic heavy drinking</u> <ul style="list-style-type: none"> <li>&lt;3d of 5+ drinks: REF</li> <li><b>≥3d of 5+ drinks: 1.12 (1.00, 1.26), P&lt;0.05</b></li> <li>Interactions NS</li> </ul>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)  <b>Other factors considered:</b> anthropometry  <b>Other:</b> N/A	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> eating pattern or diet quality</li> <li>Participant characteristics NR</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>Length of follow-up differs among participants</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>



<p><b>Schutte, 2020<sup>46</sup></b>  Prospective Cohort Study  UK Biobank study  United Kingdom  N=446,439  (N=176,950 beer/cider  N=183,930 champagne/white  wine  N=222,192 red wine  N=125,400 spirits)</p> <p><b>Participants:</b> 56y (40-69y), 54% female; excluded participants that had a previous fatal or non-fatal cardiovascular event or previous cancer diagnosis</p>	<p><b>Baseline:</b> 2006-2010  <b>Follow-up:</b> 2015-2016 (~7y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>Continuous (per week)</li> <li>Categorical <ul style="list-style-type: none"> <li><b>Other alcohol drinkers (REF):</b> drinkers that do not drink type of drink stipulated</li> <li>1-7/wk</li> <li>8-14/wk</li> <li>15-21/wk</li> <li>&gt;21/wk</li> </ul> </li> </ul> <p><b>Unit of Measure</b></p> <ul style="list-style-type: none"> <li>Beer/cider: pint (20.17 g ethanol)</li> <li>Champagne/white wine: glass (11.35 g ethanol)</li> <li>Red wine: glass (12.82 g ethanol)</li> <li>Spirits: measures (7.89 g ethanol)</li> </ul>	<p><b>Beer/cider</b>  For number of weekly pints:  <b>1.56 (1.45, 1.68), P&lt;0.0001</b></p> <p>Adj for total baseline alcohol  <b>1.70 (1.53, 1.89), P&lt;0.0001</b></p> <p>Other alcohol drinkers (REF)  1-7/wk: 0.96 (0.90, 1.02)  <b>8-14/wk: 1.26 (1.15, 1.37)</b>  <b>15-21/wk: 1.43 (1.28, 1.60)</b>  <b>&gt;21/wk: 2.19 (1.93, 2.49)</b>  <b>P-trend&lt;0.0001</b>  <i>By weekly pint, risk becomes significantly higher at ≥5 pints/wk (supplemental tables S7 and S8)</i></p> <p><b>Champagne/white wine</b>  For number of weekly glasses:  1.04 (0.95, 1.14)</p> <p>Adj for total baseline alcohol  0.97 (0.87, 1.09)</p> <p>Other alcohol drinkers (REF)  1-7/wk: <b>0.82 (0.78, 0.86)</b>  8-14/wk: <b>0.84 (0.76, 0.94)</b>  15-21/wk: <b>0.80 (0.66, 0.96)</b>  <b>&gt;21/wk: 1.27 (1.03, 1.57)</b>  <b>P-trend&lt;0.0001</b>  <i>By weekly serving, risk becomes significantly lower at 4-5 servings/wk only</i></p> <p><b>Red wine</b>  For number of weekly glasses:  0.95 (0.88, 1.03)</p>	<p><b>Key confounders:</b> Sex, age, SES (Townsend deprivation index), physical activity, smoking, total alcohol intake (g, in analyses labeled 'adj for total baseline alcohol')</p> <p><b>Other factors considered:</b> anthropometry, blood pressure, diabetes, previous cardiovascular event or cancer, beverage type (e.g., beer, wine, spirits)</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>
--	--	--	--	---

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
		<p>Adj for total baseline alcohol  <b>0.81 (0.73, 0.89), P&lt;0.0001</b></p> <p>Other alcohol drinkers (REF)  <b>1-7/wk: 0.71 (0.67, 0.74)</b>  <b>8-14/wk: 0.66 (0.61, 0.72)</b>  <b>15-21/wk: 0.60 (0.52, 0.68)</b>  <b>&gt;21/wk: 0.70 (0.59, 0.84)</b>  <b>P-trend&lt;0.0001</b>  <i>By weekly serving, risk is not significant at any single level</i></p> <p><b><u>Spirits</u></b>  For number of weekly measures:  <b>1.47 (1.35, 1.60), P&lt;0.0001</b></p> <p>Adj for total baseline alcohol  <b>1.47 (1.35, 1.60), P&lt;0.0001</b></p> <p>Other alcohol drinkers (REF)  1-7/wk: 0.96 (0.92, 1.01)  <b>8-14/wk: 1.12 (1.00, 1.25), P&lt;.05</b>  <b>15-21/wk: 1.28 (1.07, 1.53)</b>  <b>&gt;21/wk: 1.78 (1.55, 2.04)</b>  <b>P-trend&lt;0.0001</b>  <i>By weekly serving, risk becomes significantly higher at ≥7 servings/wk</i></p> <p>Data also presented on single increments per week of pints, glasses and measures.</p>		

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Shuval, 2012<sup>47</sup></b> Prospective Cohort Study, Cooper Center Longitudinal Study (CCLS) United States N=29,402  <b>Participants:</b> 43y (20-79y), 0% female; excluded those with diabetes, myocardial infarction, hypertension, stroke, cancer, or with ECG abnormalities, missing data, died during 1 <sup>st</sup> year of f/u	<b>Baseline:</b> 1993-2006 <b>Follow-up:</b> 2011 (~17.4y)  <b>Average intake</b> , drinks/wk: • Nondrinkers (0) (combined never + former) • Light drinkers (<3) • Moderate drinkers (3–14) • <b>Heavy drinkers (REF):</b> >14	• Light: 0.88 (0.75, 1.02) • Moderate: 1.0 (0.89, 1.12) • Heavy (REF) Linear, P-trend=0.1396 Curvilinear, P-trend=0.010	<b>Key confounders:</b> Sex, age, physical activity, smoking  <b>Other factors considered:</b> anthropometry (BMI), hypertension, blood pressure, diabetes, glucose, lipids  <b>Other:</b> Examination year	• <b>Key confounders NOT accounted for:</b> race/ethnicity, SES, eating pattern or diet quality, alcohol intake • Selection into the analysis was based on characteristics potentially associated with the exposure and outcome • Exposure data measurement tool not validated • Exposure data only measured once • Non-drinkers likely included never and former drinkers • No information on non-completers • No preregistered data analysis plan

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Simpson, 2016<sup>48</sup></b>  Prospective Cohort Study  Women's Health Initiative Program (WHI)  United States  N= 145,521</p> <p><b>Participants:</b> 64y (50-79y), 100% female, 2.6% Veterans  Women eligible to participate if they were postmenopausal, planned to remain in the area, did not report having been excessive drinker in past 5 years (i.e., (a) self-evaluated "excessive drinker," (b) had consumed equivalent of a fifth of liquor in 1 day, or (c) had ≥ 2 weeks of drinking ≥7/d), did not report needing to use more drugs to feel desired effects, did not report serious emotional problems that would interfere with study participation, and had an estimated survival of at least 3 years.)</p>	<p><b>Baseline:</b> 1993-1998  <b>Follow-up:</b> 2014 (16-21y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• <b>Lifelong abstainers (REF):</b> consumed alcohol &lt;12 occasions in life</li> <li>• Former drinkers: drank in past but no longer drinking</li> <li>• Infrequent drinkers: &lt;1/wk</li> <li>• <b>Moderate drinkers (REF):</b> 1-7/wk</li> <li>• Moderately heavy drinkers: 8-14/wk</li> <li>• Heavy: ≥15/wk</li> </ul>	<p><u>Veterans (n=3,719)</u></p> <ul style="list-style-type: none"> <li>• Lifelong abstainer (REF)</li> <li>• Former: NS</li> <li>• &lt;1/wk: NS</li> <li>• 1-7/wk: NS</li> <li>• 8-14/wk: NS</li> <li>• ≥15/wk: NS</li> </ul> <p><u>Non-Veterans (n=141,802)</u></p> <ul style="list-style-type: none"> <li>• Lifelong abstainer (REF)</li> <li>• <b>Former: 1.06 (1.02, 1.11)</b></li> <li>• <b>&lt;1/wk: 0.84 (0.80, 0.88)</b></li> <li>• <b>1-7/wk: 0.80 (0.76, 0.83)</b></li> <li>• <b>8-14/wk: 0.79 (0.74, 0.85)</b></li> <li>• <b>≥15/wk: 0.91 (0.85, 0.98)</b></li> </ul> <ul style="list-style-type: none"> <li>• <b>Lifelong abstainer: 1.26 (1.20, 1.32)</b></li> <li>• <b>Former: 1.34 (1.29, 1.39)</b></li> <li>• <b>&lt;1/wk: 1.05 (1.01, 1.09)</b></li> <li>• 1-7/wk (REF)</li> <li>• 8-14/wk: NS</li> <li>• <b>≥15/wk: 1.15 (1.08, 1.23)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (income, education, marital status), physical activity, smoking</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> Veteran status, WHI study arm</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data only measured once</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Skov-Ettrup, 2011<sup>49</sup></b> Prospective Cohort Study Danish National Cohort Study (DANCOS) Denmark N=26,786  <b>Participants:</b> ~45y (19-75y), 51% female; excluded participants diagnosed with ischemic heart disease at baseline, and subjects without information on alcohol intake or covariates were excluded prior to analysis	<b>Baseline:</b> 1994, 2000, 2005 <b>Follow-up:</b> 6.9y  Only moderate and light drinkers were included in this study (≤21 drinks/wk for men; ≤14 drinks/wk for women)  <u>Drinking frequency</u> <ul style="list-style-type: none"> <li>• Nondrinkers (never+former)</li> <li>• <b>1-2 days/wk (REF)</b></li> <li>• 3-4 days/wk</li> <li>• 5-6 days/wk</li> <li>• 7 days/wk</li> </ul> <u>Drinking pattern</u> <ul style="list-style-type: none"> <li>• Nondrinkers (never+former)</li> <li>• <b>Non-binge pattern (REF):</b> ≤5 drinks on all drinking days)</li> <li>• Binge pattern (&gt;5 drinks any drinking day)</li> </ul> 1 Unit = 12g ETOH	<u>Men</u> <i>Drinking frequency</i> 1-2 days/wk, REF 3-4 days/wk: NS 5-6 days/wk: NS 7 days/wk: NS  <i>Drinking pattern</i> <b>Nondrinkers: 1.27 (1.04, 1.56)</b> Non-binge: REF <b>Binge: 0.74 (0.57, 0.97)</b>  <u>Women</u> <i>Drinking frequency</i> 1-2 days/wk, REF 3-4 days/wk: NS 5-6 days/wk: NS 7 days/wk: NS  <i>Drinking pattern</i> Non-binge: REF Binge: NS	<b>Key confounders:</b> Sex, age, SES, physical activity, smoking, alcohol pattern (for alcohol intake), alcohol intake (for alcohol pattern)  <b>Other factors considered:</b> total energy intake, anthropometry, hypertension, diabetes  <b>Other:</b> N/A	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality</li> <li>• Exposure data measurement tool not validated</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<b>Sluik, 2014<sup>50</sup></b> Prospective Cohort Study European Prospective Investigation into Cancer and Nutrition (EPIC) Denmark, Germany, Italy, Netherlands, Spain, Sweden N= 258,911  <b>Participants:</b> ~52y (35-70y), 60% female, BMI ~26 kg/m <sup>2</sup> , 28% current smokers	<b>Baseline:</b> 1992-2000 <b>Follow-up:</b> ~9.9y  <b>Average intake, FFQ:</b> <ul style="list-style-type: none"> <li>0 g/d (does not distinguish between never and former)</li> <li><b>&gt;0-6 g/d (REF)</b></li> <li>&gt;6-12 g/d</li> <li>&gt;12-24 g/d</li> <li>&gt;24-60 g/d</li> <li>&gt;60 g/d</li> </ul>	<ul style="list-style-type: none"> <li>&gt;0-6 g/d: REF</li> <li><b>&gt;6-12 g/d: 0.80 (0.75, 0.85)</b></li> <li><b>&gt;12-24 g/d: 0.79 (0.75, 0.84)</b></li> <li><b>&gt;24-60 g/d: 0.87 (0.82, 0.93)</b></li> <li><b>&gt;60 g/d: 1.13 (1.04, 1.23)</b></li> </ul>	<b>Key confounders:</b> Sex, age, race/ethnicity, SES (education), eating pattern or diet quality, physical activity, smoking  <b>Other factors considered:</b> N/A  <b>Other:</b> study center, self-reported myocardial infarction, stroke or cancer at baseline,	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> alcohol pattern (for intake exposure)</li> <li>Exposure data measurement tool not validated</li> <li>Exposure data only measured once</li> <li>Length of follow-up may differ among participants (outcome assessment time NR)</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Smyth, 2015<sup>51</sup></b>  Prospective Cohort Study  Prospective Urban Rural Epidemiology (PURE) study  12 Countries: Sweden, Canada, Argentina, Brazil, Chile, Poland, South Africa, Turkey, China, Colombia, India, Zimbabwe  N= 114,970</p> <p><b>Participants:</b> ~50y (35-70y), 58% female; Current drinkers (31%): ~50y, 34% female, BMI ~25 kg/m<sup>2</sup></p>	<p><b>Baseline:</b> NR  <b>Follow-up:</b> 2014 (~4.3y)</p> <p><i>Alcohol History:</i></p> <ul style="list-style-type: none"> <li>• <b>Never drinker (REF):</b> self-reported abstinence</li> <li>• Former drinker: no alcohol for ≥1y</li> <li>• Current drinker: within the past year</li> </ul> <p><i>Intake level, current drinkers:</i></p> <ul style="list-style-type: none"> <li>• Low intake: ≤7 drinks/wk</li> <li>• Moderate intake: 7-14 drinks/wk for women; 7-21 drinks/wk for men</li> <li>• High intake: &gt;14 drinks/wk for women; &gt;21 drinks/wk for men</li> </ul> <p><i>Heavy episodic drinking, current drinkers:</i></p> <ul style="list-style-type: none"> <li>• No</li> <li>• Yes: &gt;5 drinks in 1 episode at least 1/mo</li> </ul>	<ul style="list-style-type: none"> <li>• Never drinker: REF</li> <li>• <b>Former drinker: 1.56 (1.27, 1.92)</b></li> <li>• Current drinker: 1.00 (0.87, 1.14)</li> <li>• Low intake: 0.97 (0.84, 1.13)</li> <li>• Moderate intake: 0.97 (0.79, 1.18)</li> <li>• <b>High intake: 1.31, (1.04, 1.66)</b></li> <li>• Not Heavy episodic drinkers: 0.94 (0.80, 1.10)</li> <li>• <b>Heavy episodic drinkers: 1.54 (1.27, 1.87)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, wealth index), eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> anthropometry, hypertension, diabetes, medications,</p> <p><b>Other:</b> jaundice or hepatitis, community-level clustering included as a random effect</p>	<ul style="list-style-type: none"> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• Length of follow-up differs among participants</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Soedamah-Mutha, 2013<sup>52</sup></b> Prospective Cohort Study Health Surveys for England (HSE), Scottish Health Surveys (SHS) England and Scotland N= 17,410</p> <p><b>Participants:</b> ~55y (40-95y), 57% female; excluded those diagnosed with DM, prevalent CVD, those who stopped drinking due to health conditions, and those with missing values for mortality and social class</p>	<p><b>Baseline:</b> HSE: 1997, 1998; SHS: 1998, 2003 <b>Follow-up:</b> 2008</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• None: during past 12 mo (combines never/former drinkers therefore data not extracted)</li> <li>• <b>Moderate (REF):</b> &gt;0 to 21 weekly units for men; &gt;0 to 14 weekly units for women</li> <li>• Moderately high: &gt;21 to 35 weekly units for men; &gt;14 to 21 weekly units for women</li> <li>• High: &gt;35 weekly units for men and &gt;21 weekly units for women</li> </ul> <p>1 unit = 8g ethanol = 125 ml glass of wine = ½ pint regular strength beer</p>	<ul style="list-style-type: none"> <li>• Moderate: REF</li> <li>• Moderately high: 1.08 (0.93, 1.26)</li> <li>• <b>High: 1.26 (1.08, 1.48), P=0.004</b></li> </ul>	<p><b>Key confounders:</b> sex, age, race/ethnicity, SES, physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry (BMI)</p> <p><b>Other:</b> marital status, survey, longstanding illness, changes in alcohol consumption at baseline</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>• Selection into the analysis was based on characteristics potentially associated with the exposure and outcome</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• Length of follow-up differs among participants</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>



Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Stringhini, 2011<sup>53</sup></b> Prospective Cohort Study, Whitehall II and GAZEL United Kingdom and France N=9,771 (Whitehall II) and 17,760 (GAZEL)</p> <p><b>Participants:</b> Whitehall II: 35-55y (~44y), 32% female; excluded those with missing data on health behaviors at baseline and those not followed-up for mortality</p> <p>GAZEL: 35-50y (~43y), 26% female; excluded those with missing data and those that diet before start of follow-up</p>	<p><b>Baseline:</b> Whitehall II: Phase 1, 1985-1988; GAZEL: 1990-1995 <b>Follow-up:</b> Whitehall II: Phase 3 (1991-93), Phase 5 (1997-99), and Phase 7 (2002-04); GAZEL: 1992, 1998, 2004 <b>Follow-up (outcome):</b> Whitehall II: 2009 (~19.5y); GAZEL: 2009 (~16.5y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>Abstainers: 0 units/wk (likely combines never + former)</li> <li><b>Moderate drinkers (REF):</b> 1-21 units/wk (men) or 1-14 units/wk (women)</li> <li>Heavy drinkers: &gt;21 units/wk (men) or &gt;14 units/wk (women)</li> </ul>	<p><u>Whitehall II</u> Moderate (REF) <b>Heavy: 1.25 (1.02, 1.52)</b></p> <p><u>GAZEL</u> Moderate (REF) Heavy: 1.16 (0.99, 1.36)</p> <p>P-interaction between cohort: NS</p>	<p><b>Key confounders:</b> Sex, age</p> <p><b>Other factors considered:</b> N/A</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, SES, eating pattern or diet quality, physical activity, smoking, alcohol pattern</li> <li>Exposure data measurement tool not validated</li> <li>Excluded participants had higher mortality rate and lower SES</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Syden, 2017</b><sup>54</sup></p> <p>Prospective Cohort Study</p> <p>Stockholm Public Health Survey</p> <p>Sweden</p> <p>N= 21,064</p> <p><b>Participants:</b> 46y (25-74y), 55% female, excluded those with diabetes, angina pectoris, infarct, heart failure or stroke at baseline, or missing data</p>	<p><b>Baseline:</b> 2002</p> <p><b>Follow-up:</b> 2007 (5y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• Abstainers (likely never+former)</li> <li>• Light drinkers (0–84 g 100% alcohol/wk)</li> <li>• <b>Moderate drinkers (REF):</b> men: &gt;84–252, women: 84–168 g 100% alcohol/wk</li> <li>• Heavy drinkers (men: 252, women: &gt;168 g 100% alcohol/wk)</li> </ul> <p><b>Pattern of intake:</b> Heavy episodic drinking (HED) (minimum 120g 100% alcohol (e.g. two bottles of wine/6 cans of strong beer/a half bottle of spirits)</p> <ul style="list-style-type: none"> <li>• Abstainers</li> <li>• <b>No HED, REF</b></li> <li>• HED 1–6 times/y</li> <li>• HED 1–3 times/mo</li> <li>• HED&gt;once/wk</li> </ul> <p><b>Average + Pattern</b></p> <ul style="list-style-type: none"> <li>• Abstainers</li> <li>• Light drinkers, no HED</li> <li>• Light drinkers, HED</li> <li>• <b>Moderate drinkers, no HED, REF</b></li> <li>• Moderate drinkers, HED 1–6 times/y</li> <li>• Moderate drinkers, HED&gt;monthly</li> <li>• Heavy drinkers, no HED</li> <li>• Heavy drinkers, HED 1–6 times/y</li> <li>• Heavy drinkers, HED 1–3 times/mo</li> <li>• Heavy drinkers, HED&gt;once/wk</li> </ul>	<p><u>Average intake:</u></p> <ul style="list-style-type: none"> <li>• Light consumers: 1.18 (0.85, 1.63)</li> <li>• Moderate consumers: 1.00 REF</li> <li>• <b>Heavy consumers: 2.21 (1.50, 3.26)</b></li> </ul> <p><u>Pattern: Heavy episodic drinking (HED)</u></p> <ul style="list-style-type: none"> <li>• No HED: 1.00 REF</li> <li>• <b>HED 1–6 times/y: 1.63 (1.09, 2.44)</b></li> <li>• HED 1–3 times/mo: 1.19 (0.70, 2.04)</li> <li>• <b>HED &gt;once/wk: 3.14 (2.03, 4.85)</b></li> </ul> <p><u>Average + Pattern</u></p> <ul style="list-style-type: none"> <li>• Light drinkers, no HED: 1.43 (0.94, 2.18)</li> <li>• <b>Light drinkers, HED: 2.03 (1.16, 3.55)</b></li> <li>• Moderate drinkers, no HED: 1.00 REF</li> <li>• <b>Moderate drinkers, HED 1–6 times/y: 1.85 (1.01, 3.39)</b></li> <li>• <b>Moderate drinkers, HED&gt;mo: 2.21 (1.22, 4.01)</b></li> <li>• <b>Heavy drinkers, no HED: 2.30 (1.23, 4.29)</b></li> <li>• <b>Heavy drinkers, HED 1–6 times/y: 3.93 (1.90, 8.13)</b></li> <li>• <b>Heavy drinkers, HED 1–3 times/mo: 2.48 (1.05, 5.82)</b></li> <li>• <b>Heavy drinkers, HED&gt;once a week 3.80 (1.97, 7.35)</b></li> </ul> <p><b>Findings stratified by socioeconomic position differences provided in paper.</b></p>	<p><b>Key confounders:</b> Sex, age,</p> <p><b>Other factors considered:</b> None</p> <p><b>Other:</b> None</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, SES, eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• Abstainers likely include both never and former drinkers</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<b>Tevik, 2019<sup>55</sup></b> Prospective Cohort Study Nord-Trøndelag Health Study (HUNT) Norway N=7,262  <b>Participants:</b> 74y (all ≥65y); 53% female; excluded participants with missing data on drinking frequency	<b>Baseline:</b> 2006-2008 <b>Follow up:</b> 2013  <b>Average intake:</b> <ul style="list-style-type: none"> <li>• Never</li> <li>• Former (not in past year)</li> <li>• <b>Few times/year (REF)</b> <ul style="list-style-type: none"> <li>• 1x / month</li> <li>• 2-3 days/month</li> <li>• 1 day/wk</li> <li>• 2-3 day/wk</li> <li>• 4-7 day/wk</li> </ul> </li> </ul>	Logistic regression, OR (95% CI) <ul style="list-style-type: none"> <li>• Never: NS</li> <li>• Former: NS</li> <li>• Few times/year (REF) <ul style="list-style-type: none"> <li>• 1x/mo: NS</li> <li>• 2-3 d/mo: NS</li> <li>• 1 d/wk: NS</li> <li>• 2-3 d/wk: NS</li> <li>• 4-7 d/wk: NS</li> </ul> </li> </ul> Stratifying by age group (65-74y and 75y+) did not change results  <b>Sensitivity analyses excluding never and former drinkers did not change findings</b>	<b>Key confounders:</b> Sex, age, SES (education, marital status)  <b>Other factors considered:</b> Age distribution of the study sample, hypertension, blood pressure, diabetes, lipids, overall health status, physical health, mental health  <b>Other:</b> Urban/rural living area	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> Race/ethnicity, eating pattern or diet quality, physical activity, smoking, alcohol pattern (for intake exposure)</li> <li>• Older cohort: Potential survival bias</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

<p><b>van den Brandt, 2020<sup>56</sup></b> Prospective Cohort Study Netherlands Cohort Study (NLCS) The Netherlands N=5,479</p> <p><b>Participants:</b> 68-70y, 53% female, Selected participants were all born 1916-1917; excluded participants with missing data on alcohol intake and covariates</p>	<p><b>Baseline:</b> 1986 <b>Follow-up:</b> 2006-2007</p> <p><b>Average intake</b></p> <ul style="list-style-type: none"> <li>Continuous (per 10 g/d)</li> <li>Categorical <ul style="list-style-type: none"> <li><b>Abstainers (REF)</b></li> <li>Ex-drinker (0 g/d)</li> <li>&gt;0-&lt;5 g/d</li> <li>5-&lt;10 g/d</li> <li>10-&lt;15 g/d</li> <li>15-&lt;30 g/d</li> <li>≥30 g/d</li> </ul> </li> </ul> <p><b>Pattern of alcohol drinking</b></p> <ul style="list-style-type: none"> <li><b>Only at parties (REF)</b></li> <li>Weekend and parties</li> <li>Throughout the week</li> </ul> <p><b>Binge drinking in last 6 months</b></p> <ul style="list-style-type: none"> <li><b>No (REF)</b></li> <li>Yes (&gt;6 drinks/occasion)</li> </ul> <p><b>Frequency of ≥7 drinks/occasion in last 6 months</b></p> <ul style="list-style-type: none"> <li><b>0 times (REF)</b></li> <li>1-&lt;2 times</li> <li>2-&lt;3 times</li> <li>3-&lt;5 times</li> <li>≥5 times</li> </ul>	<p><b>RR (95% CI) for reaching longevity until age 90</b></p> <p><u>Overall</u> <i>Average intake</i> Abstainers (REF) Ex-drinker: 0.84 (0.48, 1.47) &gt;0-&lt;5 g/d: 1.19 (1.07, 1.33) 5-&lt;10 g/d: 1.41 (1.21, 1.63) 10-&lt;15 g/d: 1.30 (1.10, 1.55) 15-&lt;30 g/d: 1.29 (1.10, 1.52) ≥30 g/d: 1.31 (1.06, 1.63) <b>P-trend (excluding ex-drinkers): 0.014</b></p> <p><i>Per 10 g/d (excluding ex-drinkers)</i> <b>1.05 (1.01, 1.09)</b> P-interaction: 0.168</p> <p><b>Stable Subgroup</b> (reported same intake 5y before baseline) <i>Alcohol intake</i> Abstainers (REF) &gt;0-&lt;5 g/d: 1.25 (1.09, 1.43) 5-&lt;10 g/d: 1.42 (1.18, 1.70) 10-&lt;15 g/d: 1.30 (1.05, 1.60) 15-&lt;30 g/d: 1.31 (1.08, 1.59) ≥30 g/d: 1.36 (1.05, 1.76) <b>P-trend (excluding ex-drinkers): 0.024</b></p> <p><i>Per 10 g/d (excluding ex-drinkers)</i> 1.05 (1.00, 1.11) P-interaction: 0.468</p> <p><u>Men</u> <i>Average intake</i> Abstainers (REF) Ex-drinker: 1.49 (0.69, 3.23) &gt;0-&lt;5 g/d: 1.39 (1.01, 1.90) 5-&lt;10 g/d: 1.81 (1.30, 2.53)</p>	<p><b>Key confounders:</b> Sex, age, SES (education), physical activity, smoking, alcohol pattern (for intake exposure), alcohol intake (for pattern exposure)</p> <p><b>Other factors considered:</b> total energy intake, age distribution of the study sample, anthropometry, hypertension, diabetes, beverage type (e.g., beer, wine, spirits)</p> <p><b>Other:</b> myocardial infarction, angina pectoris, stroke, cancer (excluding skin cancer)</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality</li> <li>Exposure data only measured once</li> <li>No preregistered data analysis plan</li> </ul>
--	--	--	--	---

10-<15 g/d: 1.37 (0.95, 1.97)  
15-<30 g/d: 1.43 (1.02, 1.99)  
≥30 g/d: 1.64 (1.15, 2.34)  
**P-trend (excluding ex-drinkers): 0.100**

*Per 10 g/d (excluding ex-drinkers)*  
1.04 (0.98, 1.10)

*Pattern of alcohol drinking*  
Only at parties (REF)  
Weekend and parties: NS  
Throughout the week: NS  
P-trend: NS  
P-interaction with alcohol: NS

*Binge drinking in last 6 months*  
No (REF)  
Yes: NS  
P-interaction with alcohol: NS

*Frequency of ≥7 drinks/occasion in last 6 months*  
0 times (REF)  
1-<2 times: NS  
2-<3 times: NS  
3-<5 times: NS  
≥5 times: NS  
P-trend: NS  
P-interaction with alcohol: NS

### **Women**

*Average intake*  
Abstainers (REF)  
Ex-drinker: 0.62 (0.27, 1.38)  
>0-<5 g/d: 1.17 (1.03, 1.32)  
5-<10 g/d: 1.28 (1.08, 1.52)  
10-<15 g/d: 1.38 (1.13, 1.68)  
15-<30 g/d: 1.31 (1.08, 1.60)  
≥30 g/d: 0.99 (0.69, 1.44)  
P-trend (excluding ex-drinkers): 0.078

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
		<p><i>Per 10 g/d (excluding ex-drinkers)</i> 1.05 (0.99, 1.11)</p> <p><i>Pattern of alcohol drinking</i> Only at parties (REF) Weekend and parties: NS Throughout the week: NS P-trend: NS P-interaction with alcohol: NS</p> <p><i>Binge drinking in last 6 months</i> No (REF) Yes: NS P-interaction with alcohol: NS</p> <p><i>Frequency of <math>\geq 7</math> drinks/occasion in last 6 months</i> 0 times (REF) 1-&lt;2 times: NS 2-&lt;3 times: NS 3-&lt;5 times: NS <math>\geq 5</math> times: NS P-trend: NS P-interaction with alcohol: NS</p> <p><b>Sex-specific data on beer, wine, and liquor available in paper</b></p>		

Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Van Hedel, 2018<sup>57</sup></b>  Prospective Cohort Study  Dutch GLOBE Study  The Netherlands  N=13,034</p> <p><b>Participants:</b> 25-74y, 53% female; excluded participants with chronic obstructive pulmonary disease, heart disease, stroke, renal disease, diabetes, or cancer</p>	<p><b>Baseline:</b> 1991  <b>Follow-up:</b> 2014 (23y)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>No consumption (likely combines never + former)</li> <li><b>Light consumption (1-14 drinks/wk for men, 1-7 drinks/wk for women), REF</b></li> <li>Moderate consumption</li> <li>(15-21 drinks/wk for men, 8-14 drinks/wk for women)</li> <li>Heavy consumption (22+ drinks/wk for men, 15+ drinks/wk for women)</li> </ul>	<ul style="list-style-type: none"> <li>Light = REF</li> <li>Moderate 1.12 (0.94, 1.32)</li> <li><b>Heavy: 1.71 (1.48, 1.98)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, SES (education)</p> <p><b>Other factors considered:</b> None</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> race/ethnicity, eating pattern or diet quality, physical activity, smoking, alcohol intake</li> <li>Exposure data measurement tool likely not validated</li> <li>Exposure data only measured once</li> <li>Not clear if the “no consumption” group included both never and former drinkers</li> <li>No information on non-completers</li> <li>No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Warren Andersen, 2016<sup>58</sup></b>  Prospective Cohort Study,  Southern Community Cohort  Study (SCCS)  United States  N=79,101</p> <p><b>Participants:</b> 51y (40-79y), 57% female; eligible participants spoke English and were not under treatment for cancer within the prior year; excluded those with &lt;1 year of follow-up time or missing data for smoking status, alcohol intake, physical activity, and sedentary behavior</p>	<p><b>Baseline:</b> 2002-2009  <b>Follow-up:</b> 2011</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>Nondrinker (never+former)</li> <li><b>Moderate (REF):</b> &gt;0 but ≤1 drink/d for women or ≤2 for men</li> <li>Heavy</li> </ul>	<p><u>All Participants</u></p> <ul style="list-style-type: none"> <li>Moderate (n=28,155, REF)</li> <li><b>Heavy (n=14075): 1.21 (1.13, 1.29)</b></li> </ul> <p><u>African American Men</u></p> <ul style="list-style-type: none"> <li>Moderate (n=941, REF)</li> <li><b>Heavy (n=875): 1.18 (1.07, 1.30)</b></li> </ul> <p><u>White Men</u></p> <ul style="list-style-type: none"> <li>Moderate (n=341, REF)</li> <li><b>Heavy (n=252): 1.35 (1.14, 1.59)</b></li> </ul> <p><u>African American Women</u></p> <ul style="list-style-type: none"> <li>Moderate (n=541, REF)</li> <li><b>Heavy (n=322): 1.26 (1.09, 1.45)</b></li> </ul> <p><u>White Women</u></p> <ul style="list-style-type: none"> <li>Moderate (n=268, REF)</li> <li>Heavy (n=89): 1.26 (0.99, 1.61)</li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, income, marital status, neighborhood deprivation), eating pattern or diet quality (HEI), physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry</p> <p><b>Other:</b> Enrollment source, sedentary time</p>	<ul style="list-style-type: none"> <li><b>Key confounders NOT accounted for:</b> alcohol pattern</li> <li>Exposure data only measured once</li> <li>Length of follow-up differs among participants</li> <li>No preregistered data analysis plan</li> </ul>



Articles	Time and Exposure Groups	Results ( <i>all results are HR (95% CI) unless otherwise noted</i> )	Confounders accounted for	Limitations
<p><b>Wood, 2018<sup>59</sup></b>  Prospective Cohort Study, Emerging Risk Factors Collaboration (ERFC), EPIC-CVD, UK Biobank  19 “high-income” countries  N=599,912</p> <p><b>Participants:</b> 56y (SD: 9), 21% female, included data from current drinkers in 83 prospective studies in 19 different high-income countries; excluded those with CVD at baseline</p>	<p><b>Baseline:</b> 1964-2008 (ERFC), 1990-2002 (EPIC), 2006-10 (UK Biobank)  <b>Follow-up:</b> 2013, 2009, 2016 (5.4million person years)</p> <p><b>Average intake:</b></p> <ul style="list-style-type: none"> <li>• <b>&gt;0–≤25 g/wk alcohol, REF</b></li> <li>• &gt;25–≤50 g/wk alcohol</li> <li>• &gt;50–≤75 g/wk alcohol</li> <li>• &gt;75–≤100 g/wk alcohol</li> <li>• &gt;100–≤150 g/wk alcohol</li> <li>• &gt;150–≤250 g/wk alcohol</li> <li>• &gt;250–≤350 g/wk alcohol</li> <li>• &gt;350 g/wk alcohol</li> </ul>	<p>Higher alcohol intake was associated with higher risk of ACM, with the lowest risk for &lt;100 g/wk, data not reported (Figure Only)</p> <p>Associations were similar for men and women</p>	<p><b>Key confounders:</b> Sex, age, smoking</p> <p><b>Other factors considered:</b> diabetes</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> race/ethnicity, SES, eating pattern or diet quality, physical activity, alcohol pattern</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Articles	Time and Exposure Groups	Results (all results are HR (95% CI) unless otherwise noted)	Confounders accounted for	Limitations
<p><b>Xi, 2017<sup>60</sup></b>  Prospective Cohort Study  National Health Interview Surveys  United States  N=34,754</p> <p><b>Participants:</b> &gt;18y (40% 18-39y, 37% 40-59y, 22% &gt;60y), 51% female; excluded those with missing data, who were pregnant, or who died in the first 2 years of follow up</p>	<p><b>Baseline:</b> 1997-2009  <b>Follow-up:</b> 2011 (8.2y)</p> <p><b>Average intake: frequency + pattern</b></p> <ul style="list-style-type: none"> <li>• <b>Lifetime abstainers (REF):</b> &lt;12 drinks in one's lifetime</li> <li>• Lifetime infrequent drinkers (&gt;12 lifetime drinks, &lt;12 drinks in last year)</li> <li>• Former-drinkers (&lt;12 drinks in last year)</li> <li>• Current light drinkers (&lt;3 drinks wk)</li> <li>• Current moderate drinkers (&gt;3 and &lt;14 for men and &lt;7 for women drinks/wk)</li> <li>• Current heavy drinkers (&gt;14 drinks/wk for men or &gt;7 for women)</li> </ul> <p>Binge-drinking: 5+ drinks/d</p>	<p><u>Average intake</u></p> <ul style="list-style-type: none"> <li>• Lifetime abstainers, REF</li> <li>• Lifetime infrequent drinkers: 1.02 (0.98, 1.06)</li> <li>• Former-drinkers: 1.04 (0.99, 1.09)</li> <li>• <b>Current light drinkers: 0.80 (0.77, 0.93)</b></li> <li>• <b>Current moderate drinkers: 0.81 (0.77, 0.85)</b></li> <li>• <b>Current heavy drinkers: 1.13 (1.05, 1.21)</b></li> </ul> <p><u>Binge-Drinking Status:</u></p> <ul style="list-style-type: none"> <li>• Lifetime abstainer, REF</li> <li>• <b>Drinker without binge-drinking: 0.81 (0.78, 0.85)</b></li> <li>• <b>Drinker with binge-drinking &lt;1/mo: 0.87 (0.80, 0.95)</b></li> <li>• Drinker with binge-drinking &lt;1d/wk: 0.94 (0.82, 1.07)</li> <li>• <b>Drinker with binge-drinking &gt;1d/wk: 1.16 (1.07, 1.27)</b></li> </ul>	<p><b>Key confounders:</b> Sex, age, race/ethnicity, SES (education, marital status), eating pattern or diet quality, physical activity, smoking</p> <p><b>Other factors considered:</b> anthropometry, hypertension, heart disease, stroke, cancer, diabetes</p> <p><b>Other:</b> N/A</p>	<ul style="list-style-type: none"> <li>• <b>Key confounders NOT accounted for:</b> eating pattern or diet quality, alcohol intake</li> <li>• Exposure data measurement tool not validated</li> <li>• Exposure data only measured once</li> <li>• No information on non-completers</li> <li>• No preregistered data analysis plan</li> </ul>

Table 2: Risk of bias for observational studies examining alcohol consumption and all-cause mortality<sup>vi,vii,viii</sup>

	Confounding	Selection of participants	Classification of exposures	Deviations from intended exposures	Missing data	Outcome measurement	Selection of the reported result
Almeida, 2017, <sup>1</sup> MR	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Bellavia, 2014, <sup>2</sup> PCS	Serious	Moderate	Low	Moderate	Moderate	Low	Moderate
Bergmann, 2013, <sup>3</sup> PCS	Serious	Moderate	Low	Moderate	Moderate	Low	Moderate
Bobak, 2016, <sup>4</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
Britton, 2010, <sup>5</sup> PCS	Serious	Low	Moderate	Low	Moderate	Low	Moderate
Britton, 2016, <sup>6</sup> PCS	Serious	Low	Moderate	Moderate	Low	Low	Moderate
Degerud, 2020, <sup>7</sup> PCS	Serious	Moderate	Moderate	Moderate	Low	Low	Moderate
Evans-Polce, 2016, <sup>8</sup> PCS	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Ferrari, 2014, <sup>9</sup> PCS	Serious	Moderate	Low	Moderate	Moderate	Low	Moderate
Gea, 2014, <sup>10</sup> PCS	Serious	Moderate	Low	Moderate	Low	Low	Moderate
Goulden, 2016, <sup>11</sup> PCS	Serious	Moderate	Moderate	Low	Low	Low	Moderate
Graff-Iversen, 2013, <sup>12</sup> PCS	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Hartz, 2018, <sup>13</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Holahan, 2010, <sup>14</sup> PCS	Serious	Low	Moderate	Moderate	Low	Low	Moderate
Holahan, 2015, <sup>15</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Horvat, 2018, <sup>16</sup> RCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Jackson, 2015, <sup>17</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
Jayasekara, <sup>18</sup> 2015	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Jung, 2012, <sup>19</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Kabat, 2015, <sup>20</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Kerr, 2011, <sup>21</sup> PCS	Serious	Moderate	Low	Moderate	No information	Low	Serious
Keyes, 2019, <sup>22</sup> PCS	Serious	Moderate	Moderate	Low	Low	Low	Moderate
Knott, 2015, <sup>23</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Kunzmann, 2018, <sup>24</sup> PCS	Serious	Serious	Low	Moderate	Moderate	Low	Moderate

<sup>vi</sup> A detailed description of the methodology used for assessing risk of bias is available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews> and in Part C of the following reference: Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.

<sup>vii</sup> Possible ratings of low, moderate, serious, critical, or no information determined using the "Risk of Bias for Nutrition Observational Studies" tool (RoB-NObs) (Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.)

<sup>viii</sup> Abbreviations: MR, Mendelian randomization; PCS, prospective cohort study; RCS, retrospective cohort study

Lantz, 2010, <sup>25</sup> PCS	Serious	Low	Serious	Low	Moderate	Low	Moderate
Li, 2018, <sup>26</sup> PCS	Serious	Moderate	Low	Moderate	Moderate	Low	Moderate
Licaj, 2016, <sup>27</sup> PCS	Serious	Low	Moderate	Low	Moderate	Low	Moderate
Lindschou, 2011, <sup>28</sup> PCS	Serious	Serious	Low	Moderate	Moderate	Low	Moderate
Luksiene, 2017, <sup>29</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Lundin, 2015a, <sup>30</sup> PCS	Serious	Low	Moderate	Moderate	Low	Low	Moderate
Lundin, 2015b, <sup>31</sup> PCS	Serious	Low	Moderate	Moderate	Low	Low	Moderate
McCullough, 2011, <sup>32</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Midlov, 2016, <sup>33</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Muller, 2016, <sup>34</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
Ortola, 2019, <sup>35</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
Pan, 2019, <sup>36</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Perreault, 2017, <sup>37</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Plunk, 2014, <sup>38</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Powell, 2020, <sup>39</sup> PCS	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Rhee, 2012, <sup>40</sup> PCS	Serious	Serious	Serious	Moderate	Moderate	Low	Moderate
Ricci, 2020, <sup>41</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
Rosella, 2019, <sup>42</sup> PCS	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Rostron, 2012, <sup>43</sup> PCS	Serious	Low	Moderate	Moderate	Moderate	Low	Moderate
Rundberg, 2014, <sup>44</sup> PCS	Serious	Serious	Moderate	Moderate	Serious	Low	Moderate
Schoenborn, 2014, <sup>45</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Schutte, 2020, <sup>46</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Shuval, 2012, <sup>47</sup> PCS	Serious	Moderate	Serious	Moderate	Moderate	Low	Moderate
Simpson, 2016, <sup>48</sup> PCS	Serious	Moderate	Low	Moderate	Low	Low	Moderate
Skov-Ettrup, 2011, <sup>49</sup> PCS	Serious	Moderate	Moderate	Low	Moderate	Low	Moderate
Sluik, 2014, <sup>50</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Smyth, 2015, <sup>51</sup> PCS	Serious	Moderate	Moderate	Moderate	Low	Low	Moderate
Soedamah-Muthu, 2013, <sup>52</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
Stringhini, 2011, <sup>53</sup> PCS	Serious	Low	Moderate	Low	Moderate	Low	Moderate
Syden, 2017, <sup>54</sup> PCS	Serious	Serious	Moderate	Moderate	Moderate	Low	Moderate
Tevik, 2019, <sup>55</sup> PCS	Serious	Moderate	Moderate	Moderate	Moderate	Low	Moderate
van den Brandt, 2020, <sup>56</sup> PCS	Serious	Low	Low	Moderate	Moderate	Low	Moderate
van Hedel, 2018, <sup>57</sup> PCS	Serious	Serious	Serious	Serious	Low	Low	Moderate
Warren Andersen, 2016, <sup>58</sup> PCS	Serious	Moderate	Low	Moderate	Low	Low	Moderate
Wood, 2018, <sup>59</sup> PCS	Serious	Serious	Low	Serious	Serious	Low	Moderate
Xi, 2017, <sup>60</sup> PCS	Serious	Moderate	Moderate	Serious	Moderate	Low	Moderate

## METHODOLOGY

---

The NESR team used its rigorous, protocol-driven methodology to support the 2020 Dietary Guidelines Advisory Committee in conducting this systematic review.

NESR's systematic review methodology involves:

- Developing a protocol,
- Searching for and selecting studies,
- Extracting data from and assessing the risk of bias of each included study,
- Synthesizing the evidence,
- Developing conclusion statements,
- Grading the evidence underlying the conclusion statements, and
- Recommending future research.

A detailed description of the methodology used in conducting this systematic review is available on the NESR website: <https://nesr.usda.gov/2020-dietary-guidelines-advisory-committee-systematic-reviews>, and can be found in the 2020 Dietary Guidelines Advisory Committee Report, Part C: Methodology.<sup>ix</sup> This systematic review was peer reviewed by Federal scientists, and information about the peer review process can also be found in the Committee's Report, Part C. Methodology. Additional information about this systematic review, including a description of and rationale for any modifications made to the protocol can be found in the in the 2020 Dietary Guidelines Advisory Committee Report, Chapter 11. Alcoholic Beverages.

Below are details of the final protocol for the systematic review described herein, including the:

- Analytic framework
- Literature search and screening plan
- Literature search and screening results


---

<sup>ix</sup> Dietary Guidelines Advisory Committee. 2020. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC.

# ANALYTIC FRAMEWORK

The analytic framework (**Figure 1**) illustrates the overall scope of the systematic review, including the population, the interventions and/or exposures, comparators, and outcomes of interest. It also includes definitions of key terms and identifies key confounders considered in the systematic review. The inclusion and exclusion criteria that follow provide additional information about how parts of the analytic framework were defined and operationalized for the review.

**Figure 1: Analytic framework**



# LITERATURE SEARCH AND SCREENING PLAN

## Inclusion and exclusion criteria

**Table 3** provides the inclusion and exclusion criteria for the systematic review. The inclusion and exclusion criteria are a set of characteristics used to determine which articles identified in the literature search were included in or excluded from the systematic review.

**Table 3: Inclusion and exclusion criteria**

Category	Inclusion Criteria	Exclusion Criteria
<b>Study design</b>	<ul style="list-style-type: none"> <li>Randomized controlled trials</li> <li>Mendelian randomization studies</li> <li>Non-randomized controlled trials including quasi-experimental and controlled before-and-after studies</li> <li>Prospective cohort studies</li> <li>Retrospective cohort studies</li> <li>Nested case-control studies</li> </ul>	<ul style="list-style-type: none"> <li>Uncontrolled trials</li> <li>Case-control studies</li> <li>Cross-sectional studies</li> <li>Uncontrolled before-and-after studies</li> <li>Narrative reviews</li> <li>Systematic reviews</li> <li>Meta-analyses</li> </ul>
<b>Sample size</b>	<ul style="list-style-type: none"> <li>Observational studies enrolling &gt;1,000 participants</li> </ul>	<ul style="list-style-type: none"> <li>Observational studies enrolling &lt;1,000 participants</li> </ul>
<b>Intervention/exposure</b>	<p>Average consumption of alcoholic beverages</p> <p>Per occasion consumption of alcohol beverages (i.e., number of drinks per drinking day or drinks per drinking occasion)</p> <ul style="list-style-type: none"> <li>Information on type of beverage (e.g., beer, wine, spirits) will be collected if available</li> </ul>	<ul style="list-style-type: none"> <li>Data on 'non-drinker' groups where never and former drinkers are combined</li> </ul>
<b>Comparator</b>	<p><u>Primary</u></p> <ul style="list-style-type: none"> <li>Comparisons across different average alcohol consumption or pattern of alcohol consumption among current drinkers</li> </ul> <p><u>Secondary</u></p> <ul style="list-style-type: none"> <li>Comparisons between never drinkers and current drinkers</li> </ul>	<p>No comparator</p> <ul style="list-style-type: none"> <li>Comparisons with never and former drinkers as a combined 'non-drinker' group</li> </ul>
<b>Outcomes</b>	<ul style="list-style-type: none"> <li>All-cause mortality (i.e. total mortality): the total number of deaths from all causes during a specific time period.</li> <li>(Ideally stratified by sex)</li> </ul>	<ul style="list-style-type: none"> <li>Studies that only report cause-specific mortality (total number of deaths from a specific disease, such as cardiovascular disease or cancer)</li> </ul>

Category	Inclusion Criteria	Exclusion Criteria
<b>Date of publication</b>	<ul style="list-style-type: none"> <li>January 2010 – March 2020</li> </ul>	<ul style="list-style-type: none"> <li>Articles published prior to January 2010</li> </ul>
<b>Publication status</b>	<ul style="list-style-type: none"> <li>Articles published in peer-reviewed journals</li> </ul>	<ul style="list-style-type: none"> <li>Articles not published in peer-reviewed journals, including unpublished data, manuscripts, reports, abstracts, pre-prints, and conference proceedings</li> </ul>
<b>Language of publication</b>	<ul style="list-style-type: none"> <li>Articles published in English</li> </ul>	<ul style="list-style-type: none"> <li>Articles published in languages other than English</li> </ul>
<b>Country<sup>x</sup></b>	<ul style="list-style-type: none"> <li>Studies conducted in Very High or High Human Development Countries</li> </ul>	<ul style="list-style-type: none"> <li>Studies conducted in Medium or lower Human Development Countries</li> </ul>
<b>Study participants</b>	<ul style="list-style-type: none"> <li>Human participants</li> <li>Males</li> <li>Females</li> </ul>	<ul style="list-style-type: none"> <li>Animal subjects</li> <li>Hospitalized samples</li> </ul>
<b>Age of study participants</b>	<ul style="list-style-type: none"> <li>Primarily adults 21y and older</li> <li>Studies that enroll <i>some</i> participants under 21 years old</li> </ul>	<ul style="list-style-type: none"> <li>Studies that <i>exclusively</i> enroll participants under 21 years old</li> </ul>
<b>Health status of study participants</b>	<ul style="list-style-type: none"> <li>Studies that enroll participants who are healthy and/or at risk for chronic disease, including those with obesity</li> <li>Studies that enroll <i>some</i> participants diagnosed with a disease</li> </ul>	<ul style="list-style-type: none"> <li>Studies that <i>exclusively</i> enroll participants diagnosed with a disease or hospitalized with an illness or injury. (For this criterion, studies that exclusively enroll subjects with obesity will not be excluded)</li> </ul>

<sup>x</sup> The Human Development classification was based on the Human Development Index (HDI) ranking from the year the study intervention occurred or data were collected (UN Development Program. HDI 1990-2017 HDRO calculations based on data from UNDESA (2017a), UNESCO Institute for Statistics (2018), United Nations Statistics Division (2018b), World Bank (2018b), Barro and Lee (2016) and IMF (2018). Available from: <http://hdr.undp.org/en/data>). If the study did not report the year in which the intervention occurred or data were collected, the HDI classification for the year of publication was applied. HDI values are available from 1980, and then from 1990 to present. If a study was conducted prior to 1990, the HDI classification from 1990 was applied. If a study was conducted in 2018 or 2019, the most current HDI classification was applied. When a country was not included in the HDI ranking, the current country classification from the World Bank was used instead (The World Bank. World Bank country and lending groups. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-country-and-lending-groups>).



## Electronic databases and search terms

Listed below are the databases searched to identify all potentially relevant articles that have been published to address the update to the existing systematic review.

### PubMed

- Provider: U.S. National Library of Medicine
- Date(s) Searched: March 11, 2020
- Date range searched: January 1, 2000-March 11, 2020
- Search Terms:

**#1** - "Alcohol-Induced Disorders"[Mesh] OR alcohol intake\*[tiab] OR alcohol consum\*[tiab] OR "Alcohol Drinking"[Mesh] OR alcohol drink\*[tiab] OR "Alcoholic Beverages"[Mesh] OR alcoholic beverage\*[tiab] OR beer\*[tiab] OR wine\*[tiab] OR liquor[tiab] OR liqueur\*[tiab] OR spirits[tiab] OR "Ethanol"[Mesh] OR ethanol[tiab]

**#2** - "Mortality"[Mesh] OR "mortality" [Subheading] OR mortality[tiab]

**#3** - (#1 AND #2) NOT ("Animals"[Mesh] NOT ("Animals"[Mesh] AND "Humans"[Mesh])) NOT (editorial[ptyp] OR comment[ptyp] OR news[ptyp] OR letter[ptyp] OR review[ptyp] OR systematic review[ptyp] OR systematic review[ti] OR meta-analysis[ptyp] OR meta-analysis[ti] OR meta-analyses[ti] OR retracted publication[ptyp] OR retraction of publication[ptyp] OR retraction of publication[tiab] OR retraction notice[ti]) Filters: Publication date from 2000/01/01 to 2020/03/11; English

### Cochrane Central Register of Controlled Trials (CENTRAL)

- Provider: John Wiley & Sons
- Date(s) Searched: March 11, 2020
- Date range searched: January 1, 2000-March 11, 2020
- Search Terms:

**#1** - [mh "Alcohol-Induced Disorders"] OR [mh "Alcohol Drinking"] OR [mh "Alcoholic Beverages"] OR [mh "Ethanol"]

**#2** - ("alcohol intake\*" OR "alcohol consum\*" OR "alcohol drink\*" OR "alcoholic beverage\*" OR beer\* OR wine\* OR liquor OR liqueur\* OR spirits OR ethanol):ti,ab,kw

**#3** - #1 OR #2

**#4** - ([mh "Mortality"] OR [mh /MO] OR mortality):ti,ab,kw"

**#5** - #3 AND #4

**#6** - #3 AND #4" with Publication Year from 2000 to 2020, in Trials (Word variations have been searched)

## Embase

- Provider: Elsevier
- Date(s) Searched: March 11, 2020
- Date range searched: January 1, 2000-March 11, 2020
- Search Terms:

**#1** - 'alcoholism'/exp OR 'drinking behavior'/de OR 'alcoholic beverage'/exp OR 'alcohol'/de

**#2** - 'alcohol intake\*':ab,ti OR 'alcohol consum\*':ab,ti OR 'alcohol drink\*':ab,ti OR 'alcoholic beverage\*':ab,ti OR beer\*':ab,ti OR wine\*':ab,ti OR liquor:ab,ti OR liqueur\*':ab,ti OR spirits:ab,ti

**#3** - #1 OR #2

**#4** - 'mortality'/exp OR mortality:ab,ti


**#5** - #3 AND #4

**#6** - #3 AND #4 AND ([article]/lim OR [article in press]/lim) AND [humans]/lim AND [english]/lim AND [2000-2020]/py NOT ([conference abstract]/lim OR [conference paper]/lim OR [conference review]/lim OR [editorial]/lim OR [erratum]/lim OR [letter]/lim OR [note]/lim OR [review]/lim OR [systematic review]/lim OR [meta analysis]/lim)

## LITERATURE SEARCH AND SCREENING RESULTS

The flow chart (**Figure 2**) below illustrates the literature search and screening results for articles examining the systematic review question. The results of the electronic database searches, after removal of duplicates, were screened independently by two NESR analysts using a step-wise process by reviewing titles, abstracts, and full-texts to determine which articles met the inclusion criteria. Refer to **Table 4** for the rationale for exclusion for each excluded full-text article. A manual search was done to find articles that were not identified when searching the electronic databases; all manually identified articles are also screened to determine whether they meet criteria for inclusion.

**Figure 2: Flow chart of literature search and screening results**



## Excluded articles

The table below lists the articles excluded after full-text screening, and includes categories of inclusion and exclusion criteria (see **Table 3**) that studies were excluded. At least one reason for exclusion is provided for each article, though this may not reflect all possible reasons for exclusion. Information about articles excluded after title and abstract screening is available upon request.

**Table 4: Articles excluded after full text screening with rationale for exclusion**

	Citation	Rationale
1	Aburto JM, van Raalte A. Lifespan dispersion in times of life expectancy fluctuation: the case of Central and Eastern Europe. <i>Demography</i> . 2018. 55(6):2071-2096. doi:10.1007/s13524-018-0729-9.	Study Design; Intervention/Exposure
2	Agahi N, Kelfve S, Lennartsson C, Kareholt I. Alcohol consumption in very old age and its association with survival: A matter of health and physical function. <i>Drug Alcohol Depend</i> . 2016. 159:240-5. doi:10.1016/j.drugalcdep.2015.12.022.	Sample size
3	Akbaraly TN, Ferrie JE, Berr C, Brunner EJ, Head J, Marmot MG, Singh-Manoux A, Ritchie K, Shipley, MJ, Kivimaki, M. Alternative Healthy Eating Index and mortality over 18 y of follow-up: results from the Whitehall II cohort. <i>Am J Clin Nutr</i> . 2011. 94(1):247-53. doi:10.3945/ajcn.111.013128.	Comparator
4	AlGhatrif M, Markides KS, Kuo YF, Ray LA, Moore AA. The effect of prevalent cardiovascular conditions on the association between alcohol consumption and mortality among older Mexican American men. <i>Ethn Dis</i> . 2013. 23(2):168-74.	Sample size
5	Arndt V, Rothenbacher D, Krauledat R, Daniel U, Brenner H. Age, alcohol consumption, and all-cause mortality. <i>Ann Epidemiol</i> . 2004. 14(10):750-3. doi:10.1016/j.annepidem.2004.01.004.	Publication Date
6	Baer HJ, Glynn RJ, Hu FB, Hankinson SE, Willett WC, Colditz GA, Stampfer M, Rosner B. Risk factors for mortality in the nurses' health study: a competing risks analysis. <i>Am J Epidemiol</i> . 2011. 173(3):319-29. doi:10.1093/aje/kwq368.	Comparator
7	Baglietto L, English DR, Hopper JL, Powles J, Giles GG. Average volume of alcohol consumed, type of beverage, drinking pattern and the risk of death from all causes. <i>Alcohol Alcohol</i> . 2006. 41(6):664-71. doi:10.1093/alcalc/agl087.	Publication Date
8	Bahat G, Tufan F, Bahat Z, Tufan A, Aydin Y, Akpınar TS, Erten N, Karan MA. Observational cohort study on correlates of mortality in older community-dwelling outpatients: The value of functional assessment. <i>Geriatr Gerontol Int</i> . 2015. 15(11):1219-26. doi:10.1111/ggi.12422.	Intervention/Exposure
9	Bartak M. Alcohol Consumption and Cause-Specific Mortality in Russia: A View From the Czech Republic. <i>J Stud Alcohol Drugs</i> . 2019. 80(5):503-504.	Study Design; Publication Status
10	Batty GD, Lewars H, Emslie C, Gale CR, Hunt K. Internationally recognized guidelines for 'sensible' alcohol consumption: is exceeding them actually detrimental to health and social circumstances? Evidence from a population-based cohort study. <i>J Public Health (Oxf)</i> . 2009. 31(3):360-5. doi:10.1093/pubmed/fdp063.	Publication Date
11	Bazzano LA, Gu D, Reynolds K, Chen J, Wu X, Chen CS, Duan X, Chen J, He J. Alcohol consumption and risk of coronary heart disease among Chinese men. <i>Int J Cardiol</i> . 2009. 135(1):78-85. doi:10.1016/j.ijcard.2008.03.038.	Outcome
12	Behrens G, Leitzmann MF, Sandin S, Lof M, Heid IM, Adami HO, Weiderpass E. The association between alcohol consumption and mortality: the Swedish women's lifestyle and health study. <i>Eur J Epidemiol</i> . 2011. 26(2):81-90. doi:10.1007/s10654-011-9545-x.	Comparator
13	Bellavia A, Tektonidis TG, Orsini N, Wolk A, Larsson SC. Quantifying the benefits of Mediterranean diet in terms of survival. <i>Eur J Epidemiol</i> . 2016. 31(5):527-30. doi:10.1007/s10654-016-0127-9.	Intervention/Exposure
14	Bobak M, Murphy M, Rose R, Marmot M. Determinants of adult mortality in Russia: estimates from sibling data. <i>Epidemiology</i> . 2003. 14(5):603-11. doi:10.1097/01.ede.0000082000.75818.4d.	Publication Date

Citation	Rationale
15 Bondonno NP, Lewis JR, Blekkenhorst LC, Bondonno CP, Shin JH, Croft KD, Woodman RJ, Wong G, Lim WH, Gopinath B, Flood VM, Russell J, Mitchell P, Hodgson JM. Association of flavonoids and flavonoid-rich foods with all-cause mortality: The Blue Mountains Eye Study. <i>Clin Nutr</i> . 2020. 39(1):141-150. doi:10.1016/j.clnu.2019.01.004.	Intervention/Exposure
16 Boyle SH, Mortensen L, Gronbaek M, Barefoot JC. Hostility, drinking pattern and mortality. <i>Addiction</i> . 2008. 103(1):54-9. doi:10.1111/j.1360-0443.2007.02024.x.	Publication Date
17 Breslow RA, Graubard BI. Prospective study of alcohol consumption in the United States: quantity, frequency, and cause-specific mortality. <i>Alcohol Clin Exp Res</i> . 2008. 32(3):513-21. doi:10.1111/j.1530-0277.2007.00595.x.	Outcome
18 Bridevaux IP, Bradley KA, Bryson CL, McDonnell MB, Fihn SD. Alcohol screening results in elderly male veterans: association with health status and mortality. <i>J Am Geriatr Soc</i> . 2004. 52(9):1510-7. doi:10.1111/j.1532-5415.2004.52414.x.	Intervention/Exposure
19 Britton A, Marmot M. Different measures of alcohol consumption and risk of coronary heart disease and all-cause mortality: 11-year follow-up of the Whitehall II Cohort Study. <i>Addiction</i> . 2004. 99(1):109-16. doi:10.1111/j.1360-0443.2004.00530.x.	Publication Date
20 Britton A. Moderate alcohol consumption and total mortality risk: Do not advocate drinking for 'health benefits'. <i>Nutr Metab Cardiovasc Dis</i> . 2019. 29(10):1009-1010. doi:10.1016/j.numecd.2019.06.007.	Study Design; Publication Status
21 Buckland G, Agudo A, Travier N, Huerta, JM, Cirera L, Tormo, MJ, Navarro C, Chirlaque, MD, Moreno-Iribas C, Ardanaz E, Barricarte A, Etxeberria J, Marin P, Quiros, JR, Redondo, ML, Larranaga N, Amiano P, Dorronsoro M, Arriola L, Basterretxea M, Sanchez, MJ, Molina E, Gonzalez CA. Adherence to the Mediterranean diet reduces mortality in the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Spain). <i>Br J Nutr</i> . 2011. 106(10):1581-91. doi:10.1017/s0007114511002078.	Intervention/Exposure; Comparator
22 Burke V, Zhao Y, Lee AH, Hunter E, Spargo RM, Gracey M, Smith RM, Beilin LJ, Puddey IB. Health-related behaviours as predictors of mortality and morbidity in Australian Aborigines. <i>Prev Med</i> . 2007. 44(2):135-42. doi:10.1016/j.ypmed.2006.09.008.	Publication Date
23 Byun W, Sieverdes JC, Sui X, Hooker SP, Lee CD, Church TS, Blair SN. Effect of positive health factors and all-cause mortality in men. <i>Med Sci Sports Exerc</i> . 2010. 42(9):1632-8. doi:10.1249/MSS.0b013e3181d43f29.	Comparator
24 Carlsson AC, Theobald H, Wandell PE. Health factors and longevity in men and women: a 26-year follow-up study. <i>Eur J Epidemiol</i> . 2010. 25(8):547-51. doi:10.1007/s10654-010-9472-2.	Sample size
25 Carlsson AC, Wandell PE, Gigante B, Leander K, Hellenius ML, de Faire U. Seven modifiable lifestyle factors predict reduced risk for ischemic cardiovascular disease and all-cause mortality regardless of body mass index: a cohort study. <i>Int J Cardiol</i> . 2013. 168(2):946-52. doi:10.1016/j.ijcard.2012.10.045.	Intervention/Exposure
26 Carpenter C, Dobkin C. The Effect of Alcohol Consumption on Mortality: Regression Discontinuity Evidence from the Minimum Drinking Age. <i>Am Econ J Appl Econ</i> . 2009. 1(1):164-182. doi:10.1257/app.1.1.164.	Intervention/Exposure
27 Chen LY, Hardy CL. Alcohol consumption and health status in older adults: a longitudinal analysis. <i>J Aging Health</i> . 2009. 21(6):824-47. doi:10.1177/0898264309340688.	Publication Date
28 Cheung MR. Lack of health insurance increases all cause and all cancer mortality in adults: an analysis of National Health and Nutrition Examination Survey (NHANES III) data. <i>Asian Pac J Cancer Prev</i> . 2013. 14(4):2259-63. doi:10.7314/apjcp.2013.14.4.2259.	Comparator
29 Chikritzh T, Stockwell T, Jonas H, Stevenson C, Cooper-Stanbury M, Donath S, Single E, Catalano P. Towards a standardised methodology for estimating alcohol-caused death, injury and illness in Australia. <i>Aust N Z J Public Health</i> . 2002. 26(5):443-50. doi:10.1111/j.1467-842x.2002.tb00345.x.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
30 Christensen AI, Ekholm O, Gray L, Glumer C, Juel K. What is wrong with non-respondents? Alcohol-, drug- and smoking-related mortality and morbidity in a 12-year follow-up study of respondents and non-respondents in the Danish Health and Morbidity Survey. <i>Addiction</i> . 2015. 110(9):1505-12. doi:10.1111/add.12939.	Intervention/Exposure
31 Cloud AJ, Thai A, Liao Y, Terry MB. The impact of cancer prevention guideline adherence on overall mortality in a high-risk cohort of women from the New York site of the Breast Cancer Family Registry. <i>Breast Cancer Res Treat</i> . 2015. 149(2):537-46. doi:10.1007/s10549-014-3234-x.	Intervention/Exposure; Outcome; Health Status

	Citation	Rationale
32	Colpani V, Oppermann K, Spritzer PM. Causes of death and associated risk factors among climacteric women from Southern Brazil: a population based-study. <i>BMC Public Health</i> . 2014. 14:194. doi:10.1186/1471-2458-14-194.	Country
33	Connor J, Kydd R, Shield K, Rehm J. The burden of disease and injury attributable to alcohol in New Zealanders under 80 years of age: marked disparities by ethnicity and sex. <i>N Z Med J</i> . 2015. 128(1409):15-28.	Study Design; Intervention/Exposure; Outcome
34	Dai J, Mukamal KJ, Krasnow RE, Swan GE, Reed T. Higher usual alcohol consumption was associated with a lower 41-y mortality risk from coronary artery disease in men independent of genetic and common environmental factors: the prospective NHLBI Twin Study. <i>Am J Clin Nutr</i> . 2015. 102(1):31-9. doi:10.3945/ajcn.114.106435.	Sample size
35	Daly M, Hall PA, Allan JL. Time Perspective and All-Cause Mortality: Evidence From the English Longitudinal Study of Ageing. <i>Ann Behav Med</i> . 2019. 53(5):486-492. doi:10.1093/abm/kay046.	Intervention/Exposure
36	Davis JS, Banfield E, Lee HY, Peng HL, Chang S, Wood AC. Lifestyle behavior patterns and mortality among adults in the NHANES 1988-1994 population: A latent profile analysis. <i>Prev Med</i> . 2019. 120:131-139. doi:10.1016/j.ypmed.2019.01.012.	Intervention/Exposure
37	Davletov K, McKee M, Berkinbayev S, Battakova Z, Vujnovic M, Rechel B. Regional differences in cardiovascular mortality in Kazakhstan: further evidence for the 'Russian mortality paradox'?. <i>Eur J Public Health</i> . 2015. 25(5):890-4. doi:10.1093/eurpub/ckv019.	Outcome
38	Dawson DA. Alcohol consumption, alcohol dependence, and all-cause mortality. <i>Alcohol Clin Exp Res</i> . 2000. 24(1):72-81.	Publication Date
39	De la Torre R, Corella D, Castaner O, Martinez-Gonzalez MA, Salas-Salvado J, Vila J, Estruch R, Sorli JV, Aros F, Fiol M, Ros E, Serra-Majem L, Pinto X, Gomez-Gracia E, Lapetra J, Ruiz-Canela M, Basora J, Asensio EM, Covas MI, Fito M. Protective effect of homovanillyl alcohol on cardiovascular disease and total mortality: virgin olive oil, wine, and catechol-methylation. <i>Am J Clin Nutr</i> . 2017. 105(6):1297-1304. doi:10.3945/ajcn.116.145813.	Intervention/Exposure
40	de Vegt F, Dekker JM, Groeneveld WJ, Nijpels G, Stehouwer CD, Bouter LM, Heine RJ. Moderate alcohol consumption is associated with lower risk for incident diabetes and mortality: the Hoorn Study. <i>Diabetes Res Clin Pract</i> . 2002. 57(1):53-60. doi:10.1016/s0168-8227(02)00013-x.	Publication Date
41	Ding D, Rogers K, van der Ploeg H, Stamatakis E, Bauman AE. Traditional and Emerging Lifestyle Risk Behaviors and All-Cause Mortality in Middle-Aged and Older Adults: Evidence from a Large Population-Based Australian Cohort. <i>PLoS Med</i> . 2015. 12(12):e1001917. doi:10.1371/journal.pmed.1001917.	Comparator
42	Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to alcohol consumption: a prospective study among male British doctors. <i>Int J Epidemiol</i> . 2005. 34(1):199-204. doi:10.1093/ije/dyh369.	Publication Date
43	Doniec K, Stefler D, Murphy M, Gugushvili A, McKee M, Marmot M, Bobak M, King, L. Education and mortality in three Eastern European populations: findings from the PrivMort retrospective cohort study. <i>Eur J Public Health</i> . 2019. 29(3):549-554. doi:10.1093/eurpub/cky254.	Intervention/Exposure; Outcome
44	Ebbert JO, Janney CA, Sellers TA, Folsom AR, Cerhan JR. The association of alcohol consumption with coronary heart disease mortality and cancer incidence varies by smoking history. <i>J Gen Intern Med</i> . 2005. 20(1):14-20. doi:10.1111/j.1525-1497.2005.40129.x.	Publication Date
45	Egeberg A, Fowler JF, Jr, Gislason GH, Thyssen JP. Nationwide Assessment of Cause-Specific Mortality in Patients with Rosacea: A Cohort Study in Denmark. <i>Am J Clin Dermatol</i> . 2016. 17(6):673-679. doi:10.1007/s40257-016-0217-1.	Health Status
46	Eguchi E, Iso H, Tanabe N, Wada Y, Yatsuya H, Kikuchi S, Inaba Y, Tamakoshi A. Healthy lifestyle behaviours and cardiovascular mortality among Japanese men and women: the Japan collaborative cohort study. <i>Eur Heart J</i> . 2012. 33(4):467-77. doi:10.1093/eurheartj/ehr429.	Outcome
47	Eliassen M, Becker U, Gronbaek M, Juel K, Tolstrup JS. Alcohol-attributable and alcohol-preventable mortality in Denmark: an analysis of which intake levels contribute most to alcohol's harmful and beneficial effects. <i>Eur J Epidemiol</i> . 2014. 29(1):15-26. doi:10.1007/s10654-013-9855-2.	Outcome

	Citation	Rationale
48	Elovainio M, Hakulinen C, Pulkki-Raback L, Virtanen M, Josefsson K, Jokela M, Vahtera J, Kivimäki M. Contribution of risk factors to excess mortality in isolated and lonely individuals: an analysis of data from the UK Biobank cohort study. <i>Lancet Public Health</i> . 2017. 2(6):e260-e266. doi:10.1016/s2468-2667(17)30075-0.	Intervention/Exposure
49	Embersson JR, Shaper AG, Wannamethee SG, Morris RW, Whincup PH. Alcohol intake in middle age and risk of cardiovascular disease and mortality: accounting for intake variation over time. <i>Am J Epidemiol</i> . 2005. 161(9):856-63. doi:10.1093/aje/kwi111.	Publication Date
50	Eyawo O, McGinnis KA, Justice AC, Fiellin DA, Hahn JA, Williams EC, Gordon AJ, Marshall BDL, Kraemer KL, Crystal S, Gaither JR, Edelman EJ, Bryant KJ, Tate JP. Alcohol and Mortality: Combining Self-Reported (AUDIT-C) and Biomarker Detected (PEth) Alcohol Measures Among HIV Infected and Uninfected. <i>Journal of acquired immune deficiency syndromes</i> (1999). 2018. 77(2):135-143. doi:10.1097/QAI.0000000000001588.	Sample size
51	Farchi G, Fidanza F, Giampaoli S, Mariotti S, Menotti A. Alcohol and survival in the Italian rural cohorts of the Seven Countries Study. <i>Int J Epidemiol</i> . 2000. 29(4):667-71. doi:10.1093/ije/29.4.667.	Publication Date
52	Fichter MM, Quadflieg N, Fischer UC. Severity of alcohol-related problems and mortality: results from a 20-year prospective epidemiological community study. <i>Eur Arch Psychiatry Clin Neurosci</i> . 2011. 261(4):293-302. doi:10.1007/s00406-010-0141-y.	Intervention/Exposure
53	Fillmore KM, Kerr WC, Bostrom A. Changes in drinking status, serious illness and mortality. <i>J Stud Alcohol</i> . 2003. 64(2):278-85. doi:10.15288/jsa.2003.64.278.	Publication Date
54	Ford ES, Zhao G, Tsai J, Li C. Low-risk lifestyle behaviors and all-cause mortality: findings from the National Health and Nutrition Examination Survey III Mortality Study. <i>Am J Public Health</i> . 2011. 101(10):1922-9. doi:10.2105/ajph.2011.300167	Comparator
55	Freiberg MS, Chang YF, Kraemer KL, Robinson JG, Adams-Campbell LL, Kuller LL. Alcohol consumption, hypertension, and total mortality among women. <i>Am J Hypertens</i> . 2009. 22(11):1212-8. doi:10.1038/ajh.2009.172.	Publication Date
56	Friesema IH, Zwietering PJ, Veenstra MY, Knottnerus JA, Garretsen HF, Kester AD, Lemmens PH. The effect of alcohol intake on cardiovascular disease and mortality disappeared after taking lifetime drinking and covariates into account. <i>Alcohol Clin Exp Res</i> . 2008. 32(4):645-51. doi:10.1111/j.1530-0277.2007.00612.x.	Publication Date
57	Friesema IH, Zwietering PJ, Veenstra MY, Knottnerus JA, Garretsen HF, Lemmens PH. Alcohol intake and cardiovascular disease and mortality: the role of pre-existing disease. <i>J Epidemiol Community Health</i> . 2007. 61(5):441-6. doi:10.1136/jech.2006.050419.	Publication Date
58	Furuya H, Young AF, Powers JR, Byles JE. Alcohol consumption and physical health-related quality of life in older women using the transformation of SF-36 to account for death. <i>Nihon Arukoru Yakubutsu Igakkai Zasshi</i> . 2008. 43(2):97-109.	Outcome
59	Gargiulo G, Testa G, Cacciatore F, Mazzella F, Galizia G, Della-Morte D, Langellotto A, Pirozzi G, Ferro G, Ferrara N, Rengo F, Abete P. Moderate alcohol consumption predicts long-term mortality in elderly subjects with chronic heart failure. <i>J Nutr Health Aging</i> . 2013. 17(5):480-5. doi:10.1007/s12603-012-0430-4.	Comparator; Health Status
60	Gaziano JM, Gaziano TA, Glynn RJ, Sesso HD, Ajani UA, Stampfer MJ, Manson JE, Hennekens CH, Buring JE. Light-to-moderate alcohol consumption and mortality in the Physicians' Health Study enrollment cohort. <i>J Am Coll Cardiol</i> . 2000. 35(1):96-105. doi:10.1016/s0735-1097(99)00531-8.	Publication Date
61	Glover J, Gold, JA. Alcohol Use Increasing Among Adults 65 and Older. <i>Wmj</i> . 2017. 116(3):182-183.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
62	Golden, RN, Maroney, L. Addressing a deadly epidemic. <i>Wmj</i> . 2009. 108(7):377-8.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
63	Gray, CM, Carroll, RJ, Lentjes, MAH, Keogh, RH. Correcting for measurement error in fractional polynomial models using Bayesian modelling and regression calibration, with an application to alcohol and mortality. <i>Biom J</i> . 2019. 61(3):558-573. doi:10.1002/bimj.201700279.	Outcome

	Citation	Rationale
64	Greenfield, TK, Rehm J, Rogers, JD. Effects of depression and social integration on the relationship between alcohol consumption and all-cause mortality. <i>Addiction</i> . 2002. 97(1):29-38. doi:10.1046/j.1360-0443.2002.00065.x.	Publication Date
65	Greenlee H, Strizich G, Lovasi, GS, Kaplan, RC, Biggs, ML, Li, CI, Richardson J, Burke, GL, Fitzpatrick, AL, Fretts, AM, Psaty, BM, Fried, LP. Concordance With Prevention Guidelines and Subsequent Cancer, Cardiovascular Disease, and Mortality: A Longitudinal Study of Older Adults. <i>Am J Epidemiol</i> . 2017. 186(10):1168-1179. doi:10.1093/aje/kwx150.	Intervention/Exposure
66	Gronbaek M, Becker U, Johansen D, Gottschau A, Schnohr P, Hein, HO, Jensen G, Sorensen, TI. Type of alcohol consumed and mortality from all causes, coronary heart disease, and cancer. <i>Ann Intern Med</i> . 2000. 133(6):411-9. doi:10.7326/0003-4819-133-6-200009190-00008.	Publication Date
67	Gronbaek M, Johansen D, Becker U, Hein, HO, Schnohr P, Jensen G, Vestbo J, Sorensen, TI. Changes in alcohol intake and mortality: a longitudinal population-based study. <i>Epidemiology</i> . 2004. 15(2):222-8. doi:10.1097/01.ede.0000112219.01955.56.	Publication Date
68	Gun, RT, Pratt N, Ryan P, Gordon I, Roder, D. Tobacco and alcohol-related mortality in men: estimates from the Australian cohort of petroleum industry workers. <i>Aust N Z J Public Health</i> . 2006. 30(4):318-24. doi:10.1111/j.1467-842x.2006.tb00842.x.	Publication Date
69	Hallgren, MA, Hogberg P, Andreasson, S. Alcohol consumption and harm among elderly Europeans: falling between the cracks. <i>Eur J Public Health</i> . 2010. 20(6):616-7. doi:10.1093/eurpub/ckq111.	Study Design
70	Halme, JT, Seppa K, Alho H, Poikolainen K, Pirkola S, Aalto, M. Alcohol consumption and all-cause mortality among elderly in Finland. <i>Drug Alcohol Depend</i> . 2010. 106(2-3):212-8. doi:10.1016/j.drugalcdep.2009.08.017.	Comparator
71	Hamer M, O'Donovan G, Stamatakis, E. Lifestyle risk factors, obesity and infectious disease mortality in the general population: Linkage study of 97,844 adults from England and Scotland. <i>Prev Med</i> . 2019. 123:65-70. doi:10.1016/j.ypmed.2019.03.002.	Outcome
72	Harriss, LR, English, DR, Hopper, JL, Powles J, Simpson, JA, O'Dea K, Giles, GG, Tonkin, AM. Alcohol consumption and cardiovascular mortality accounting for possible misclassification of intake: 11-year follow-up of the Melbourne Collaborative Cohort Study. <i>Addiction</i> . 2007. 102(10):1574-85. doi:10.1111/j.1360-0443.2007.01947.x.	Outcome
73	Hart, CL, Davey Smith G, Gruer L, Watt, GC. The combined effect of smoking tobacco and drinking alcohol on cause-specific mortality: a 30 year cohort study. <i>BMC Public Health</i> . 2010. 10:789. doi:10.1186/1471-2458-10-789.	Comparator
74	Hart CL, Smith GD. Alcohol consumption and mortality and hospital admissions in men from the Midspan collaborative cohort study. <i>Addiction</i> . 2008. 103(12):1979-86. doi:10.1111/j.1360-0443.2008.02373.x.	Publication Date
75	Haupt TH, Rasmussen LJH, Kallermose T, Ladelund S, Andersen O, Pisinger C, Eugen-Olsen J. Healthy lifestyles reduce suPAR and mortality in a Danish general population study. <i>Immun Ageing</i> . 2019. 16:1. doi:10.1186/s12979-018-0141-8.	Intervention/Exposure; Outcome; Health Status
76	Hayward RD, Krause, N. The effect of belonging to an alcohol-proscribing religious group on the relationship between moderate alcohol consumption and mortality. <i>Soc Sci Med</i> . 2014. 101:1-8. doi:10.1016/j.socscimed.2013.10.039.	Comparator
77	Heitz AE, Baumgartner RN, Baumgartner KB, Boone SD. Healthy lifestyle impact on breast cancer-specific and all-cause mortality. <i>Breast Cancer Res Treat</i> . 2018. 167(1):171-181. doi:10.1007/s10549-017-4467-2.	Health Status
78	Henry JA, Moloney C, Rivas C, Goldin RD. Increase in alcohol related deaths: is hepatitis C a factor?. <i>J Clin Pathol</i> . 2002. 55(9):704-7. doi:10.1136/jcp.55.9.704.	Intervention/Exposure; Outcome
79	Hernandez-Hernandez A, Gea A, Ruiz-Canela M, Toledo E, Beunza JJ, Bes-Rastrollo M, Martinez-Gonzalez MA. Mediterranean Alcohol-Drinking Pattern and the Incidence of Cardiovascular Disease and Cardiovascular Mortality: The SUN Project. <i>Nutrients</i> . 2015. 7(11):9116-26. doi:10.3390/nu7115456.	Outcome
80	Hernández-Vásquez A, Vargas-Fernández R, Rebatta-Acuña A, Bendezu-Quispe, G. Trends in mortality due to gastrointestinal diseases attributed to alcohol use in Peru from 2003 to 2016. <i>Revista de gastroenterología del Perú : organo oficial de la Sociedad de Gastroenterología del Perú</i> . 2019. 39(3):239-245.	Intervention/Exposure; Outcome
81	Heroux M, Janssen I, Lam M, Lee, DC, Hebert, JR, Sui X, Blair, SN. Dietary patterns and the risk of mortality: impact of cardiorespiratory fitness. <i>Int J Epidemiol</i> . 2010. 39(1):197-209. doi:10.1093/ije/dyp191.	Intervention/Exposure
82	Hoiseth G, Magnus P, Knudsen GP, Jansen MD, Naess O, Tambs K, Morland J. Is ADH1C genotype relevant for the cardioprotective effect of alcohol? <i>Alcohol</i> . 2013. 47(2):81-4. doi:10.1016/j.alcohol.2012.12.005.	Outcome



	Citation	Rationale
83	Holahan CJ, Brennan PL, Schutte KK, Holahan CK, Hixon JG, Moos RH. Drinking Level Versus Drinking Pattern and Cigarette Smoking Among Older Adults. <i>Alcohol Clin Exp Res</i> . 2018. 42(4):795-802. doi:10.1111/acer.13607.	Sample size
84	Holahan CJ, Schutte KK, Brennan PL, Holahan CK, Moos RH. Episodic heavy drinking and 20-year total mortality among late-life moderate drinkers. <i>Alcohol Clin Exp Res</i> . 2014. 38(5):1432-8. doi:10.1111/acer.12381.	Sample size
85	Holahan CJ, Schutte KK, Brennan PL, North RJ, Holahan CK, Moos BS, Moos RH. Wine consumption and 20-year mortality among late-life moderate drinkers. <i>J Stud Alcohol Drugs</i> . 2012. 73(1):80-8. doi:10.15288/jsad.2012.73.80.	Sample size
86	Howie EK, Sui X, Lee DC, Hooker SP, Hebert JR, Blair SN. Alcohol consumption and risk of all-cause and cardiovascular disease mortality in men. <i>J Aging Res</i> . 2011. 2011:805062. doi:10.4061/2011/805062.	Comparator
87	Hulsegge G, Looman M, Smit HA, Daviglus ML, van der Schouw YT, Verschuren WM. Lifestyle Changes in Young Adulthood and Middle Age and Risk of Cardiovascular Disease and All-Cause Mortality: The Doetinchem Cohort Study. <i>J Am Heart Assoc</i> . 2016. 5(1). doi:10.1161/jaha.115.002432.	Comparator
88	Huynh QL, Reid CM, Chowdhury EK, Huq MM, Billah B, Wing LM, Tonkin AM, Simons LA, Nelson MR. Prediction of cardiovascular and all-cause mortality at 10 years in the hypertensive aged population. <i>Am J Hypertens</i> . 2015. 28(5):649-56. doi:10.1093/ajh/hpu213.	Intervention/Exposure; Health Status
89	Ikehara S, Iso H, Toyoshima H, Date C, Yamamoto A, Kikuchi S, Kondo T, Watanabe Y, Koizumi A, Wada Y, Inaba Y, Tamakoshi A. Alcohol consumption and mortality from stroke and coronary heart disease among Japanese men and women: the Japan collaborative cohort study. <i>Stroke</i> . 2008. 39(11):2936-42. doi:10.1161/strokeaha.108.520288.	Outcome
90	Im PK, Millwood IY, Chen Y, Guo Y, Du H, Kartsonaki C, Bian Z, Tan Y, Su J, Li Y, Yu C, Lv J, Li L, Yang L, Chen Z. Problem drinking, wellbeing and mortality risk in Chinese men: findings from the China Kadoorie Biobank. <i>Addiction</i> . 2019. doi:10.1111/add.14873.	Country
91	Ivey KL, Hodgson JM, Croft KD, Lewis JR, Prince RL. Flavonoid intake and all-cause mortality. <i>Am J Clin Nutr</i> . 2015. 101(5):1012-20. doi:10.3945/ajcn.113.073106.	Intervention/Exposure
92	Ivey KL, Jensen MK, Hodgson JM, Eliassen AH, Cassidy A, Rimm EB. Association of flavonoid-rich foods and flavonoids with risk of all-cause mortality. <i>Br J Nutr</i> . 2017. 117(10):1470-1477. doi:10.1017/s0007114517001325.	Intervention/Exposure
93	Jakovljevic B, Stojanov V, Paunovic K, Belojevic G, Milic N. Alcohol consumption and mortality in Serbia: twenty-year follow-up study. <i>Croat Med J</i> . 2004. 45(6):764-8.	Publication Date
94	Jankovic N, Geelen A, Streppel MT, de Groot LC, Kieft-de Jong JC, Orfanos P, Bamia C, Trichopoulou A, Boffetta P, Bobak M, Pikhart H, Kee F, O'Doherty MG, Buckland G, Woodside J, Franco OH, Ikram MA, Struijk EA, Pajak A, Malyutina S, Kubinova R, Wennberg M, Park Y, Bueno-de-Mesquita HB, Kampman E, Feskens EJ. WHO guidelines for a healthy diet and mortality from cardiovascular disease in European and American elderly: the CHANCES project. <i>Am J Clin Nutr</i> . 2015. 102(4):745-56. doi:10.3945/ajcn.114.095117.	Outcome
95	Jayanama K, Theou O, Blodgett JM, Cahill L, Rockwood K. Frailty, nutrition-related parameters, and mortality across the adult age spectrum. <i>BMC Med</i> . 2018. 16(1):188. doi:10.1186/s12916-018-1176-6.	Intervention/Exposure; Outcome
96	Jeong HG, Kim TH, Lee JJ, Lee SB, Park JH, Huh Y, Chin HJ, Jhoo JH, Lee DY, Woo JI, Kim KW. Impact of alcohol use on mortality in the elderly: results from the Korean Longitudinal Study on Health and Aging. <i>Drug Alcohol Depend</i> . 2012. 121(1-2):133-9. doi:10.1016/j.drugalcdep.2011.08.017.	Sample size
97	Jiang H, Livingston M, Room R, Gan Y, English D, Chenhall R. Can public health policies on alcohol and tobacco reduce a cancer epidemic? Australia's experience. <i>BMC Med</i> . 2019. 17(1):213. doi:10.1186/s12916-019-1453-z.	Intervention/Exposure; Outcome
98	Johansen D, Gronbaek M, Overvad K, Schnohr P, Andersen PK. Generalized additive models applied to analysis of the relation between amount and type of alcohol and all-cause mortality. <i>Eur J Epidemiol</i> . 2005. 20(1):29-36. doi:10.1007/s10654-004-2172-z.	Comparator
99	Johnson JE, Finney JW, Moos RH. Predictors of 5-year mortality following inpatient/residential group treatment for substance use disorders. <i>Addict Behav</i> . 2005. 30(7):1300-16. doi:10.1016/j.addbeh.2005.01.005.	Intervention/Exposure; Health Status

	Citation	Rationale
100	Kamsa-ard S, Promthet S, Lewington S, Burrett JA, Sherliker P, Kamsa-ard S, Suwanrungruang K, Parkin DM. Alcohol consumption and mortality: the Khon Kaen Cohort Study, Thailand. <i>J Epidemiol</i> . 2014. 24(2):154-60. doi:10.2188/jea.je20130092.	Country
101	Katcher BS, Reiter RB, Aragon TJ. Estimating alcohol-related premature mortality in San Francisco: use of population-attributable fractions from the global burden of disease study. <i>BMC Public Health</i> . 2010. 10:682. doi:10.1186/1471-2458-10-682.	Comparator; Outcome
102	Kerr WC, Fillmore KM, Marvy P. Beverage-specific alcohol consumption and cirrhosis mortality in a group of English-speaking beer-drinking countries. <i>Addiction</i> . 2000. 95(3):339-46. doi:10.1046/j.1360-0443.2000.9533394.x.	Outcome
103	Keys A, Menott A, Karvonen, MJ, Aravanjs C, Blackburn H, Buzina R, Djordjevic, BS, Dontas, AS, Fidanza F, Keys, MH, Kromhout D, Nedeljkovic S, Punsar S, Seccareccia F, Toshima, H. The diet and 15-year death rate in the seven countries study. <i>American Journal of Epidemiology</i> . 2017. 185(11):1130-1142. doi:10.1093/aje/kwx101.	Publication Date
104	Khang, YH, Lynch, JW, Yang S, Harper S, Yun, SC, Jung-Choi K, Kim, HR. The contribution of material, psychosocial, and behavioral factors in explaining educational and occupational mortality inequalities in a nationally representative sample of South Koreans: relative and absolute perspectives. <i>Soc Sci Med</i> . 2009. 68(5):858-66. doi:10.1016/j.socscimed.2008.12.003.	Publication Date
105	Khaw, KT, Wareham N, Bingham S, Welch A, Luben R, Day, N. Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. <i>PLoS Med</i> . 2008. 5(1):e12. doi:10.1371/journal.pmed.0050012.	Publication Date
106	Kiiskinen T, Mars, NJ, Palviainen T, Koskela J, Ramo, JT, Ripatti P, Ruotsalainen S, Palotie A, Madden, PAF, Rose, RJ, Kaprio J, Salomaa V, Makela P, Havulinna, AS, Ripatti, S. Genomic prediction of alcohol-related morbidity and mortality. <i>Transl Psychiatry</i> . 2020. 10(1):23. doi:10.1038/s41398-019-0676-2.	Intervention/exposure
107	Kim, MK, Ko, MJ, Han, JT. Alcohol consumption and mortality from all-cause and cancers among 1.34 million Koreans: the results from the Korea national health insurance corporation's health examinee cohort in 2000. <i>Cancer Causes Control</i> . 2010. 21(12):2295-302. doi:10.1007/s10552-010-9656-9.	Comparator
108	Kinder, LS, Bryson, CL, Sun H, Williams, EC, Bradley, KA. Alcohol screening scores and all-cause mortality in male Veterans Affairs patients. <i>J Stud Alcohol Drugs</i> . 2009. 70(2):253-60. doi:10.15288/jsad.2009.70.253.	Comparator
109	King, DE, Mainous, AG, 3rd, Geesey, ME. Adopting moderate alcohol consumption in middle age: subsequent cardiovascular events. <i>Am J Med</i> . 2008. 121(3):201-6. doi:10.1016/j.amjmed.2007.12.004.	Outcome
110	King, DE, Mainous, AG, 3rd, Matheson, EM, Everett, CJ. Impact of healthy lifestyle on mortality in people with normal blood pressure, LDL cholesterol, and C-reactive protein. <i>Eur J Prev Cardiol</i> . 2013. 20(1):73-9. doi:10.1177/1741826711425776.	Intervention/Exposure
111	Klatsky, AL, Friedman, GD, Armstrong, MA, Kipp, H. Wine, liquor, beer, and mortality. <i>Am J Epidemiol</i> . 2003. 158(6):585-95. doi:10.1093/aje/kwg184.	Publication Date
112	Klatsky, AL, Udaltsova, N. Alcohol Drinking and Total Mortality Risk. <i>Annals of Epidemiology</i> . 2007. 17(5 SUPPL.):S63-S67. doi:10.1016/j.annepidem.2007.01.014.	Publication Date
113	Knoops, KT, de Groot, LC, Kromhout D, Perrin, AE, Moreiras-Varela O, Menotti A, van Staveren, WA. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. <i>Jama</i> . 2004. 292(12):1433-9. doi:10.1001/jama.292.12.1433.	Publication Date
114	Koch, MB, Diderichsen F, Gronbaek M, Juel, K. What is the association of smoking and alcohol use with the increase in social inequality in mortality in Denmark? A nationwide register-based study. <i>BMJ Open</i> . 2015. 5(5):e006588. doi:10.1136/bmjopen-2014-006588.	Intervention/Exposure; Outcome
115	Kontro, TK, Sarna S, Kaprio J, Kujala, UM. Mortality and health-related habits in 900 Finnish former elite athletes and their brothers. <i>Br J Sports Med</i> . 2018. 52(2):89-95. doi:10.1136/bjsports-2017-098206.	Intervention/Exposure; Comparator
116	Korotayev A, Khaltourina D, Meshcherina K, Zamiatnina, E. Distilled Spirits Overconsumption as the Most Important Factor of Excessive Adult Male Mortality in Europe. <i>Alcohol Alcohol</i> . 2018. 53(6):742-752. doi:10.1093/alcalc/agy054.	Intervention/Exposure
117	Kovše K, Tomšič S, Ponikvar, BM, Nadrag, P. Alcohol related harm in Slovenia. <i>Zdravniški Vestnik</i> . 2012. 81(2):119-127.	Language
118	Kristenson H, Osterling A, Nilsson, JA, Lindgarde, F. Prevention of alcohol-related deaths in middle-aged heavy drinkers. <i>Alcohol Clin Exp Res</i> . 2002. 26(4):478-84. doi:10.1111/j.1530-0277.2002.tb02564.x.	Intervention/Exposure; Outcome

Citation	Rationale
119 Krokstad S, Ding D, Grunseit, AC, Sund, ER, Holmen, TL, Rangul V, Bauman, A. Multiple lifestyle behaviours and mortality, findings from a large population-based Norwegian cohort study - The HUNT Study. <i>BMC Public Health</i> . 2017. 17(1):58. doi:10.1186/s12889-016-3993-x.	Intervention/Exposure
120 Krueger, PM, Saint Onge, JM, Chang, VW. Race/ethnic differences in adult mortality: the role of perceived stress and health behaviors. <i>Soc Sci Med</i> . 2011. 73(9):1312-22. doi:10.1016/j.socscimed.2011.08.007.	Comparator
121 Kvaavik E, Batty, GD, Ursin G, Huxley R, Gale, CR. Influence of individual and combined health behaviors on total and cause-specific mortality in men and women: the United Kingdom health and lifestyle survey. <i>Arch Intern Med</i> . 2010. 170(8):711-8. doi:10.1001/archinternmed.2010.76.	Comparator
122 Laaksonen M, Talala K, Martelin T, Rahkonen O, Roos E, Helakorpi S, Laatikainen T, Prattala, R. Health behaviours as explanations for educational level differences in cardiovascular and all-cause mortality: a follow-up of 60 000 men and women over 23 years. <i>Eur J Public Health</i> . 2008. 18(1):38-43. doi:10.1093/eurpub/ckm051.	Publication Date
123 Laatikainen T, Manninen L, Poikolainen K, Vartiainen, E. Increased mortality related to heavy alcohol intake pattern. <i>J Epidemiol Community Health</i> . 2003. 57(5):379-84. doi:10.1136/jech.57.5.379.	Publication Date
124 Laine, JE, Baltar, VT, Stringini S, Gandini M, Chadeau-Hyam M, Kivimaki M, Severi G, Perduca V, Hodge, AM, Dugue, PA, Giles, GG, Milne, RL, Barros H, Sacerdote C, Krogh V, Panico S, Tumino R, Goldberg M, Zins M, Delpierre C, Vineis, P. Reducing socio-economic inequalities in all-cause mortality: a counterfactual mediation approach. <i>Int J Epidemiol</i> . 2019. doi:10.1093/ije/dyz248.	Intervention/Exposure
125 Lang I, Guralnik J, Wallace, RB, Melzer, D. What level of alcohol consumption is hazardous for older people? Functioning and mortality in U.S. and English national cohorts. <i>J Am Geriatr Soc</i> . 2007. 55(1):49-57. doi:10.1111/j.1532-5415.2006.01007.x.	Publication Date
126 Lapidus L, Bengtsson C, Bergfors E, Bjorkelund C, Spak F, Lissner, L. Alcohol intake among women and its relationship to diabetes incidence and all-cause mortality: the 32-year follow-up of a population study of women in Gothenburg, Sweden. <i>Diabetes Care</i> . 2005. 28(9):2230-5. doi:10.2337/diacare.28.9.2230.	Publication Date
127 Laranjeira E, Szrek, H. Going beyond life expectancy in assessments of health systems' performance: life expectancy adjusted by perceived health status. <i>Int J Health Econ Manag</i> . 2016. 16(2):133-161. doi:10.1007/s10754-015-9183-z.	Study Design
128 Larsson, SC, Kaluza J, Wolk, A. Combined impact of healthy lifestyle factors on lifespan: two prospective cohorts. <i>J Intern Med</i> . 2017. 282(3):209-219. doi:10.1111/joim.12637.	Intervention/Exposure; Comparator
129 Lee, DH, Folsom, AR, Jacobs, DR, Jr. Iron, zinc, and alcohol consumption and mortality from cardiovascular diseases: the Iowa Women's Health Study. <i>Am J Clin Nutr</i> . 2005. 81(4):787-91. doi:10.1093/ajcn/81.4.787.	Outcome
130 Lee I, Kim S, Kang, H. Lifestyle Risk Factors and All-Cause and Cardiovascular Disease Mortality: Data from the Korean Longitudinal Study of Aging. <i>Int J Environ Res Public Health</i> . 2019. 16(17). doi:10.3390/ijerph16173040.	Comparator
131 Lee, JH, Lee, DH. Comparison of injury pattern and clinical outcomes between young adults and elderly patients with alcohol-related injury in South Korea 2011-2016. <i>PeerJ</i> . 2019. 7:e7704. doi:10.7717/peerj.7704.	Intervention/Exposure; Health Status
132 Lee, SJ, Sudore, RL, Williams, BA, Lindquist K, Chen, HL, Covinsky, KE. Functional limitations, socioeconomic status, and all-cause mortality in moderate alcohol drinkers. <i>J Am Geriatr Soc</i> . 2009. 57(6):955-62. doi:10.1111/j.1532-5415.2009.02184.x.	Publication Date
133 Levantesi G, Marfisi R, Mozaffarian D, Franzosi, MG, Maggioni A, Nicolosi, GL, Schweiger C, Silletta M, Tavazzi L, Tognoni G, Marchioli R. Wine consumption and risk of cardiovascular events after myocardial infarction: results from the GISSI-Prevenzione trial. <i>Int J Cardiol</i> . 2013. 163(3):282-287. doi:10.1016/j.ijcard.2011.06.053.	Health Status
134 Lhachimi SK, Nusselder WJ, Smit HA, Baili P, Bennett K, Fernandez E, Kulik MC, Lobstein T, Pomerleau J, Boshuizen HC, Mackenbach JP. Potential health gains and health losses in eleven EU countries attainable through feasible prevalences of the life-style related risk factors alcohol, BMI, and smoking: a quantitative health impact assessment. <i>BMC Public Health</i> . 2016. 16:734. doi:10.1186/s12889-016-3299-z.	Study Design; Outcome
135 Liao Y, McGee, DL, Cao G, Cooper RS. Alcohol intake and mortality: findings from the National Health Interview Surveys (1988 and 1990). <i>Am J Epidemiol</i> . 2000. 151(7):651-9. doi:10.1093/oxfordjournals.aje.a010259.	Publication Date

	Citation	Rationale
136	Lin Y, Kikuchi S, Tamakoshi A, Wakai K, Kawamura T, Iso H, Ogimoto I, Yagyu K, Obata Y, Ishibashi T. Alcohol consumption and mortality among middle-aged and elderly Japanese men and women. <i>Ann Epidemiol.</i> 2005. 15(8):590-7. doi:10.1016/j.annepidem.2004.10.010.	Publication Date
137	Makela P, Paljarvi T, Poikolainen K. Heavy and nonheavy drinking occasions, all-cause and cardiovascular mortality and hospitalizations: a follow-up study in a population with a low consumption level. <i>J Stud Alcohol.</i> 2005. 66(6):722-8. doi:10.15288/jsa.2005.66.722.	Comparator
138	Malinski MK, Sesso HD, Lopez-Jimenez F, Buring JE, Gaziano JM. Alcohol consumption and cardiovascular disease mortality in hypertensive men. <i>Arch Intern Med.</i> 2004. 164(6):623-8. doi:10.1001/archinte.164.6.623.	Publication Date
139	Malyutina S, Bobak M, Kurilovitch S, Gafarov V, Simonova G, Nikitin Y, Marmot M. Relation between heavy and binge drinking and all-cause and cardiovascular mortality in Novosibirsk, Russia: a prospective cohort study. <i>Lancet.</i> 2002. 360(9344):1448-54. doi:10.1016/s0140-6736(02)11470-x.	Publication Date
140	Manuel DG, Perez R, Sanmartin C, Taljaard M, Hennessy D, Wilson K, Tanuseputro P, Manson H, Bennett C, Tuna M, Fisher S, Rosella LC. Measuring Burden of Unhealthy Behaviours Using a Multivariable Predictive Approach: Life Expectancy Lost in Canada Attributable to Smoking, Alcohol, Physical Inactivity, and Diet. <i>PLoS Med.</i> 2016. 13(8):e1002082. doi:10.1371/journal.pmed.1002082.	Comparator
141	Maraldi C, Volpato S, Kritchevsky SB, Cesari M, Andresen E, Leeuwenburgh C, Harris TB, Newman AB, Kanaya A, Johnson KC, Rodondi N, Pahor M. Impact of inflammation on the relationship among alcohol consumption, mortality, and cardiac events: the health, aging, and body composition study. <i>Arch Intern Med.</i> 2006. 166(14):1490-7. doi:10.1001/archinte.166.14.1490.	Publication Date
142	Marteau TM, White M, Rutter H, Petticrew M, Mytton OT, McGowan JG, Aldridge RW. Increasing healthy life expectancy equitably in England by 5 years by 2035: could it be achieved?. <i>Lancet.</i> 2019. 393(10191):2571-2573. doi:10.1016/s0140-6736(19)31510-7.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
143	Martikainen P, Makela P, Peltonen R, Myrskylä M. Income differences in life expectancy: the changing contribution of harmful consumption of alcohol and smoking. <i>Epidemiology.</i> 2014. 25(2):182-90. doi:10.1097/ede.0000000000000064.	Intervention/Exposure; Outcome
144	Martin J, Barry J, Goggin D, Morgan K, Ward M, O'Suilleabhain T. Alcohol-attributable mortality in Ireland. <i>Alcohol Alcohol.</i> 2010. 45(4):379-86. doi:10.1093/alcalc/agg032.	Intervention/Exposure; Outcome
145	Martin-Diener E, Meyer J, Braun J, Tarnutzer S, Faeh D, Rohrmann S, Martin BW. The combined effect on survival of four main behavioural risk factors for non-communicable diseases. <i>Prev Med.</i> 2014. 65:148-52. doi:10.1016/j.ypmed.2014.05.023.	Comparator
146	Marugame T, Yamamoto S, Yoshimi I, Sobue T, Inoue M, Tsugane S. Patterns of alcohol drinking and all-cause mortality: results from a large-scale population-based cohort study in Japan. <i>Am J Epidemiol.</i> 2007. 165(9):1039-46. doi:10.1093/aje/kwk112.	Publication Date
147	McCartney DL, Hillary RF, Stevenson AJ, Ritchie SJ, Walker RM, Zhang Q, Morris SW, Birmingham ML, Campbell A, Murray AD, Whalley HC, Gale CR, Porteous DJ, Haley CS, McRae AF, Wray NR, Visscher PM, McIntosh AM, Evans KL, Deary IJ, Marioni RE. Epigenetic prediction of complex traits and death. <i>Genome Biol.</i> 2018. 19(1):136. doi:10.1186/s13059-018-1514-1.	Intervention/Exposure; Outcome
148	McCartney G, Shipley M, Hart C, Davey-Smith G, Kivimäki M, Walsh D, Watt GC, Batty GD. Why do males in Scotland die younger than those in England? evidence from three prospective cohort studies. <i>PLoS ONE.</i> 2012. 7(7). doi:10.1371/journal.pone.0038860.	Study Design; Intervention/Exposure
149	McCaul KA, Almeida OP, Hankey GJ, Jamrozik K, Byles JE, Flicker L. Alcohol use and mortality in older men and women. <i>Addiction.</i> 2010. 105(8):1391-400. doi:10.1111/j.1360-0443.2010.02972.x.	Comparator
150	Mehta NK, House JS, Elliott MR. Dynamics of health behaviours and socioeconomic differences in mortality in the USA. <i>J Epidemiol Community Health.</i> 2015. 69(5):416-22. doi:10.1136/jech-2014-204248.	Intervention/Exposure
151	Melo APS, Franca EB, Malta DC, Garcia LP, Mooney M, Naghavi M. Mortality due to cirrhosis, liver cancer, and disorders attributed to alcohol use: Global Burden of Disease in Brazil, 1990 and 2015. <i>Rev Bras Epidemiol.</i> 2017. 20Suppl 01(Suppl 01):61-74. doi:10.1590/1980-5497201700050006.	Outcome

	Citation	Rationale
152	Meyer J, Doring A, Herder C, Roden M, Koenig W, Thorand B. Dietary patterns, subclinical inflammation, incident coronary heart disease and mortality in middle-aged men from the MONICA/KORA Augsburg cohort study. <i>Eur J Clin Nutr.</i> 2011. 65(7):800-7. doi:10.1038/ejcn.2011.37.	Intervention/Exposure
153	Mills JA, Beach, SRH, Dogan M, Simons, RL, Gibbons, FX, Long, JD, Philibert, R. A Direct Comparison of the Relationship of Epigenetic Aging and Epigenetic Substance Consumption Markers to Mortality in the Framingham Heart Study. <i>Genes (Basel).</i> 2019. 10(1). doi:10.3390/genes10010051.	Intervention/Exposure
154	Millwood IY, Walters RG, Mei XW, Guo Y, Yang L, Bian Z, Bennett DA, Chen Y, Dong C, Hu R, Zhou G, Yu B, Jia W, Parish S, Clarke R, Davey Smith G, Collins R, Holmes MV, Li L, Peto R, Chen, Z. Conventional and genetic evidence on alcohol and vascular disease aetiology: a prospective study of 500 000 men and women in China. <i>Lancet.</i> 2019. 393(10183):1831-1842. doi:10.1016/s0140-6736(18)31772-0.	Country
155	Mink PJ, Scrafford CG, Barraj LM, Harnack L, Hong CP, Nettleton JA, Jacobs, DR, Jr. Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. <i>Am J Clin Nutr.</i> 2007. 85(3):895-909. doi:10.1093/ajcn/85.3.895.	Outcome
156	Miyazaki M, Ue H. Japanese alcoholic beverage and all cause mortality in Japanese adult men. <i>J Epidemiol.</i> 2001. 11(5):219-23. doi:10.2188/jea.11.219.	Publication Date
157	Moore AA, Giuli L, Gould R, Hu P, Zhou K, Reuben D, Greendale G, Karlamangla A. Alcohol use, comorbidity, and mortality. <i>J Am Geriatr Soc.</i> 2006. 54(5):757-62. doi:10.1111/j.1532-5415.2006.00728.x.	Publication Date
158	Morc, LS, Johansen D, Lokkegaard E, Hundrup YA, Gronbaek M. Drinking pattern and mortality in Danish nurses. <i>Eur J Clin Nutr.</i> 2008. 62(6):817-22. doi:10.1038/sj.ejcn.1602799.	Publication Date
159	Mukamal KJ, Chen CM, Rao SR, Breslow RA. Alcohol consumption and cardiovascular mortality among U.S. adults, 1987 to 2002. <i>J Am Coll Cardiol.</i> 2010. 55(13):1328-35. doi:10.1016/j.jacc.2009.10.056.	Outcome
160	Murray RP, Connett JE, Tyas SL, Bond R, Ekuma O, Silversides CK, Barnes GE. Alcohol volume, drinking pattern, and cardiovascular disease morbidity and mortality: is there a U-shaped function?. <i>Am J Epidemiol.</i> 2002. 155(3):242-8. doi:10.1093/aje/155.3.242.	Outcome
161	Muscari A, Bianchi G, Conte C, Forti P, Magalotti D, Pandolfi P, Vaccheri A, Zoli M. No Direct Survival Effect of Light to Moderate Alcohol Drinking in Community-Dwelling Older Adults. <i>J Am Geriatr Soc.</i> 2015. 63(12):2526-2533. doi:10.1111/jgs.13837.	Comparator
162	Nagy J, Sipka S, Sipka Jr S, Kocsis J, Horvath Z. The Hardness of Drinking Water Negatively while Socio-Economic Deprivation Positively Correlate with the Age-Adjusted Mortality Rates due to Cardiovascular Diseases in Hungarian Wine Regions. <i>Int J Environ Res Public Health.</i> 2019. 16(18). doi:10.3390/ijerph16183437.	Intervention/Exposure; Outcome
163	Naimi TS, Xuan Z, Brown DW, Saitz R. Confounding and studies of 'moderate' alcohol consumption: the case of drinking frequency and implications for low-risk drinking guidelines. <i>Addiction.</i> 2013. 108(9):1534-43. doi:10.1111/j.1360-0443.2012.04074.x.	Outcome
164	Nakamura Y, Ueshima H, Kadota A, Hozawa A, Okamura T, Kadowaki S, Kadowaki T, Hayakawa T, Kita Y, Abbott RD, Okayama A. Alcohol intake and 19-year mortality in diabetic men: NIPPON DATA80. <i>Alcohol.</i> 2009. 43(8):635-41. doi:10.1016/j.alcohol.2009.06.005.	Publication Date
165	Nakaya N, Kurashima K, Yamaguchi J, Ohkubo T, Nishino Y, Tsubono Y, Shibuya D, Fukudo S, Fukao A, Tsuji I, Hisamichi S. Alcohol consumption and mortality in Japan: the Miyagi Cohort Study. <i>J Epidemiol.</i> 2004. 14 Suppl 1:S18-25. doi:10.2188/jea.14.s18.	Publication Date
166	Nandi A, Glymour MM, Subramanian SV. Association among socioeconomic status, health behaviors, and all-cause mortality in the United States. <i>Epidemiology.</i> 2014. 25(2):170-7. doi:10.1097/ede.0000000000000038.	Comparator
167	Nelson MR, Alkhateeb AN, Ryan P, Willson K, Gartlan JG, Reid CM. Physical activity, alcohol and tobacco use and associated cardiovascular morbidity and mortality in the Second Australian National Blood Pressure study cohort. <i>Age Ageing.</i> 2010. 39(1):112-6. doi:10.1093/ageing/afp186.	Outcome
168	Nicholson A, Bobak M, Murphy M, Rose R, Marmot M. Alcohol consumption and increased mortality in Russian men and women: a cohort study based on the mortality of relatives. <i>Bull World Health Organ.</i> 2005. 83(11):812-9. doi:s0042-96862005001100010.	Publication Date

	Citation	Rationale
169	Nielsen NR, Schnohr P, Jensen G, Gronbaek M. Is the relationship between type of alcohol and mortality influenced by socio-economic status?. <i>J Intern Med</i> . 2004. 255(2):280-8. doi:10.1046/j.1365-2796.2003.01268.x.	Publication Date
170	Nielsen NR, Thygesen LC, Johansen D, Jensen G, Gronbaek M. The influence of duration of follow-up on the association between alcohol and cause-specific mortality in a prospective cohort study. <i>Ann Epidemiol</i> . 2005. 15(1):44-55. doi:10.1016/j.annepidem.2004.04.002.	Publication Date
171	Nordahl H, Lange T, Osler M, Diderichsen F, Andersen I, Prescott E, Tjonneland A, Frederiksen BL, Rod NH. Education and cause-specific mortality: the mediating role of differential exposure and vulnerability to behavioral risk factors. <i>Epidemiology</i> . 2014. 25(3):389-96. doi:10.1097/ede.000000000000080.	Intervention/Exposure; Other (e.g., duplicative data, not primary research)
172	Norstrom T, Razvodovsky Y. Per capita alcohol consumption and alcohol-related harm in Belarus, 1970-2005. <i>Eur J Public Health</i> . 2010. 20(5):564-8. doi:10.1093/eurpub/ckq011.	Intervention/Exposure; Comparator
173	Nybo H, Petersen HC, Gaist D, Jeune B, Andersen K, McGue M, Vaupel JW, Christensen K. Predictors of mortality in 2,249 nonagenarians--the Danish 1905-Cohort Survey. <i>J Am Geriatr Soc</i> . 2003. 51(10):1365-73. doi:10.1046/j.1532-5415.2003.51453.x.	Publication Date
174	Odegaard AO, Koh WP, Gross MD, Yuan JM, Pereira MA. Combined lifestyle factors and cardiovascular disease mortality in Chinese men and women: the Singapore Chinese health study. <i>Circulation</i> . 2011. 124(25):2847-54. doi:10.1161/circulationaha.111.048843.	Intervention/Exposure; Outcome
175	Odegaard AO, Koh WP, Yuan JM, Pereira MA. Beverage habits and mortality in Chinese adults. <i>J Nutr</i> . 2015. 145(3):595-604. doi:10.3945/jn.114.200253.	Intervention/Exposure; Comparator
176	Onat A, Hergenc G, Kucukdurmaz Z, Ugur M, Kaya Z, Can G, Yuksel H. Moderate and heavy alcohol consumption among Turks: long-term impact on mortality and cardiometabolic risk. <i>Turk Kardiyol Dern Ars</i> . 2009. 37(2):83-90.	Country
177	Ostergren O, Martikainen P, Tarkiainen L, Elstad JI, Bronnum-Hansen H. Contribution of smoking and alcohol consumption to income differences in life expectancy: evidence using Danish, Finnish, Norwegian and Swedish register data. <i>J Epidemiol Community Health</i> . 2019. 73(4):334-339. doi:10.1136/jech-2018-211640.	Study Design; Intervention/Exposure; Outcome
178	Ozasa K. Alcohol use and mortality in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). i. 2007. 8 Suppl:81-8.	Publication Date
179	Paganini-Hill A, Kawas CH, Corrada MM. Type of alcohol consumed, changes in intake over time and mortality: the Leisure World Cohort Study. <i>Age Ageing</i> . 2007. 36(2):203-9. doi:10.1093/ageing/af184.	Publication Date
180	Paganini-Hill A. Lifestyle practices and cardiovascular disease mortality in the elderly: the leisure world cohort study. i. 2011. 2011:983764. doi:10.4061/2011/983764.	Outcome
181	Park S, Hong JP, Choi SH, Ahn MH. Clinical and laboratory predictors of all causes deaths and alcohol-attributable deaths among discharged alcohol-dependent patients. <i>Alcohol Clin Exp Res</i> . 2013. 37(2):270-5. doi:10.1111/j.1530-0277.2012.01943.x.	Intervention/Exposure; Health Status
182	Paschall M, Lipton, RI. Wine preference and related health determinants in a U.S. national sample of young adults. <i>Drug Alcohol Depend</i> . 2005. 78(3):339-44. doi:10.1016/j.drugalcdep.2004.12.004.	Outcome
183	Pedersen JO, Heitmann BL, Schnohr P, Gronbaek M. The combined influence of leisure-time physical activity and weekly alcohol intake on fatal ischaemic heart disease and all-cause mortality. <i>Eur Heart J</i> . 2008. 29(2):204-12. doi:10.1093/eurheartj/ehm574.	Publication Date
184	Perez-Perez E, Cruz-Lopez L, Hernandez-Llanes NF, Gallegos-Cari A, Camacho-Solis RE, Mendoza-Melendez MA. Years of life lost (Yll) attributable to alcohol consumption in Mexico City. <i>Cien Saude Colet</i> . 2016. 21(1):37-44. doi:10.1590/1413-81232015211.09472015.	Intervention/Exposure
185	Perlman F, Bobak M. Socioeconomic and behavioral determinants of mortality in posttransition Russia: a prospective population study. <i>Ann Epidemiol</i> . 2008. 18(2):92-100. doi:10.1016/j.annepidem.2007.07.093.	Publication Date
186	Perreira KM, Sloan FA. Excess alcohol consumption and health outcomes: a 6-year follow-up of men over age 50 from the health and retirement study. <i>Addiction</i> . 2002. 97(3):301-10. doi:10.1046/j.1360-0443.2002.00067.x.	Publication Date

	Citation	Rationale
187	Petersen KE, Johnsen NF, Olsen A, Albieri V, Olsen LK, Dragsted LO, Overvad K, Tjønneland A, Egeberg R. The combined impact of adherence to five lifestyle factors on all-cause, cancer and cardiovascular mortality: a prospective cohort study among Danish men and women. <i>Br J Nutr</i> . 2015. 113(5):849-58. doi:10.1017/s0007114515000070.	Intervention/Exposure
188	Plavinski SL, Plavinskaya SI, Klimov AN. Social factors and increase in mortality in Russia in the 1990s: prospective cohort study. <i>Bmj</i> . 2003. 326(7401):1240-2. doi:10.1136/bmj.326.7401.1240.	Intervention/Exposure; Other (e.g., duplicative data, not primary research)
189	Prinelli F, Yannakoulia M, Anastasiou CA, Adorni F, Di Santo SG, Musicco M, Scarmeas N, Correa Leite ML. Mediterranean diet and other lifestyle factors in relation to 20-year all-cause mortality: A cohort study in an Italian population. <i>British Journal of Nutrition</i> . 2015. 113(6):1003-1011. doi:10.1017/S0007114515000318.	Comparator
190	Probst C, Rehm J. Alcohol use, opioid overdose and socioeconomic status in Canada: A threat to life expectancy?. <i>Cmaj</i> . 2018. 190(44):E1294-e1295. doi:10.1503/cmaj.180806.	Study Design; Other (e.g., duplicative data, not primary research)
191	Rao TS, Andrade C. Alcohol intake, morbidity, and mortality. <i>Indian J Psychiatry</i> . 2016. 58(1):1-3. doi:10.4103/0019-5545.174352.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
192	Razvodovsky YE. Alcohol attributable mortality in Belarus. <i>Alcoholism</i> . 2012. 48(1):13-22.	Intervention/Exposure
193	Razvodovsky YE. Estimation of alcohol attributable fraction of mortality in Russia. <i>Adicciones</i> . 2012. 24(3):247-52.	Study Design; Intervention/Exposure
194	Rehm J, Greenfield TK, Rogers JD. Average volume of alcohol consumption, patterns of drinking, and all-cause mortality: results from the US National Alcohol Survey. <i>Am J Epidemiol</i> . 2001. 153(1):64-71. doi:10.1093/aje/153.1.64.	Publication Date
195	Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. <i>Lancet</i> . 2009. 373(9682):2223-33. doi:10.1016/s0140-6736(09)60746-7.	Study Design; Intervention/Exposure
196	Rehm J, Monteiro M. Alcohol consumption and burden of disease in the Americas: implications for alcohol policy. <i>Rev Panam Salud Publica</i> . 2005. 18(4-5):241-8. doi:10.1590/s1020-49892005000900003.	Intervention/Exposure; Outcome
197	Rehm J, Patra J, Taylor, B. Harm, Benefits, and Net Effects on Mortality of Moderate Drinking of Alcohol Among Adults in Canada in 2002. <i>Annals of Epidemiology</i> . 2007. 17(5 SUPPL.):S81-S86. doi:10.1016/j.annepidem.2007.01.018.	Intervention/Exposure; Outcome
198	Rehm J, Room R, Monteiro M, Gmel G, Graham K, Rehn N, Sempos, CT, Jernigan, D. Alcohol as a risk factor for global burden of disease. <i>Eur Addict Res</i> . 2003. 9(4):157-64. doi:10.1159/000072222.	Intervention/Exposure; Outcome
199	Rehm J, Scafato, E. Indicators of alcohol consumption and attributable harm for monitoring and surveillance in European Union countries. <i>Addiction</i> . 2011. 106 Suppl 1:4-10. doi:10.1111/j.1360-0443.2010.03323.x.	Outcome
200	Rehm J, Sulkowska U, Manczuk M, Boffetta P, Powles J, Popova S, Zatonski, W. Alcohol accounts for a high proportion of premature mortality in central and eastern Europe. <i>Int J Epidemiol</i> . 2007. 36(2):458-67. doi:10.1093/ije/dyl294.	Intervention/Exposure; Outcome
201	Reile R, Stickley A, Leinsalu, M. Large variation in predictors of mortality by levels of self-rated health: Results from an 18-year follow-up study. <i>Public Health</i> . 2017. 145:59-66. doi:10.1016/j.puhe.2016.12.034.	Intervention/Exposure; Comparator
202	Renaud S, Lanzmann-Petithory D, Gueguen R, Conard, P. Alcohol and mortality from all causes. <i>Biol Res</i> . 2004. 37(2):183-7. doi:10.4067/s0716-97602004000200002.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
203	Renaud SC, Gueguen R, Conard P, Lanzmann-Petithory D, Orgogozo JM, Henry O. Moderate wine drinkers have lower hypertension-related mortality: a prospective cohort study in French men. <i>Am J Clin Nutr</i> . 2004. 80(3):621-5. doi:10.1093/ajcn/80.3.621.	Comparator
204	Rey G, Boniol M, Jouglu E. Estimating the number of alcohol-attributable deaths: methodological issues and illustration with French data for 2006. <i>Addiction</i> . 2010. 105(6):1018-29. doi:10.1111/j.1360-0443.2010.02910.x.	Outcome

	Citation	Rationale
205	Ricci C, Wood A, Muller D, Gunter MJ, Agudo A, Boeing H, van der Schouw YT, Warnakula S, Saieva C, Spijkerman A, Sluijs I, Tjonneland A, Kyro C, Weiderpass E, Kuhn T, Kaaks R, Sanchez MJ, Panico S, Agnoli C, Palli D, Tumino R, Engstrom G, Melander O, Bonnet F, Boer JMA, Key TJ, Travis RC, Overvad K, Verschuren WMM, Quiros JR, Trichopoulou A, Papatesta, EM, Peppas E, Iribas CM, Gavrila D, Forslund AS, Jansson JH, Matullo G, Arriola L, Freisling H, Lassale C, Tzoulaki I, Sharp SJ, Forouhi NG, Langenberg C, Saracchi R, Sweeting M, Brennan P, Butterworth AS, Riboli E, Wareham NJ, Danesh J, Ferrari P. Alcohol intake in relation to non-fatal and fatal coronary heart disease and stroke: EPIC-CVD case-cohort study. <i>Bmj</i> . 2018. 361:k934. doi:10.1136/bmj.k934.	Outcome
206	Ringmets I, Tuusov J, Lang K, Vali M, Parna K, Tonisson M, Helander A, McKee M, Leon DA. Alcohol and premature death in Estonian men: a study of forensic autopsies using novel biomarkers and proxy informants. <i>BMC Public Health</i> . 2012. 12:146. doi:10.1186/1471-2458-12-146.	Intervention/Exposure; Outcome
207	Rizzuto D, Orsini N, Qiu C, Wang HX, Fratiglioni L. Lifestyle, social factors, and survival after age 75: population based study. <i>Bmj</i> . 2012. 345:e5568. doi:10.1136/bmj.e5568.	Comparator
208	Robinson M, Shipton D, Walsh D, Whyte B, McCartney G. Regional alcohol consumption and alcohol-related mortality in Great Britain: novel insights using retail sales data. <i>BMC Public Health</i> . 2015. 15:1. doi:10.1186/1471-2458-15-1.	Intervention/Exposure
209	Rogers RG, Boardman JD, Pendergast PM, Lawrence EM. Drinking problems and mortality risk in the United States. <i>Drug Alcohol Depend</i> . 2015. 151:38-46. doi:10.1016/j.drugalcdep.2015.02.039.	Intervention/Exposure
210	Romelsjo A, Allebeck P, Andreasson S, Leifman A. Alcohol, mortality and cardiovascular events in a 35 year follow-up of a nationwide representative cohort of 50,000 Swedish conscripts up to age 55. <i>Alcohol Alcohol</i> . 2012. 47(3):322-7. doi:10.1093/alcalc/ags021.	Comparator
211	Rosen M, Haglund B. Follow-up of an age-period-cohort analysis on alcohol-related mortality trends in Sweden 1970-2015 with predictions to 2025. <i>Scand J Public Health</i> . 2019. 47(4):446-451. doi:10.1177/1403494817752521.	Intervention/Exposure
212	Ruf E, Baumert J, Meisinger C, Doring A, Ladwig KH. Are psychosocial stressors associated with the relationship of alcohol consumption and all-cause mortality?. <i>BMC Public Health</i> . 2014. 14:312. doi:10.1186/1471-2458-14-312.	Comparator
213	Ryu M, Gombojav B, Nam CM, Lee Y, Han K. Modifying effects of resting heart rate on the association of binge drinking with all-cause and cardiovascular mortality in older Korean men: the Kangwha Cohort Study. <i>J Epidemiol</i> . 2014. 24(4):274-80. doi:10.2188/jea.je20130101.	Comparator
214	Sadakane A, Gotoh T, Ishikawa S, Nakamura Y, Kayaba K. Amount and frequency of alcohol consumption and all-cause mortality in a Japanese population: the JMS Cohort Study. <i>J Epidemiol</i> . 2009. 19(3):107-15. doi:10.2188/jea.je20081003.	Publication Date
215	Saito E, Inoue M, Sawada N, Charvat H, Shimazu T, Yamaji T, Iwasaki M, Sasazuki S, Mizoue T, Iso H, Tsugane S. Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. <i>J Epidemiol</i> . 2018. 28(3):140-148. doi:10.2188/jea.JE20160200.	Intervention/Exposure
216	Sakaue S, Akiyama M, Hirata M, Matsuda K, Murakami Y, Kubo M, Kamatani Y, Okada Y. Functional variants in ADH1B and ALDH2 are non-additively associated with all-cause mortality in Japanese population. <i>Eur J Hum Genet</i> . 2019. doi:10.1038/s41431-019-0518-y.	Intervention/Exposure
217	Schnohr C, Hojbjerg L, Riegels M, Ledet L, Larsen T, Schultz-Larsen K, Petersen L, Prescott E, Gronbaek M. Does educational level influence the effects of smoking, alcohol, physical activity, and obesity on mortality? A prospective population study. <i>Scand J Public Health</i> . 2004. 32(4):250-6. doi:10.1080/14034940310019489.	Publication Date
218	Schooling CM, Sun W, Ho SY, Chan WM, Tham MK, Ho KS, Leung GM, Lam TH. Moderate alcohol use and mortality from ischaemic heart disease: a prospective study in older Chinese people. <i>PLoS One</i> . 2008. 3(6):e2370. doi:10.1371/journal.pone.0002370.	Outcome
219	Sempos CT, Rehm J, Wu T, Crespo CJ, Trevisan M. Average volume of alcohol consumption and all-cause mortality in African Americans: the NHEFS cohort. <i>Alcohol Clin Exp Res</i> . 2003. 27(1):88-92. doi:10.1097/01.Alc.0000046597.92232.73.	Publication Date



	Citation	Rationale
220	Sethi M, Heyer JH, Wall S, DiMaggio C, Shinseki M, Slaughter D, Frangos SG. Alcohol use by urban bicyclists is associated with more severe injury, greater hospital resource use, and higher mortality. <i>Alcohol</i> . 2016. 53:1-7. doi:10.1016/j.alcohol.2016.03.005.	Intervention/Exposure; Outcome
221	Shaw BA, Agahi N. A prospective cohort study of health behavior profiles after age 50 and mortality risk. <i>BMC public health</i> . 2012. 12:803.	Intervention/Exposure; Comparator
222	Shield KD, Gmel G, Gmel G, Makela P, Probst C, Room R, Rehm J. Life-time risk of mortality due to different levels of alcohol consumption in seven European countries: implications for low-risk drinking guidelines. <i>Addiction</i> . 2017. 112(9):1535-1544. doi:10.1111/add.13827.	Intervention/Exposure
223	Shield KD, Gmel G, Kehoe-Chan T, Dawson DA, Grant BF, Rehm J. Mortality and potential years of life lost attributable to alcohol consumption by race and sex in the United States in 2005. <i>PLoS One</i> . 2013. 8(1):e51923. doi:10.1371/journal.pone.0051923.	Study Design; Intervention/Exposure; Outcome
224	Shield KD, Kehoe T, Taylor B, Patra J, Rehm J. Alcohol-attributable burden of disease and injury in Canada, 2004. <i>Int J Public Health</i> . 2012. 57(2):391-401. doi:10.1007/s00038-011-0247-7.	Intervention/Exposure; Outcome
225	Shield KD, Rehm J, Gmel G, Rehm MX, Allamani A. Alcohol consumption, alcohol dependence, and related mortality in Italy in 2004: effects of treatment-based interventions on alcohol dependence. <i>Subst Abuse Treat Prev Policy</i> . 2013. 8:21. doi:10.1186/1747-597x-8-21.	Intervention/Exposure; Comparator
226	Shield KD, Rehm J. Russia-specific relative risks and their effects on the estimated alcohol-attributable burden of disease. <i>BMC Public Health</i> . 2015. 15:482. doi:10.1186/s12889-015-1818-y.	Intervention/Exposure; Outcome
227	Shield KD, Taylor B, Kehoe T, Patra J, Rehm J. Mortality and potential years of life lost attributable to alcohol consumption in Canada in 2005. <i>BMC Public Health</i> . 2012. 12:91. doi:10.1186/1471-2458-12-91.	Intervention/Exposure; Comparator
228	Sidorenkov O, Nilssen O, Grijbovski AM. Determinants of cardiovascular and all-cause mortality in northwest Russia: a 10-year follow-up study. <i>Ann Epidemiol</i> . 2012. 22(1):57-65. doi:10.1016/j.annepidem.2011.08.008.	Comparator
229	Sidorenkov O, Nilssen O, Grijbovski AM. Metabolic syndrome in Russian adults: associated factors and mortality from cardiovascular diseases and all causes. <i>BMC Public Health</i> . 2010. 10:582. doi:10.1186/1471-2458-10-582.	Intervention/Exposure; Outcome
230	Simons LA, McCallum J, Friedlander Y, Ortiz M, Simons J. Moderate alcohol intake is associated with survival in the elderly: the Dubbo Study. <i>Med J Aust</i> . 2000. 173(3):121-4.	Publication Date
231	Simons LA, Simons J, Friedlander Y, McCallum J. Predictors of long-term mortality in the elderly: the Dubbo Study. <i>Intern Med J</i> . 2011. 41(7):555-60. doi:10.1111/j.1445-5994.2009.02106.x.	Intervention/Exposure; Comparator
232	Sipetic S, Bjegovic-Mikanovic V, Vlajinac H, Marinkovic J, Jankovic S, Terzic Z, Atanackovic-Markovic Z, Saulic A, Laaser U. The burden of disease preventable by risk factor reduction in Serbia. <i>Vojnosanit Pregl</i> . 2013. 70(5):445-51. doi:10.2298/vsp111024049s.	Outcome
233	Sipila P, Rose RJ, Kaprio J. Drinking and mortality: long-term follow-up of drinking-discordant twin pairs. <i>Addiction</i> . 2016. 111(2):245-54. doi:10.1111/add.13152.	Comparator
234	Smigielski J, Bielecki W, Drygas W. Health and life style-related determinants of survival rate in the male residents of the city of Lodz. <i>Int J Occup Med Environ Health</i> . 2013. 26(3):337-48. doi:10.2478/s13382-013-0104-4.	Comparator
235	Smigielski J, Jegier A, Hanke W, Bielecki W, Drygas W. The effect of selected lifestyle factors and diet on mortality of men with documented physical fitness in the city of Lodz. <i>Int J Occup Med Environ Health</i> . 2013. 26(4):535-44. doi:10.2478/s13382-013-0112-4.	Sample size
236	Sofi F, Abbate R, Gensini GF, Casini A, Trichopoulou A, Bamia C. Identification of change-points in the relationship between food groups in the Mediterranean diet and overall mortality: an 'a posteriori' approach. <i>Eur J Nutr</i> . 2012. 51(2):167-72. doi:10.1007/s00394-011-0202-7.	Intervention/Exposure; Outcome
237	Sotos-Prieto M, Mattei J, Cook NR, Hu FB, Willett WC, Chiuve SE, Rimm EB, Sesso HD. Association Between a 20-Year Cardiovascular Disease Risk Score Based on Modifiable Lifestyles and Total and Cause-Specific Mortality Among US Men and Women. <i>J Am Heart Assoc</i> . 2018. 7(21):e010052. doi:10.1161/jaha.118.010052.	Intervention/Exposure; Comparator

	Citation	Rationale
238	Sovic S, Vitale K, Brborovic O, Dzakula A, Tiljak H. Association of behavioral cardiovascular risk factors with mortality in Croatian adult population: the CroHort study. <i>Coll Antropol</i> . 2012. 36 Suppl 1:177-82. doi:10.5671/ca.2012361s.177.	Intervention/Exposure; Comparator
239	Stahre M, Roeber J, Kanny D, Brewer RD, Zhang X. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. <i>Prev Chronic Dis</i> . 2014. 11:E109. doi:10.5888/pcd11.130293.	Intervention/Exposure; Outcome
240	StatBite: Alcohol-related deaths from chronic causes among American Indians and Alaska Natives. <i>J Natl Cancer Inst</i> . 2008. 100(19):1349. doi:10.1093/jnci/djn350.	Study Design; Publication Status
241	Strandberg AY, Strandberg TE, Salomaa VV, Pitkala K, Miettinen TA. Alcohol consumption, 29-y total mortality, and quality of life in men in old age. <i>Am J Clin Nutr</i> . 2004. 80(5):1366-71. doi:10.1093/ajcn/80.5.1366.	Publication Date
242	Streppel MT, Ocke MC, Boshuizen HC, Kok FJ, Kromhout D. Long-term wine consumption is related to cardiovascular mortality and life expectancy independently of moderate alcohol intake: the Zutphen Study. <i>J Epidemiol Community Health</i> . 2009. 63(7):534-40. doi:10.1136/jech.2008.082198.	Publication Date
243	Stringhini S, Carmeli C, Jokela M, Avendaño M, Muennig P, Guida F, Ricceri F, d'Errico A, Barros H, Bochud M, Chadeau-Hyam M, Clavel-Chapelon F, Costa G, Delpierre C, Fraga S, Goldberg M, Giles GG, Krogh V, Kelly-Irving M, Layte R, Lasserre AM, Marmot MG, Preisig M, Shipley MJ, Vollenweider P, Zins M, Kawachi I, Steptoe A, Mackenbach JP, Vineis P, Kivimäki M, Alenius H, Avendano M, Carra L, Castagné R, Courtin E, Dugué PA, Elliott P, Gares V, Giles G, Greco D, Hodge A, Irving MK, Karisola P, Lang T, Lepage B, Mackenbach J, McCrory C, Milne R, Nusselder W, Panico S, Petrovic D, Polidoro S, Raitakari O, Ribeiro AI, Robinson O, Valverde JR, Sacerdote C, Satolli R, Severi G, Tumino R. Socioeconomic status and the 25 × 25 risk factors as determinants of premature mortality: a multicohort study and meta-analysis of 1.7 million men and women. <i>The Lancet</i> . 2017. 389(10075):1229-1237. doi:10.1016/S0140-6736(16)32380-7.	Comparator; Outcome
244	Stringhini S, Sabia S, Shipley M, Brunner E, Nabi H, Kivimäki M, Singh-Manoux A. Association of socioeconomic position with health behaviors and mortality. <i>Jama</i> . 2010. 303(12):1159-66. doi:10.1001/jama.2010.297.	Study design
245	Stringhini S, Zaninotto P, Kumari M, Kivimäki M, Lassale C, Batty GD. Socio-economic trajectories and cardiovascular disease mortality in older people: the English Longitudinal Study of Ageing. <i>Int J Epidemiol</i> . 2018. 47(1):36-46. doi:10.1093/ije/dyx106.	Comparator
246	Suadicani P, Hein HO, Gyntelberg F. Wine intake, ABO phenotype, and risk of ischemic heart disease and all-cause mortality: the Copenhagen Male Study--a 16-year follow-up. <i>Alcohol</i> . 2008. 42(7):575-82. doi:10.1016/j.alcohol.2008.05.007.	Publication Date
247	Suadicani P, Hein HO, von Eyben FE, Gyntelberg F. Metabolic and lifestyle predictors of ischemic heart disease and all-cause mortality among normal weight, overweight, and obese men: a 16-year follow-up in the Copenhagen Male Study. <i>Metab Syndr Relat Disord</i> . 2009. 7(2):97-104. doi:10.1089/met.2008.0041.	Intervention/Exposure
248	Sull JW, Yi SW, Nam CM, Ohrr H. Binge drinking and mortality from all causes and cerebrovascular diseases in Korean men and women: a Kangwha cohort study. <i>Stroke</i> . 2009. 40(9):2953-8. doi:10.1161/strokeaha.109.556027.	Publication Date
249	Sun W, Schooling CM, Chan WM, Ho KS, Lam TH, Leung GM. Moderate alcohol use, health status, and mortality in a prospective Chinese elderly cohort. <i>Ann Epidemiol</i> . 2009. 19(6):396-403. doi:10.1016/j.annepidem.2009.01.011.	Publication Date
250	Tae HK, Shin J, Noh S, Kyo JS, Park JK, Ahn JS, Min S. Alcohol-related mortality in a rural community in Korea over an eight-year period. <i>Psychiatry Investigation</i> . 2007. 4(2):96-103.	Publication Date
251	Taylor B, Rehm J, Room R, Patra J, Bondy S. Determination of lifetime injury mortality risk in Canada in 2002 by drinking amount per occasion and number of occasions. <i>Am J Epidemiol</i> . 2008. 168(10):1119-25; discussion 1126-31. doi:10.1093/aje/kwn215.	Intervention/Exposure; Outcome
252	Taylor BJ, Shield KD, Rehm JT. Combining best evidence: a novel method to calculate the alcohol-attributable fraction and its variance for injury mortality. <i>BMC Public Health</i> . 2011. 11:265. doi:10.1186/1471-2458-11-265.	Outcome
253	Taylor J. Effects of alcohol on cardiovascular disease: originally thought to be cardioprotective, the pendulum is now swinging against alcohol, with more deaths due to drinking than lives saved. <i>Eur Heart J</i> . 2013. 34(9):627-9.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
254	Theobald H, Bygren LO, Carstensen J, Engfeldt P. A moderate intake of wine is associated with reduced total mortality and reduced mortality from cardiovascular disease. <i>J Stud Alcohol</i> . 2000. 61(5):652-6. doi:10.15288/jsa.2000.61.652.	Publication Date

	Citation	Rationale
255	Theobald H, Johansson SE, Bygren LO, Engfeldt P. The effects of alcohol consumption on mortality and morbidity: a 26-year follow-up study. <i>J Stud Alcohol</i> . 2001. 62(6):783-9. doi:10.15288/jsa.2001.62.783.	Publication Date
256	Thomson CA, McCullough ML, Wertheim BC, Chlebowski RT, Martinez ME, Stefanick ML, Rohan TE, Manson JE, Tindle HA, Ockene J, Vitolins MZ, Wactawski-Wende J, Sarto GE, Lane DS, Neuhaus ML. Nutrition and physical activity cancer prevention guidelines, cancer risk, and mortality in the women's health initiative. <i>Cancer Prev Res (Phila)</i> . 2014. 7(1):42-53. doi:10.1158/1940-6207.Capr-13-0258.	Intervention/Exposure
257	Thorpe RJ, Jr, Wilson-Frederick SM, Bowie JV, Coa K, Clay OJ, LaVeist TA, Whitfield KE. Health behaviors and all-cause mortality in African American men. <i>Am J Mens Health</i> . 2013. 7(4 Suppl):8s-18s. doi:10.1177/1557988313487552.	Comparator
258	Thygesen LC, Johansen C, Keiding N, Giovannucci E, Gronbaek M. Effects of sample attrition in a longitudinal study of the association between alcohol intake and all-cause mortality. <i>Addiction</i> . 2008. 103(7):1149-59. doi:10.1111/j.1360-0443.2008.02241.x.	Publication Date
259	Tognon G, Lissner L, Saebye D, Walker KZ, Heitmann BL. The Mediterranean diet in relation to mortality and CVD: a Danish cohort study. <i>Br J Nutr</i> . 2014. 111(1):151-9. doi:10.1017/s0007114513001931.	Comparator
260	Tolstrup JS, Jensen MK, Tjonneland A, Overvad K, Gronbaek M. Drinking pattern and mortality in middle-aged men and women. <i>Addiction</i> . 2004. 99(3):323-30. doi:10.1111/j.1360-0443.2003.00667.x.	Publication Date
261	Tolvanen E, Seppa K, Lintonen T, Paavilainen P, Jylha M. Old people, alcohol use and mortality. A ten-year prospective study. <i>Aging Clin Exp Res</i> . 2005. 17(5):426-33. doi:10.1007/bf03324633.	Publication Date
262	Tomkins S, Collier T, Oralov A, Saburova L, McKee M, Shkolnikov V, Kiryanov N, Leon DA. Hazardous alcohol consumption is a major factor in male premature mortality in a typical Russian city: prospective cohort study 2003-2009. <i>PLoS One</i> . 2012. 7(2):e30274. doi:10.1371/journal.pone.0030274.	Intervention/Exposure
263	Trevisan M, Schisterman E, Mennotti A, Farchi G, Conti S. Drinking pattern and mortality: the Italian Risk Factor and Life Expectancy pooling project. <i>Ann Epidemiol</i> . 2001. 11(5):312-9. doi:10.1016/s1047-2797(00)00183-6.	Intervention/Exposure; Comparator
264	Trias-Llimos S, Janssen F. Alcohol and gender gaps in life expectancy in eight Central and Eastern European countries. <i>Eur J Public Health</i> . 2018. 28(4):687-692. doi:10.1093/eurpub/cky057.	Intervention/Exposure
265	Trias-Llimos S, Martikainen P, Makela P, Janssen F. Comparison of different approaches for estimating age-specific alcohol-attributable mortality: The cases of France and Finland. <i>PLoS One</i> . 2018. 13(3):e0194478. doi:10.1371/journal.pone.0194478.	Intervention/Exposure
266	Trichopoulou A, Bamia C, Trichopoulos D. Anatomy of health effects of Mediterranean diet: Greek EPIC prospective cohort study. <i>Bmj</i> . 2009. 338:b2337. doi:10.1136/bmj.b2337.	Publication Date
267	Tsubono Y, Koizumi Y, Nakaya N, Fujita K, Takahashi H, Hozawa A, Suzuki Y, Kuriyama S, Tsuji I, Fukao A, Hisamichi S. Health practices and mortality in Japan: combined effects of smoking, drinking, walking and body mass index in the Miyagi Cohort Study. <i>J Epidemiol</i> . 2004. 14 Suppl 1:S39-45. doi:10.2188/jea.14.s39.	Intervention/Exposure
268	Turiano NA, Chapman BP, Gruenewald TL, Mroczek DK. Personality and the leading behavioral contributors of mortality. <i>Health Psychol</i> . 2015. 34(1):51-60. doi:10.1037/hea0000038.	Intervention/Exposure
269	Tverdal A, Magnus P, Selmer R, Thelle D. Consumption of alcohol and cardiovascular disease mortality: a 16 year follow-up of 115,592 Norwegian men and women aged 40-44 years. <i>Eur J Epidemiol</i> . 2017. 32(9):775-783. doi:10.1007/s10654-017-0313-4.	Outcome
270	Urtamo A, Jyväkorpi SK, Kautiainen H, Pitkälä KH, Strandberg TE. Major cardiovascular disease (CVD) risk factors in midlife and extreme longevity. <i>Aging Clinical and Experimental Research</i> . 2020. 32(2):299-304. doi:10.1007/s40520-019-01364-7.	Intervention/exposure; Sample size
271	van Dam RM, Li T, Spiegelman D, Franco OH, Hu FB. Combined impact of lifestyle factors on mortality: prospective cohort study in US women. <i>Bmj</i> . 2008. 337:a1440. doi:10.1136/bmj.a1440.	Publication Date
272	Veronese N, Li Y, Manson JE, Willett WC, Fontana L, Hu FB. Combined associations of body weight and lifestyle factors with all cause and cause specific mortality in men and women: prospective cohort study. <i>Bmj</i> . 2016. 355:i5855. doi:10.1136/bmj.i5855.	Intervention/Exposure

	Citation	Rationale
273	Vetrano DL, Collamati A, Magnavita N, Sowa A, Topinkova E, Finne-Soveri H, van der Roest HG, Tobiasz-Adamczyk B, Giovannini S, Ricciardi W, Bernabei R, Onder G, Poscia A. Health determinants and survival in nursing home residents in Europe: Results from the SHELTER study. <i>Maturitas</i> . 2018. 107:19-25. doi:10.1016/j.maturitas.2017.09.014.	Intervention/Exposure; Health Status
274	Walker S. Russia's alcohol problem. <i>Bmj</i> . 2011. 343:d5240. doi:10.1136/bmj.d5240.	Study Design
275	Wannamethee SG, Shaper AG. Taking up regular drinking in middle age: effect on major coronary heart disease events and mortality. <i>Heart</i> . 2002. 87(1):32-6. doi:10.1136/heart.87.1.32.	Publication Date
276	Waskiewicz A, Sygnowska E, Drygas W. Relationship between alcohol consumption and cardiovascular mortality--the Warsaw Pol-MONICA Project. <i>Kardiol Pol</i> . 2004. 60(6):552-62; discussion 563.	Publication Date
277	Wellmann J, Heidrich J, Berger K, Doring A, Heuschmann PU, Keil U. Changes in alcohol intake and risk of coronary heart disease and all-cause mortality in the MONICA/KORA-Augsburg cohort 1987-97. <i>Eur J Cardiovasc Prev Rehabil</i> . 2004. 11(1):48-55. doi:10.1097/01.hjr.0000118174.70522.20.	Publication Date
278	White J, Greene G, Kivimaki M, Batty GD. Association between changes in lifestyle and all-cause mortality: the Health and Lifestyle Survey. <i>J Epidemiol Community Health</i> . 2018. 72(8):711-714. doi:10.1136/jech-2017-210363.	Intervention/Exposure
279	Whitfield JB, Heath AC, Madden PAF, Landers JG, Martin NG. Effects of high alcohol intake, alcohol-related symptoms and smoking on mortality. <i>Addiction</i> . 2018. 113(1):158-166. doi:10.1111/add.14008.	Comparator
280	Whitley E, Batty GD, Hunt K, Popham F, Benzeval M. The role of health behaviours across the life course in the socioeconomic patterning of all-cause mortality: the west of Scotland twenty-07 prospective cohort study. <i>Ann Behav Med</i> . 2014. 47(2):148-57. doi:10.1007/s12160-013-9539-x.	Intervention/Exposure
281	Williams EC, Bradley KA, Gupta S, Harris AHS. Association between alcohol screening scores and mortality in Black, Hispanic, and White male veterans. <i>Alcoholism: Clinical and Experimental Research</i> . 2012. 36(12):2132-2140. doi:10.1111/j.1530-0277.2012.01842.x.	Intervention/Exposure
282	Wojtyniak B, Moskalewicz J, Stokwiszewski J, Rabczenko D. Gender-specific mortality associated with alcohol consumption in Poland in transition. <i>Addiction</i> . 2005. 100(12):1779-89. doi:10.1111/j.1360-0443.2005.01247.x.	Intervention/Exposure; Outcome
283	Woo J, Ho SC, Yu AL. Lifestyle factors and health outcomes in elderly Hong Kong Chinese aged 70 years and over. <i>Gerontology</i> . 2002. 48(4):234-40. doi:10.1159/000058356.	Publication Date
284	Wood AM, Kaptoge S, Butterworth AS, Willeit P, Warnakula S, Bolton T, Paige E, Paul DS, Sweeting M, Burgess S, Bell S, Astle W, Stevens D, Koulman A, Selmer RM, Verschuren WMM, Sato S, Njølstad I, Woodward M, Salomaa V, Nordestgaard BG, Yeap BB, Fletcher A, Melander O, Kuller LH, Balkau B, Marmot M, Koenig W, Casiglia E, Cooper C, Arndt V, Franco OH, Wennberg P, Gallacher J, de la Cámara AG, Völzke H, Dahm CC, Dale CE, Bergmann MM, Crespo CJ, van der Schouw YT, Kaaks R, Simons LA, Lagiou P, Schoufour JD, Boer JMA, Key TJ, Rodriguez B, Moreno-Iribas C, Davidson KW, Taylor JO, Sacerdote C, Wallace RB, Quiros JR, Tumino R, Blazer DG, Linneberg A, Daimon M, Panico S, Howard B, Skeie G, Strandberg T, Weiderpass E, Psaty BM, Kromhout D, Salamanca-Fernandez E, Kiechl S, Krumholz HM, Grioni S, Palli D, Huerta JM, Price J, Sundström J, Arriola L, Arima H, Travis RC, Panagiotakos DB, Karakatsani A, Trichopoulou A, Kühn T, Grobbee DE, Barrett-Connor E, van Schoor N, Boeing H, Overvad K, Kauhanen J, Wareham N, Langenberg C, Forouhi N, Wennberg M, Després, JP, Cushman M, Cooper JA, Rodriguez CJ, Sakurai M, Shaw JE, Knuiman M, Voortman T, Meisinger C, Tjønneland A, Brenner H, Palmieri L, Dallongeville J, Brunner, EJ, Assmann G, Trevisan M, Gillum RF, Ford I, Sattar N, Lazo M, Thompson SG, Ferrari P, Leon, DA, Davey Smith G, Peto R, Jackson R, Banks E, Di Angelantonio E, Danesh J, Veikko S, Gómez de la Cámara A, Rimm EB, Dallongeville JP, Gillum RF, Ford IF, Thompson S. Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599 912 current drinkers in 83 prospective studies. <i>The Lancet</i> . 2018. 391(10129):1513-1523. doi:10.1016/S0140-6736(18)30134-X.	Duplicate
285	Woodside JV, Yarnell JW, Patterson CC, Arveiler D, Amouyel P, Ferrieres J, Kee F, Evans A, Bingham A, Ducimetiere P. Do lifestyle behaviours explain socioeconomic differences in all-cause mortality, and fatal and non-fatal cardiovascular events? Evidence from middle aged men in France and Northern Ireland in the PRIME Study. <i>Prev Med</i> . 2012. 54(3-4):247-53. doi:10.1016/j.ypmed.2012.01.017.	Intervention/Exposure

	Citation	Rationale
286	Wu CY, Chou YC, Huang N, Chou YJ, Hu HY, Li CP. Association of body mass index with all-cause and cardiovascular disease mortality in the elderly. <i>PLoS One</i> . 2014. 9(7):e102589. doi:10.1371/journal.pone.0102589.	Intervention/Exposure; Comparator
287	Wyshak G. Behavioral practices and mortality in women former college athletes and nonathletes. <i>Health Care Women Int</i> . 2003. 24(9):808-21. doi:10.1080/07399330390229920.	Intervention/Exposure
288	Xu WH, Zhang XL, Gao YT, Xiang YB, Gao LF, Zheng W, Shu XO. Joint effect of cigarette smoking and alcohol consumption on mortality. <i>Prev Med</i> . 2007. 45(4):313-9. doi:10.1016/j.ypmed.2007.05.015.	Publication Date
289	Yang L, Zhou M, Sherliker P, Cai Y, Peto R, Wang L, Millwood I, Smith M, Hu Y, Yang G, Chen, Z. Alcohol drinking and overall and cause-specific mortality in China: nationally representative prospective study of 220,000 men with 15 years of follow-up. <i>Int J Epidemiol</i> . 2012. 41(4):1101-13. doi:10.1093/ije/dys075.	Country
290	Yayci N, Agritmis H, Turla A, Koc, S. Fatalities due to methyl alcohol intoxication in Turkey: an 8-year study. <i>Forensic Sci Int</i> . 2003. 131(1):36-41. doi:10.1016/s0379-0738(02)00376-6.	Intervention/Exposure; Outcome; Country
291	Yi, SW, Yoo, SH, Sull, JW, Ohrr, H. Association between alcohol drinking and cardiovascular disease mortality and all-cause mortality: Kangwha Cohort Study. <i>J Prev Med Public Health</i> . 2004. 37(2):120-6.	Language
292	Zaccardi F, Franks PW, Dudbridge F, Davies MJ, Khunti K, Yates T. Mortality risk comparing walking pace to handgrip strength and a healthy lifestyle: A UK Biobank study. <i>Eur J Prev Cardiol</i> . 2019:2047487319885041. doi:10.1177/2047487319885041.	Other (e.g., duplicative data, not primary research)
293	Zaridze D, Lewington S, Boroda A, Scelo G, Karpov R, Lazarev A, Konobeevskaya I, Igitov V, Terechova T, Boffetta P, Sherliker P, Kong X, Whitlock G, Boreham J, Brennan P, Peto R. Alcohol and mortality in Russia: prospective observational study of 151,000 adults. <i>Lancet</i> . 2014. 383(9927):1465-1473. doi:10.1016/s0140-6736(13)62247-3.	Outcome
294	Zaridze D. Alcohol and health in Russia: good news at last. <i>Lancet</i> . 2017. 390(10102):1616-1618. doi:10.1016/s0140-6736(17)32387-5.	Study Design; Publication Status; Other (e.g., duplicative data, not primary research)
295	Zazpe I, Sanchez-Tainta A, Toledo E, Sanchez-Villegas A, Martinez-Gonzalez MA. Dietary patterns and total mortality in a Mediterranean cohort: the SUN project. <i>J Acad Nutr Diet</i> . 2014. 114(1):37-47. doi:10.1016/j.jand.2013.07.024.	Intervention/Exposure
296	Zhang QL, Zhao LG, Zhang W, Li HL, Gao J, Han LH, Zheng W, Shu XO, Xiang YB. Combined impact of known lifestyle factors on total and cause-specific mortality among Chinese men: a prospective cohort study. <i>Sci Rep</i> . 2017. 7(1):5293. doi:10.1038/s41598-017-05079-5.	Country
297	Zhao G, Ronda E, Cea L, Pulido J, Barrio G, Regidor E. Mortality by cause of death and risk behaviors in farmers versus non-farmers: the importance of avoiding the healthy worker effect. <i>Int Arch Occup Environ Health</i> . 2019. 92(4):599-608. doi:10.1007/s00420-018-1396-2.	Intervention/Exposure
298	Zhao W, Ukawa S, Okada E, Wakai K, Kawamura T, Ando M, Tamakoshi A. The associations of dietary patterns with all-cause mortality and other lifestyle factors in the elderly: An age-specific prospective cohort study. <i>Clin Nutr</i> . 2019. 38(1):288-296. doi:10.1016/j.clnu.2018.01.018.	Intervention/Exposure
299	Zhu N, Yu C, Guo Y, Bian Z, Han Y, Yang L, Chen Y, Du H, Li H, Liu F, Chen J, Chen Z, Lv J, Li L. Adherence to a healthy lifestyle and all-cause and cause-specific mortality in Chinese adults: a 10-year prospective study of 0.5 million people. <i>Int J Behav Nutr Phys Act</i> . 2019. 16(1):98. doi:10.1186/s12966-019-0860-z.	Country